



**RECONNAISSANCE INVENTORY AND PRIORITIZATION
OF EXISTING AND POTENTIAL BOTTOMLANDS
IN THE UPPER COLORADO RIVER BASIN
1993-1994**



**Final Report
July 1995**

U. S. FISH AND WILDLIFE SERVICE



Digitized by the Internet Archive
in 2012 with funding from
LYRASIS Members and Sloan Foundation

<http://archive.org/details/reconnaissancein00irvi>

**RECONNAISSANCE INVENTORY AND PRIORITIZATION
OF EXISTING AND POTENTIAL BOTTOMLANDS
IN THE UPPER COLORADO RIVER BASIN
1993-1994**

Final Report

July 1995

**David B. Irving^a
Fishery Biologist**

and

**Bob D. Burdick^b
Fishery Biologist**


U. S. FISH AND WILDLIFE SERVICE

^a Colorado River Fishery Project, 266 West 100 North, Suite 2, Vernal, Utah 84078

^b Colorado River Fishery Project, 764 Horizon Drive, South Annex A, Grand Junction, Colorado 81506

This project was funded by the Recovery Implementation Program for Endangered Fish Species in the Upper Colorado River Basin. This Program is a joint effort of the U. S. Fish and Wildlife Service, U. S. Bureau of Reclamation, Western Area Power Administration, States of Colorado, Utah, and Wyoming, Upper Basin water users, and environmental organizations.

Recovery Implementation Program for the Endangered Fishes of the Upper Colorado River Basin, U. S. Fish and Wildlife Service, P. O. Box 25486, Denver Federal Center, Denver, Colorado 80225



DISCLAIMER

Mention of trade names, commercial products, or firms and businesses does not constitute endorsement or recommendation for use by the authors, the Fish and Wildlife Service, U. S. Department of Interior, or the Recovery Implementation Program.

TABLE OF CONTENTS

LIST OF TABLES	v
LIST OF FIGURES	v
LIST OF APPENDICES	viii
ACKNOWLEDGMENTS	xi
EXECUTIVE SUMMARY	xiii
INTRODUCTION	1
Background	1
Purpose and Objectives	2
STUDY AREA	3
METHODS	3
Inventory	3
Identification of Bottomlands	3
Classification of Bottomlands	6
Scoring and Prioritization of Sites	7
Geomorphology	10
Ecology	12
Hydrology	12
Stage vs. Discharge Relationship	12
Channel Cross Sections	13
River Discharge	13
Land Ownership Status	14
RESULTS AND DISCUSSION	14
Photographic Data	14
Water Discharge on Photograph Dates	15
Distribution of Bottomland Habitats	15
Green River	15
Yampa River	15
White River	17
Colorado River	17
Site Scoring and Prioritization	20
Green River Drainage	21
Colorado River Drainage	23
Ecology	25
Hydrology	25
Green River Drainage	25
Stage vs. Discharge Relationship	25
Channel Cross Sections	34
Colorado River Drainage	36
Stage vs. Discharge Relationship	36
Channel Cross Sections	45

TABLE OF CONTENTS (cont'd)

Land Ownership Status	47
Green River Drainage	47
Colorado River Drainage	49
RECOMMENDATIONS	52
LITERATURE CITED	55

LIST OF APPENDICES

Appendix A--The Floodplain Habitat Restoration Program; Commonly Asked Questions; Right-of-Entry Permit Form	
Appendix B--Existing and needed inventory maps and photos	
Appendix C--Complete and partial inventory plan for the Green and Colorado rivers, 1993	
Appendix D--Bottomland habitat sites, river miles, historical and actual flooded acreages	
Appendix E--Parameters used to categorize and rank floodplain restoration sites	
Appendix F--Scoring and ranking of bottomland sites	
Appendix G--Land ownership acreage for the Colorado and Gunnison rivers	
Appendix H--Nine initial criteria and sub-criteria considered for scoring and ranking bottomland sites	
Appendix I--Mean daily flow exceedences	

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1. Paired comparison analysis of the four selection criteria. . . .	9
2. The top 11 ranked bottomland sites adjacent to the Green (GR) River, 1993	21
3. The top 13 ranked bottomland sites adjacent to the Colorado (CO) and Gunnison (GU) rivers, 1993	23
4. Brief description of the five bottomland sites selected for demonstration projects in the Colorado and Green River drainages, Colorado and Utah	26
5. Range of flows and dates that stage-discharge data were collected in the Green River drainage, 1994	34
6. Estimated overbank flooding and frequency of flooding at the seven bottomland habitat sites in the Green River (GR) . . .	35
7. Range of flows and dates that stage-discharge data were collected in the Colorado and Gunnison river drainages, 1994	36
8. Estimated overbank flooding and frequency of flooding at the nine bottomland habitat sites in the Colorado and Gunnison rivers.	46

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1. Bottomland habitat inventory of the Upper Colorado River Basin, Colorado and Utah, 1993	4
2. Schematic of the bed profile of the three major bottomland habitat classifications at various flow regimes in the Upper Colorado River Basin	11
3. Water discharge (cfs) at peak and low-flow periods on the Green, Yampa, White, Colorado, and Gunnison rivers, Colorado and Utah, 1992 and 1993	16
4. Actual and potential (historic floodplain) acres (hectares) of bottomland habitat adjacent to the Green River, Colorado and Utah, 1993	18

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
5. Areas inundated during high water (May and June) and low water (September) in 1993 compared with areas that historically flooded on 240 miles of the Colorado River from the Green River confluence upstream to Rifle, Colorado and 75 miles of the Gunnison River from the Colorado River confluence upstream to the North Fork confluence	19
6. Actual and potential (historic floodplain) acres (hectares) of the top eleven ranked bottomland habitat sites adjacent to the Green River, Utah, 1993	22
7. Areas inundated during high water (May and June) and low water (September) in 1993 compared with areas that were historically flooded for the top 13 ranked bottomland sites adjacent to the Colorado and Gunnison rivers	24
8. River profile and stage versus discharge relationship for the Meril Snow Ranch bottomland habitat site adjacent to the Green River, Utah, 1994.	27
9. River profile and stage versus discharge relationship for the Little Stewart Lake bottomland habitat site adjacent to the Green River, Utah, 1994.	28
10. River profile and stage versus discharge relationship for the Baeser Bend bottomland habitat site adjacent to the Green River, Utah, 1994	29
11. River profile and stage versus discharge relationship for the Johnson Bottom bottomland habitat site adjacent to the Green River, Utah, 1994	30
12. River profile and stage versus discharge relationship for the Leota Pond 7 bottomland habitat site adjacent to the Green River, Utah, 1994	31
13. River profile and stage versus discharge relationship for the Sheppard Bottom bottomland habitat site adjacent to the Green River, Utah, 1994	32
14. River profile and stage versus discharge relationship for the Woods Bottom (Old Charlie Wash) bottomland habitat site adjacent to the Green River, Utah, 1994	33

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
15. River profile (top) and stage vs. discharge relationship (bottom) for Adobe Creek bottomland habitat site adjacent to the Colorado River, river mile 159.9	37
16. River profile (top) and stage vs. discharge relationship (bottom) for 30-29 Road bottomland habitat site adjacent to the Colorado River, river mile 174.4	38
17. River profile (top) and stage vs. discharge relationship (bottom) for Griffith's bottomland habitat site adjacent to the Colorado River, river mile 176.0	39
18. River profile (top) and stage vs. discharge relationship (bottom) for Clifton Sanitation No. 1 bottomland habitat site adjacent to the Colorado River, river mile 178.5.	40
19. River profile (top) and stage vs. discharge relationship (bottom) for EXXON bottomland habitat site adjacent to the Colorado River, river mile 220.0	41
20. River profile (top) and stage vs. discharge relationship (bottom) for Battlement Mesa bottomland habitat site adjacent to the Colorado River, river mile 221.3	42
21. River profile (top) and stage vs. discharge relationship (bottom) for Johnson Boy's Slough bottomland habitat site adjacent to the Gunnison River, river mile 53.6.	43
22. River profile (top) and stage vs. discharge relationship (bottom) for Confluence Park bottomland habitat site adjacent to the Gunnison River, river mile 57.1	44
23. River profile for Pike's bottomland habitat site adjacent to the Colorado River, river mile 179.1	45
24. Land ownership status in the Browns Park National Wildlife Refuge bottomland habitat area (rm 368-393), Escalante Ranch to Brennan Bottom area (rm 262-310), Ouray National Wildlife Refuge area (rm 249-265), Ouray to Pariette Draw area (rm 238-251), and Desert Spring to Green and Colorado river confluence area (rm 0-249), Green River, Colorado and Utah, 1993	48
25. Land ownership status of bottomland sites inventoried in five stream reaches adjacent to the Colorado River, Colorado and Utah, extending from Rifle, Colorado (river mile 241) downstream to the Green River confluence (river mile 0).	50

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
26. Land ownership status of bottomland sites inventoried in three stream reaches adjacent to the Gunnison River, Colorado extending from the North Fork confluence (river mile 75) downstream to the Colorado River confluence (river mile 0.7).	51

LIST OF APPENDICES

<u>Appendix</u>	<u>Page</u>
A. Floodplain Habitat Restoration Program; Commonly asked questions; Right-of-entry permit form.	A-1-A-4
B. Table B.1. Existing and needed inventory maps, aerial photographs, infrared photography, and videos of the Green River Basin, Colorado and Utah, 1993	B-1
C. Table C.1. Plan for complete or partial infrared photography inventories for the Green River Basin, Colorado and Utah, 1993.	C-1
D. Table D.1. Bottomland habitat sites, river mile, and actual and potential inundated acres (hectares), Green River, Colorado and Utah, May and September, 1993.	D-1-D-2
D. Table D.2. Bottomland habitat sites, river mile, actual (May [runoff] and September [post-runoff] 1993) inundated area (hectares), and historical inundated area (hectares), Colorado and Gunnison rivers, Colorado and Utah, May and September 1993	D-3-D-6
E. Table E.1. The 10 parameters from 35-mm color slide photo-imagery used to categorize and rank bottomland habitats important to endangered fishes in the Green River, Colorado and Utah, 1993 (Part 1)	E-1,E-3,E-5,E-7
E. Table E.2. The 10 parameters from 35-mm color slide photo-imagery used to categorize and rank bottomland habitats important to endangered fishes in the Green River, Colorado and Utah, 1993 (Part 2).	E-2,E-4,E-6,E-8
E. Table E.3. The 10 parameters from 35-mm color slide photo-imagery used to categorize and rank bottomland habitats important to endangered fishes in the Yampa River, Colorado, 1993 (Part 1)	E-9

LIST OF APPENDICES

<u>Appendix</u>	<u>Page</u>
E. Table E.4. The 10 parameters from 35-mm color slide photo-imagery used to categorize and rank bottomland habitats important to endangered fishes in the Yampa River, Colorado, 1993 (Part 2)	E-9
E. Table E.5. The 10 parameters from 35-mm color slide photo-imagery used to categorize and rank bottomland habitats important to endangered fishes in the White River, Colorado and Utah, 1993 (Part 1).	E-10
E. Table E.6. The 10 parameters from 35-mm color slide photo-imagery used to categorize and rank bottomland habitats important to endangered fishes in the White River, Colorado and Utah, 1993 (Part 2)	E-10
E. Table E.7-E.16. The 10 parameters used to rank 158 potential floodplain restoration sites important to endangered fishes of the Upper Colorado River Basin. . . .	E-11-E-30
F. Table F.1. Selection criteria and associated sub-criteria raw score values multiplied by the average weighted point values to produce the total score and rank for each bottomland habitat site in the Green, Yampa, and White rivers, Colorado and Utah, 1993	F-1-F-3
F. Table F.2. Selection criteria and associated sub-criteria raw score values multiplied by the average weighted point values to produce the total score and rank for each bottomland sites in the Colorado and Gunnison rivers, Colorado and Utah, 1993	F-4-F-10
G. Table G.1. Land ownership status by area (hectares) and percentage for 158 bottomland sites adjacent to the Colorado and Gunnison rivers	G-1
H. Table H.1. The initial nine criteria and sub-criteria considered for selection criteria for scoring and ranking bottomland sites in the Upper Colorado River Basin	H-1
I. Figure I.1. Percent of time flow (cfs) is exceeded in the Green River before (1947-1966) and after (1966-1993) Flaming Gorge Dam	I-1

LIST OF APPENDICES

<u>Appendix</u>	<u>Page</u>
I. Figure I.2. Mean daily flow exceedences for May (top) and June (bottom) for the Colorado River at the Stateline USGS stream gauge during the post-water development period, 1952-1993	I-2
I. Figure I.3. Mean daily flow exceedences for May (top) and June (bottom) for the Colorado River at the Cameo USGS stream gauge during the post-water development period, 1952-1993	I-3
I. Figure I.4. Mean daily flow exceedences for May (top) and June (bottom) for Plateau Creek (near the confluence with the Colorado River) USGS stream gauge during the post-water development period, 1952-1993	I-4
I. Figure I.5. Mean daily flow exceedences for May (top) and June (bottom) for the Gunnison River at the Whitewater USGS stream gauge during the post-water development period, 1952-1993	I-5
I. Figure I.6. Mean daily flow exceedences for May (top) and June (bottom) for the Gunnison River at the Delta USGS stream gauge during the post-water development period, 1952-1993	I-6

ACKNOWLEDGMENTS

We would like to thank the present and former Coordinator of the Floodplain Habitat Restoration Program from the U.S. Fish and Wildlife Service, Pat Nelson and Dr. Richard Wydoski, respectively, and John Hawkins, Larval Fish Laboratory, Colorado State University, Ft. Collins, for their guidance and assistance in developing appropriate research techniques and data evaluation. George Smith, Regional Hydrologist of the Fish and Wildlife Service, Denver, Colorado, and his staff provided hydrology support. Without support from key personnel in the Bureau of Reclamation, the hydrology portion of the study could not have been accomplished. Chuck Gallagher was instrumental in designing and establishing the field-based camera-recording equipment to obtain staff-discharge information. Terry Ayer, Blake Bailey, Burt Ives, Bob Levine, and Ted Juntilla from the Grand Junction, Colorado, Bureau of Reclamation Projects Office and Tom Jordan, Donavan Johnston and S. Lujan from the Provo, Utah, Office of the Bureau of Reclamation surveyed river cross sections. Pat Mangan and Andrew Montano from the Environmental Planning and Assessment Branch of the Bureau of Reclamation in Denver assisted with river cross sections. Bob Norman, Bureau of Reclamation in Grand Junction, and Tim Modde and Frank Pfeifer, Fish and Wildlife Service, provided administrative support. Both Bob and Jeff Baumgartner, The Nature Conservancy, often offered useful technical advice to the project. Gordon Mueller, Richard Wydoski, and Robert Muth provided useful comments and improved an earlier version of this report.

We are grateful for the field work, assistance, and advice provided by Steve Brock, Dan Schaad, and Kelli Stone from the Ouray National Wildlife Refuge, and Tom Hatch and Gary Baker from the Colorado River Fishery Project (CRFP) in Vernal, Utah. We would like to thank CRFP biological technicians, John Rosendale, Sara Persselin, Chris Flann, Quentin Bradwich, and Bruce Bonar for their assistance in collecting field data. Mike Baker, Mike Montagne, and Greg Taylor assembled staff gauges and fabricated ammo boxes for housing cameras for the hydrology portion of the study.

A special thanks goes to Greg Whitbeck and Mary Smuin from the U. S. Natural Resources Conservation Service, Vernal, Utah for the use of their camera equipment and photography skills. Carl Landini and his staff from the Consolidated Farm Service Agency in Grand Junction provided office space and digitizing equipment to obtain acreage estimates of bottomland habitat areas.

Dave Webster from the Farm Service Agency and Doyle Dow, real estate appraiser from the Bureau of Reclamation, were pivotal in working with local landowners in the Vernal, Utah, and Grand Junction and Delta, Colorado localities. We would also like to thank Randy Arment of Research Air, Inc. and Hank Heines of Dinaland Aviation for their skill and safety in assisting in the aerial reconnaissance survey work.

Aerial photographs of the Colorado and Gunnison rivers were provided by Grady McNure and Ken Jacobson of the U.S. Army Corps of Engineers Regulatory

Office, Grand Junction, Dr. Jack Stanford, University of Montana, and Paul Obert, U. S. Natural Resource Conservation Service, Delta. Ms. Judy Armstrong made historical photos available from the archive section at the Museum of Western Colorado.

Finally, we would like to thank each of the landowners for allowing us access to or through their land to each of the bottomland sites to conduct the on-ground hydrology screening--without their voluntary willingness to participate, much of the on-site hydrology and screening work could not have been accomplished. Those landowners include Mr. Oby Blanchard; Mr. Don Sheley, President of the Board, Clifton Sanitation No. 1; Mr. Frank Hyde and Mr. Brian Woods, Clifton Sanitation No. 2; Mr. L.O. Griffith; Mr. Kurt Mill, Colorado Division of Parks and Outdoor Recreation; Mr. Ed Gardner, Whitewater Building Materials; Mr. Harold Pike; EXXON Corporation, Houston, Texas; Mr. Bill Wilde, Battlement Mesa Partners; Mr. Wendell Johnson; Mr. Kenneth Johnson; Mr. Fred Wild; Mr. Ray Toole; and representatives from the City of Delta.

EXECUTIVE SUMMARY

The purpose of this study was to inventory all potential flooded bottomland habitats in the Upper Colorado River Basin that were believed to be important for recovery of endangered fishes, especially razorback sucker, and rank (prioritize) the bottomland sites. All potential bottomland habitats adjacent to stream reaches in the Upper Colorado River Basin (871 river miles) that were considered important for recovery of endangered fishes were inventoried in 1993. A total of 293 bottomland habitat sites in the Green River (135 sites) and Colorado River (158) drainages was identified, categorized, and classified using a five-step procedure to objectively prioritize bottomland sites. Each site was scored and prioritized (ranked) from four selection criteria (land ownership, proximity to a known or planned razorback sucker spawning area or proximity to recent adult captures, June hydrological connection to the river, and potential for a network [complex] of bottomland sites) that were based on available photographic imagery and the most current biological information. Prioritizing the bottomland sites was also used to establish staff gauge and river transect locations in 1994 to determine flows necessary to inundate a specific bottomland site.

In the Green River drainage, the largest flooded bottomland areas were located in the Browns Park National Wildlife Refuge reach (1,371 acres [555 ha]) and between Escalante Ranch and Pariette Draw (6,093 acres [2,466 ha]). The greatest concentration of flooded bottomland sites were within the Ouray National Wildlife Refuge (2,265 acres [917 ha]). In the Colorado River drainage, there were three distinct areas where bottomland habitats were concentrated. These were usually in large floodplain areas near agricultural centers. The first area was the stream reach between Rifle and Debeque, Colorado (2,248 acres [910 ha]), the second was the stream reach in the Grand Valley between Palisade and Loma, Colorado (4,564 acres [1,847 ha]), and the third area between the Cisco Boat Landing and McGraw and Hotel Bottoms (1,214 acres [491 ha]) upstream from Moab, Utah. Moab Slough, near Moab, Utah, was the largest, single bottomland area along the Colorado River encompassing approximately 875 acres (354 ha). In the Gunnison River drainage, the largest and only major floodplain area was between Austin, Colorado, and Roubideau Creek (1,739 acres [704 ha]; 17 river miles [27 rkm]).

The average inundated area during both the May and June runoff and the low-flow period in September was larger in the Green River (May and June: mean = 59 acres [24 ha]; September: mean = 19 acres [8 ha]) than any of the sites inundated in both the Colorado River (May and June: mean = 22 acres [9 ha]; September: mean = 10 acres [4 ha]) or Gunnison River (May and June: mean = 17 acres [7 ha]; September: mean = 3 acres [1 ha]).

In the Green River, the bottomland habitat sites between Pariette Draw and Escalante Ranch (32 sites) were ranked 1 through 11. Bottomland habitat in the Ouray National Wildlife Refuge (six sites) were the highest ranked (1st and 2nd) sites. The Browns Park National Wildlife Refuge (eight sites) sites ranked (12th-14th) the next highest, followed by lower ranked (17th-18th) sites from Pariette Draw to the Green and Colorado river confluence (92 sites). Based on these rankings, there were five general bottomland habitat

areas in the Green River. The top ranked sites were in the Ouray National Wildlife Refuge, the next in the Jensen to Brennan Bottom area, followed by sites downstream of Ouray. The lowest ranked sites were in the Browns Park National Wildlife Refuge area and from Desert Spring and Moon Bottom to the Green and Colorado river confluence. A vast majority (75%) of the bottomland habitat types in the Green River sub-basin were floodplain terraces. Federal agencies own most (75%) of the bottomlands in the vicinity of Browns Park National Wildlife Refuge. Forty-two percent of the land between Escalante Ranch and Brennan Bottom was in private ownership. All of the Ouray National Wildlife Refuge bottomland sites are federally owned. Seventy-one percent of the bottomland sites were on tribal lands between Ouray and Pariette Draw. The majority of bottomland habitat between Desert Spring and the Green and Colorado river confluence is federal land.

Unlike the Green River drainage, the top 13 ranked sites in the Colorado and Gunnison river drainages were scattered among four general areas. Along the Colorado River, there were 10 bottomland habitat sites within the Debeque area, eight sites within the 15-mile reach, and five sites within the 18-mile reach. There were three sites within the vicinity of Delta, Colorado, along the Gunnison River.

The top ranked site was the Walter Walker State Wildlife Area in the 18-mile reach. Four other sites in the immediate vicinity ranked 6th and 10th. Johnson Boy's Slough along the Gunnison River ranked 2nd, the Clifton Pond area along the Colorado River ranked 3rd, and the Debeque I-70 Slough ranked 4th. Six other sites immediately downstream of the Debeque I-70 Slough ranked 9th and 11th. Floodplain terraces comprised 37% of the bottomland habitat types, gravel-pit ponds 21%, and side channels 20%. Of the 110 bottomland sites identified and inventoried along the 241 miles of the Colorado River, 51% or about 6,355 acres (2,572 ha) were in private ownership. Of the 48 bottomland sites along 75 miles of the Gunnison River, 70% or about 2,221 acres (899 ha) were in private ownership. Fifteen percent (1,826 acres [739 ha]) of the ownership was in public (state and federal) ownership along the Colorado River; 22% (697 acres [282 ha]) was in public (state and federal) ownership along the Gunnison River. The greatest amount of bottomland habitat along the Colorado River owned by the federal government (83%) was within Canyonlands National Park. The federal government does not own any bottomlands between Rifle and Loma. The federal government owned or controlled only 8 percent of the bottomland habitat along the Gunnison River within a canyon-bound area between Roubideau Creek (rm 50 [rkm 81]) and the Colorado River confluence.

Hydrological parameters such as the annual frequency, magnitude, and duration of flooding at a site provide pivotal information that can be used to determine if a site is suitable for enhancement or restoration. This information could be used as an additional selection criterion to screen sites.

In 1994, channel cross sections were established where temporary staff gauges had been previously placed at seven bottomland sites adjacent to the Green River and at nine sites adjacent to the Colorado and Gunnison rivers to collect flow and elevation data from May through July. Stage vs. discharge

relationships for each site were determined to estimate a range of flows necessary to inundate the bottomland site. However, the flows that were estimated to flood each of these bottomland sites should be considered provisional in this report until empirical verification can be obtained for these sites. Flow information was also used to determine the frequency that predicted flows might now occur at each bottomland site.

In the Green River drainage, of the seven sites evaluated, the Woods Bottom (Old Charlie Wash) bottomland site within the Ouray National Wildlife Refuge floods the most frequently, between 95-100% (i. e., between 9-1/2 and 10 years out of 10). Sheppard and Johnson Bottoms flood approximately 65-75% and 40-45% annually, respectively. The Baeser Bend site floods less frequently, only about 10-20% (one to two years out of 10). In the Colorado River drainage, of the nine sites evaluated, the Griffith bottomland site in the 15-mile reach of the Colorado River floods the most frequently, about 50% (one year out of two). The frequency of flooding at the Confluence Park bottomland site (an oxbow) on the Gunnison River now occurs only about 10-15% (one to one and one-half years out of 10) since the construction of upstream dams; a tertiary side channel at the Adobe Creek site is inundated about 20-25% (two to two and one-half years out of 10). Many of the other sites are now inundated less frequently because of the loss of the (historical) high-volume years, which are important to inundating off-channel habitats in the floodplain. Not only has the magnitude of runoff flows in May and June been reduced by about 45% in both the Colorado and Gunnison rivers, and by 22% in the Green River but the duration of flooding has also been reduced. Much of this is a result of upstream water development projects, primarily transmountain diversions and reservoir storage. The construction of levees and dikes, particularly along the Colorado and Gunnison rivers, has further reduced the likelihood of flooding by further isolating the floodplain from the main stream reaches and preventing water access during runoff. Given the present hydrologic regime, current water availability, and future compact water development in the Upper Colorado River Basin, inundating bottomland sites might be best accomplished by removing portions of- or entire human-made dikes or natural obstructions (e.g., levees) to increase flooding frequency. Those sites that flood more frequently under the existing hydrologic regime will also flood more frequently in the future, and will provide more benefits to endangered fish.

The priority recommendation was to collect additional flow and elevation data during runoff in 1995 at seven bottomland sites established in 1994 on the Green River and at four bottomland sites (Pike's and Adobe Creek on the Colorado River and Confluence Park and Johnson Boy's Slough on the Gunnison River) established in 1994 to verify and refine stage vs. discharge relationships. These data are particularly important for estimating flows necessary to inundate bottomland sites and vital in determining which sites can be practically restored. Other recommendations include, 1) completing contaminant screening for the remaining 19 bottomland sites investigated in 1994, 2) determining if three bottomland depression sites along the Colorado River between Moab, Utah, and the Green River confluence have suitable habitat for restoration, 3) obtaining a complete set of current color aerial photographs of the entire Upper Colorado River Basin, and 4) developing alternative or additional selection criteria, if warranted by site visits, to

re-prioritize bottomland sites identified in this inventory.

INTRODUCTION

Background

Historically, Upper Colorado River Basin floodplains were inundated more frequently by flows during spring runoff, but today floodplains are not regularly connected to the river because of channelization by either levees and dikes or rip-rap near population centers in agricultural areas. Introduced salt cedar or tamarisk (*Tamarisk spp.*) has become established along the shoreline of major rivers resulting in sediment deposition and stabilized banks that further reduce the connectivity of the river with historic flooded bottomlands (Graf 1978). The periodicity of out-of-channel flooding in the Upper Colorado River has dramatically decreased following the onset of transmountain water diversions, irrigation diversions, and the construction of mainstem dams (Osmundson and Kaeding 1991). The construction of dikes coupled with the reduction of high spring flows has altered the natural hydrograph and either reduced or eliminated regular flooding of bottomlands. It is believed that flooded bottomlands may have served as nursery areas for the razorback sucker *Xyrauchen texanus* (McAda 1977; Osmundson and Kaeding 1991). In addition, Osmundson and Kaeding (1991) suggested that oxbow lakes and flooded pastures in the Grand Valley near Grand Junction, Colorado, were historically the primary spawning habitats of the razorback sucker. They also suggested that low-velocity, off-channel habitats were used by adult Colorado squawfish (*Ptychocheilus lucius*) during high stream flows.

Numerous studies (Grabowski and Hiebert 1989; Tyus and Karp 1989; Wydoski and Wick 1994) have suggested the importance of seasonal flooding to river productivity. Flooded bottomland habitats were considered important to adult razorback sucker for feeding prior to and after spawning and also important as nursery areas (Tyus and Karp 1989). Adult fish may have used these off-channel habitats for "velocity shelters" to escape high water velocities during spring runoff.

Turbid rivers in the upper basin are not very productive for zooplankton that are essential food for early life stages of razorback sucker. When compared to the riverine environment and river backwaters, inundated bottomland habitats produce the higher densities of zooplankton (Grabowski and Hiebert 1989; Mabey and Shiozawa 1993). Although predation by non-native fishes has been documented to be a limiting factor in survival of larval razorback sucker in the lower basin (Minckley et al. 1991), starvation may also limit survival (Marsh and Langhorst 1988; Papoulias and Minckley 1990). It is hypothesized that the loss of these productive flooded bottomland habitats is limiting recruitment of razorback sucker in the Upper Colorado River Basin because of insufficient quantity of appropriate-sized food organisms at the right time (Wydoski and Wick 1994).

Most flooded bottomland habitats occur in broad valleys along low-gradient stream reaches. Bottomlands are off- or out-of-channel habitats that include oxbow lakes, former side channels in broad valley floodplains, ponds, and wetland depressions and terraces. Less numerous flooded bottomland habitats are found in canyon-bound stream reaches. These are wetland

depressions that are currently not connected to the river and receive water from groundwater infiltration, wetland terraces that are only inundated during flooded periods, or canyon mouths that are "ribbon-like" channels in narrow canyons adjacent to the main river flow. During high flow events, some of these off-channel features are temporarily connected to the river. Re-establishing connectivity to the mainstem river may provide off-channel habitats for endangered fishes that are essential for effective management of the riverine ecosystem and recovery of endangered fishes. Former natural riverine features could be integrated into the historic floodplain by removing portions of human-made dikes or natural obstructions (e.g., gravel and sand bars).

Habitat development and enhancement is identified as one of the five principal elements of the Recovery Program (U. S. Fish and Wildlife Service [FWS] 1987). Protection, restoration, and enhancement of inundated bottomland habitat along mainstem riparian corridors are believed to be important for recovery of razorback sucker. Riparian enhancement in the Upper Colorado River Basin can be accomplished by providing sufficient flows to inundate bottomlands in a manner that approximates the natural hydrograph. If sufficient flows cannot be obtained regularly, dikes and levees should be breached at intervals to allow inundation of lowlands during high spring flows. Restoration of flooded bottomland habitats will provide food that should increase survival of larvae and juveniles of endangered fishes as well as provide resting and feeding areas for adults. Before razorback sucker are reintroduced into historical stream reaches of the Upper Colorado and Gunnison rivers, a habitat enhancement program (i. e., flooded bottomland management) must be implemented if recovery is to be realized.

Purpose and Objectives

The purpose of this study was to summarize all potential flooded bottomland habitats in the Upper Colorado River Basin that were believed to be important for recovery of endangered fishes, especially razorback sucker. The inventory of these habitats provides managers with candidate sites for acquisition, restoration, and management. The objectives of the study were to:

1. Identify bottomlands adjacent to Upper Colorado River Basin mainstem rivers representing potential endangered fish habitat.
2. Categorize and classify the bottomlands identified according to potential value for endangered fish recovery.
3. Rank (prioritize) mainstem bottomlands by their perceived value to endangered fish recovery.

The end product is an inventory of mainstem flooded bottomlands categorized by size (area), extent of flooding, fish and water access, upland vegetation types, ownership, and related criteria. Sites were ranked as to potential contribution to endangered fish recovery.

Another objective was to conduct preliminary hydrology screening at a few (up to 20 sites) bottomland sites that ranked highest from the prioritization. We believed that predicting flows necessary for overbank flooding to inundate a bottomland site was necessary for determining sites that could be practically restored or enhanced.

STUDY AREA

The study area included the mainstem river channels and adjacent bottomland habitats in the Upper Colorado River Basin (Figure 1). River miles [(rm); river kilometers (rkm)] inventoried were 398 rm (640.5 rkm) in the Green River, 50 rm (80.5 rkm) in the Yampa River, 104.5 rm (168.2 rkm) in the White River, 241 rm (387.9 rkm) in the Colorado River, and 75 rm (120.7 rkm) in the Gunnison River.

METHODS

Inventory

A two-step approach was used to accomplish the bottomland inventory. First, potential bottomland habitats adjacent to mainstem rivers were identified and inventoried in 1993. Inventorying sites included cataloging various information from different photographic imagery. Second, each bottomland habitat site was classified using a numerical score from four selection criteria, and prioritized in 1994.

Identification of Bottomlands

All existing information necessary to conduct this aspect of the study was identified and assembled. This included topographical maps (7-1/2 minute quadrangle maps), land status maps, and various aerial photographic imagery that included videography, color infrared photographs, and 35-mm colored slides. Information sources included the U. S. Army Corps of Engineers, U. S. Bureau of Reclamation (BR), U. S. Bureau of Land Management, U. S. Natural Resources Conservation Service (NRCS), Argonne National Laboratories, and FWS.

A Cessna 206 fixed-winged aircraft was used to conduct low-level reconnaissance photography flights over bottomland habitats along the Green and Colorado rivers in 1993. The NRCS provided equipment and assistance during each photography session. A 35-mm camera loaded with color slide film was attached to the underside of the plane to facilitate direct overhead photography of bottomland sites. The plane was flown at an altitude of 5,000 to 6,000 feet (above ground level) directly overhead to allow for photo-image coverage of the main river channel and adjacent floodplain areas.

Aerial photographs were taken along the Green, Yampa, and White rivers in the Green River drainage and the Colorado and Gunnison rivers in the Colorado River drainage during 1993. The beginning and ending locations for flights in these river drainages were as follows:



Figure 1. Bottomland habitat inventory of the Upper Colorado River Basin, Colorado and Utah, 1993.

1. Green River drainage

- a. Green River from Browns Park to the Green and Colorado river confluence (rm 0-398 [rkm 0-640.5]).
- b. Yampa River from Cross Mountain to the Yampa and Green river confluence (rm 0-50 [rkm 0-80.5]).
- c. White River from Taylor Draw Dam to the White and Green river confluence (rm 0-104.5 [rkm 0-168.2]).

2. Colorado River drainage

- a. Colorado River from Rifle, Colorado to the Colorado and Green river confluence (rm 0-241 [rkm 0-387.9]).
- b. Gunnison River from the Gunnison and North Fork river confluence to the Gunnison and Colorado river confluence (rm 0-75, [rkm 0-120.7]).

Aerial photographs were taken during high, medium, and low flows in the Green River drainage and during high and low flows in the Colorado River drainage. Flows were determined using the U.S. Geological Survey (USGS) stream gauge on the Green River at Jensen, Utah, the Stateline stream gauge on the Colorado River near the Utah and Colorado stateline, and the Gunnison River stream gauge at Whitewater, Colorado. The target water discharge (cubic feet per second [cfs]) and date for aerial photographs were as follows:

1. Green River drainage

- a. High flows between 18,000 and 20,000 cfs on 25 May 1993 in the Green, Yampa, and White rivers.
- b. Medium flows between 6,000 and 12,000 cfs on 15 and 30 June 1993 in the Green River between Split Mountain and Sand Wash.
- c. Low flows between 1,000 and 2,000 cfs on 28 September 1993 in the Green, Yampa, and White rivers.

2. Colorado River drainage

- a. High flows between 18,000 and 20,000 cfs in the Colorado River and 7,000 and 9,000 cfs in the Gunnison River on 10 June 1993.
- b. Low flows between 4,000 and 5,000 cfs in the Colorado River and 2,000 and 3,000 cfs in the Gunnison River on 28 September 1993.

After consultation with personnel from the NRCS, it was determined that one photograph per river mile was necessary to adequately photo-document bottomland habitat areas. It was estimated that it would take approximately

23-28 rolls¹ of color slide film to photograph the 871 miles of river bottomland areas delineated in the Green, Yampa, White, Colorado, and Gunnison rivers.

A list identifying the location of each bottomland habitat site was formulated from the aerial 35-mm color slides and color infrared photographs. All slides shot per photo session were meticulously cataloged according to film roll number and slide sequence number, date, river, river mile, and river location, and then stored in slide holders and binders for future reference.

Once the color slides were organized by river and river mile, they were viewed to identify and locate specific bottomland areas of potential endangered fish habitat. The bottomland areas on these slides were then computer digitized or overlaid with a transparent area grid to estimate flooded acreage. Estimates of area were within $\pm 10\%$.

i-----

$$Rolls = \left[\left(\frac{1 Photo}{Riv.Mi.} \right) \times \left(\frac{871 Riv.Mi.}{Drainage} \right) \times \left(\frac{1 Roll}{36 Photos} \right) \right] \approx 24 Rolls (36 Exposure)$$

Classification of Bottomlands

Ten categories of information were used in cataloging, categorizing, and prioritizing bottomland habitats in the Upper Colorado River Basin. These parameters included (see Appendix E):

1. Basin, sub-basin, river
2. Bottomland Location: site, river mile, township, range, section
3. Runoff (May and June)
 - a. Total potential (historical) inundated area in acres and hectares; the total area that could be potentially flooded during historic or the pre-water development era
Note: the 100-year floodplain was used to estimate the historical inundated area
 - b. Inundated (yes/no)
 - c. Total inundated area (acres/hectares); the actual or realistic area flooded
 - d. Hydrological connection to mainstem river
 - 1) yes/no
 - 2) number of connections
 - 3) connections up- or downstream
 - 4) other water sources (groundwater, gravity feed, tributary/stream)
4. Post-runoff (September)
 - a. Total potential (historical) inundated area in acres and hectares
Note: the 100-year floodplain was used to estimate the

- historical inundated area
- b. Inundated (yes/no)
- c. Total inundated area (acres/hectares)
- d. Hydrological connection to mainstem river
 - 1) yes/no
 - 2) number of connections
 - 3) connections up- or downstream
 - 4) other water sources (groundwater, gravity feed, tributary/stream)
- 5. Levy/Berm
 - a. Levy/berm dividing mainstem and bottomland
 - 1) yes/no
 - 2) human-made or natural
 - b. Levy width of length (meters)
- 6. Vegetation (upland)
 - a. Yes/no
 - b. Predominant type(s)
- 7. Proximity to Endangered Fishes
 - a. River mile
 - b. Species
 - c. Life stage
- 8. Predominant Floodplain Habitat Type (terrace, depression, gravel-pit pond, canyon mouth)
- 9. Land Ownership (Federal, State, Private, Tribal)
- 10. Photo Imagery
 - a. Type (ASCS-35-mm, U.S. Army Corps of Engineers, University of Montana, Color Infrared)
 - b. Catalog No.

Scoring and Prioritization of Sites

Several classification systems were evaluated for suitability in classifying and ranking bottomland habitats that were perceived to be potentially valuable for providing endangered fish habitat (Cowardin et al. 1979; Brinson 1992; The Nature Conservancy 1992; Lyon 1993). Constraints of time and personnel precluded using these detailed classification systems for the bottomland inventory. Instead, a five-step procedure was developed in 1994 by a three-member team² to score and rank bottomland sites. Prioritizing sites assisted in identifying locations where staff gauges and river transects would be established in 1994 to predict flows necessary to inundate a bottomland site. The five steps were:

Step 1: Initially, criteria were developed to assist in discriminating and prioritizing bottomland sites using available photographic imagery and the most current biological information. The initial list included area, ownership, proximity to razorback sucker use (historical and current), water

² -----
 USFWS: Pat Nelson, Coordinator of the Floodplain Habitat Restoration Program, Region 6, Denver, Colorado; Bob Burdick, Fishery Biologist, Colorado River Fishery Project (FWS), Grand Junction, Colorado; and David Irving, Fishery Biologist, Colorado River Fishery Project, (FWS), Vernal, Utah. The meeting was facilitated by Chuck Gallagher (BR), Grand Junction, Colorado.

duration (June-September), hydrological connection to river during runoff (May and June), wetland feature (yes/no), suitability for fish or water control structures, presently diked or rip-rapped (yes/no), and potential for a network of sites (within 25 river miles of known historical or current razorback sucker use)(Appendix H; Table H.1.).

Step 2: Because many of the bottomland sites had not been visited prior to the prioritization process, five of the initial criteria considered were discarded. The five criteria eliminated were, 1) area, 2) wetland feature, 3) suitability for fish or water control structures, 4) presently diked or rip-rapped, and 5) water duration. Other criteria necessary to prioritize sites such as screening for contaminants, densities of non-native fishes and their species composition, and geomorphology would require site visits. Because there was not enough time to visit all sites prior to the high water of 1994, criteria were chosen that could be used with available photographic imagery to determine bottomland sites where preliminary hydrology screening would be conducted in the spring of 1994. From the initial nine criteria, four selection criteria with common characteristics among bottomland sites were chosen. These four criteria were used to evaluate, score, and rank bottomland sites for planning the 1994 hydrology studies. These four selection criteria are considered preliminary and could be revised, substituted, or supplemented if warranted by site visits. If alternative or additional criteria are developed, all bottomland sites listed in this inventory would have to be re-scored, re-ranked, and a new prioritization list generated. The four selection criteria used were:

- A. Status of land ownership
- B₁. Green River: proximity to a known or planned razorback sucker spawning area
- B₂. Colorado River: proximity to recent adult razorback sucker captures (1974-1993)
- C. Having a June hydrological connection to the river
- D. Potential for a network (complex) of bottomland sites

Step 3: The selection criteria were compared against each other to obtain a weighting factor. A paired analysis allowed a comparison of one criterion against the other three criteria. A point value was subjectively assigned to each selection criteria as follows:

- 4 = major importance
- 3 = medium importance
- 2 = minor importance
- 1 = no preference

Each team member, assigned a point value for each selection criterion. The point values of team member were totaled and averaged. The average was rounded to the nearest whole number to give a total average point value. Next, the selection criteria were classified with the highest total average point value being the most important selection criteria and the lowest total average point value being the least important criteria (Table 1).

Step 4: Sub-criteria for each of the four selection criteria were necessary in order to ensure that bottomland sites could be classified without tie-scores. Each selection criterion was then further divided into sub-criteria that were subjectively assigned a point value as follows:

Criterion A: Status of land ownership

- 3 = Federal ownership
- 2 = State ownership
- 1 = Private or Tribal ownership

Table 1. Paired comparison analysis of the four selection criteria^a. When a criterion is compared to itself it receives a point value of 1 (e.g., A:A = 1). When two criteria are compared, the one that is selected as being more important receives the point value. For example, B:A means that criterion B was perceived to contribute more to endangered fish recovery than criterion A, and therefore received the point value. Because criterion B was chosen over criterion A, the A:B comparison received no point value.

	A ^a	B ^a	C ^a	D ^a
A	A:B 0	B:A (3+3+4)/3=3.3	C:A (4+4+4)/3=4.0 ^b	D:B 0
B	A:B 0	B:B 1	C:B 0	D:B 0
C	A:C 0	B:C (4+4+4)/3=4.0	C:C 1	D:C (3+3+3)/3=3.0
D	A:D 0	B:D (3+3+4)/3=3.3	C:D 0	D:D 1
Total Average Point Value ^c	(1+0+0+0)=1	(3.3+1+4.0+3.3)=12	(4.0+0+1+0)=5	(3.3+0+3.0+1)=7

^a Criterion A = Land ownership status, Criterion B = Proximity to a known or planned razorback sucker spawning area (Green River) or proximity to recent adult razorback sucker captures (Colorado River), Criterion C = June hydrological connection to river, Criterion D = Potential for a network (complex) of bottomland sites.

^b The equation (3+3+4)/3 = 3.3 represents two events. The (3+3+4) portion of the equation is the three point values assigned the criterion by each team member. The product of the equation or 3.3 is the average weighted point value of the selected criterion.

^c The total average point value is the sum of the average weighted point values for each selected criterion. (These values were rounded to the nearest whole number.)

Criterion B₁ (Green River): Proximity to a known or planned razorback sucker spawning area

- 5 = within 1-25 river miles downstream of a spawning area
- 4 = within 26-50 river miles downstream of a spawning area
- 3 = within 51-75 river miles downstream of a spawning area
- 2 = more than 75 river miles downstream of a spawning area
- 1 = upstream of a spawning area

Criterion B₂ (Colorado River): Proximity to recent adult razorback sucker captures (using data from 1974-1993, [Burdick 1992; Quarterone 1993])

- 5 = 100 captures
- 4 = 50-99 captures
- 3 = 15-49 captures
- 2 = 6-14 captures
- 1 = 0-5 captures

Criterion C: Having a June hydrological connection to the river

- 3 = yes
- 2 = unsure
- 1 = no

Criterion D: Potential for network (complex) of bottomland sites

- 5 = yes, ≥ 8 sites
- 4 = yes, 6-7 sites
- 3 = yes, 4-5 sites
- 2 = yes, 2-3 sites
- 1 = no, only 1 site

Step 5: The sub-criteria raw scores were then multiplied by the total average weighted point values to produce a total score for each bottomland site. The total score was sorted from the lowest to the highest with the highest total score receiving the lowest ordinal rank and lowest total score the highest ordinal rank. The lowest ordinal rank was the top priority site and the highest ordinal rank was the lowest priority site in the bottomland inventory. The Green River drainage was scored and prioritized independently from the Colorado and Gunnison river drainages.

Geomorphology

Three general geomorphological categories were used to simplify bottomland classification and identify habitat types. These categories were floodplain depression, floodplain terrace, or gravel-pit pond (Figure 2). These three categories were the most numerous encountered in the inventory, although there were other habitat types such as canyon mouths.




Figure 2. Schematic of the bed profile of the three major bottomland habitat classifications at various flow regimes in the Upper Colorado River Basin.

Ecology

Five candidate sites were selected prior to the bottomland inventory and were proposed as sites to conduct demonstration studies aimed at testing hypotheses regarding flooded bottomland restoration for endangered fishes. Wetland hydrology, water chemistry, and vegetation and invertebrate inventory studies were completed at these five bottomland sites in spring 1994. The studies were conducted at two sites along the Colorado River, in the Moab Slough bottomland site (Cooper and Severn 1994a), and at the 29-5/8 Road gravel-pit pond site (Cooper and Severn 1994b). The Escalante State Wildlife Area bottomland site (Cooper and Severn 1994c) on the Gunnison River and two sites along the Green River, the Escalante Ranch bottomland (Cooper and Severn 1994d) and one area within the Ouray National Wildlife Refuge (Old Charlie Wash [Woods Bottom] Cooper and Severn 1994e) were also studied. A conceptual plan was developed for each of these five sites that discussed experimental strategies, research, and evaluation that could be used to restore and enhance various types of bottomland habitats to assist recovery of endangered fishes. (Moab-Collins 1994; Escalante Ranch-Richardson 1994; Old Charlie-Modde and Irving 1994; Escalante State Wildlife Area-Burdick 1994a and 29-5/8 Road gravel pit pond-Burdick 1994b).

Hydrology

Stage vs. Discharge Relationship

Hydrological parameters such as the timing, magnitude, and duration of flooding at a site are pivotal information to determine if a site is suitable for enhancement or restoration. This information was used to screen sites by determining how often or frequent does a site flood and how long (duration) does the site remain wetted following flooding.

Temporary staff gauges were placed at seven bottomland sites adjacent to the Green River and at nine sites adjacent to the Colorado and Gunnison rivers to collect flow vs. elevation data from May through July 1994. These data would be used to establish stage vs. discharge relationships for each site and to estimate a range of flows necessary to inundate the bottomland site. Staff gauge placement corresponded nearest low areas along the river channel where water would enter into the adjacent bottomland areas. However, in some instances, the staff gauge was not placed at the lowest point because cameras would have been inundated at high water. Automatic, programmable 35-mm cameras were placed inside plexiglass-faced ammunition boxes and were aimed at each staff gauge to photo-document water level changes over time. Each camera was set to take pictures that ranged from 4 to 12 hours. Film was exchanged about every 10 to 14 days. The elevation of the top of the staff gauge was surveyed as a future reference point. Two ground reference control points were established at each site. In the Colorado and Gunnison rivers, the control point was arbitrarily set at a relative elevation of 1000.0 feet (305 m). In the Green River drainage, elevation control points were absolute (feet above sea level). Reinforcing rod with an aluminum survey cap served as the control point headpin.

The staff gauge water-level changes were recorded on computer spreadsheets. Daily streamflow data were obtained from USGS stream gauges nearest the bottomland site for dates and times when stage elevations were recorded. This information was used with BR flow models to determine a stage vs. discharge relationship for each site. From several hundred possible equations, this model provides the equation which best describes the stage vs. discharge relationship and the regression correlation coefficient. The information was used to estimate flows necessary to inundate each of the bottomland sites.

Channel Cross Sections

To determine if channel configuration (depth and width) at each of these bottomland sites would remain constant or change over time with various discharges, it was necessary to obtain streambed profile information. If there were a change, this might affect the amount of flow necessary to inundate the site or the amount of fill material that would have to be removed to allow water access to the bottomland site. Therefore, transects were established at each bottomland site where staff gauges had previously been located. Kevlar cable was stretched across the entire channel between two steel fence posts. Channel cross sections usually encompassed the entire 10-year floodplain. Transects were surveyed by BR personnel from high points from the right shoreline to the left shoreline that crossed secondary and tertiary channels, backwaters, and islands. All distance and elevation measurements were obtained with an electronic distance meter (EDM), that corrected for up- and downstream distances from the transect. Across the transect, elevations were recorded at every 10-foot interval or at every significant grade break. Bed width was plotted against either absolute (Green River) or relative (Colorado and Gunnison rivers) elevations.

River Discharge

Mean daily streamflows were correlated with the dates when staff-gauge data were obtained. Also, mean daily flows for pre- and post-water development periods were obtained to generate flow exceedences. The period of record at a stream gauging station that best represented present water availability which would inundate a bottomland site was used. The Jensen, Utah gauge was used for bottomland sites adjacent to the Green River. The pre-water development period, 1947-1966, and post-water development period, 1966-1993, were analyzed at the Jensen gauge. For the two sites on the Gunnison River, the Delta gauge (1977-1993) was used for the Confluence Park bottomland site and the Whitewater gauge (1967-1993) near Grand Junction for the Johnson Boy's Slough site. Stateline gauge (1952-1993) was used for the Adobe Creek site along the Colorado River. The Cameo gauge (1952-1993) was used for the two sites between Debeque and Rifle, Colorado. For the four bottomland sites in the 15-mile reach, a different approach was used to determine streamflows. Although there is a USGS stream gauge immediately downstream of the Grand Valley Diversion, there are only three years of flow data. Therefore, the mean daily flows at the Cameo and Plateau Creek (1970-1993) stream gauges were summed and the water diversions at Government

Highline and Grand Valley Diversion were subtracted.

Land Ownership Status

A team³ composed of fisheries and realty experts contacted landowners who owned bottomlands along the Green, Colorado, and Gunnison rivers in August and September 1994 to discuss options for acquisition, lease, or easement. Either on-site visits or telephone contacts were made with landowners of the sites that were initially selected for hydrology screening. For the Green River, these bottomland sites included the Meril Snow Ranch, Little Stewart Lake, Baeser Bend, Brennan Bottom, and Pariette Draw. For the Gunnison River, these bottomland sites included Johnson Boy's Slough and Confluence Park. For the Colorado River, Adobe Creek, Griffith's, Clifton Sanitation No. 1, Pike's, and Battlement Mesa. During each visit, team members explained the purpose of the Recovery Program for the Endangered Fishes of the Upper Colorado, distributed the pamphlet entitled, "Swimming Upstream: The endangered fish of the Colorado River", and provided a prepared statement of commonly asked questions with answers (see Appendix A).

Permission to access private property was requested to collect additional biological data, conduct on-site screening for contaminants, and to acquire river channel and bottomland profile survey data. A Right-of-Entry Permit form was developed for signature to all parties for legal purposes (see Appendix A).

RESULTS AND DISCUSSION

Photographic Data

Prior to 1993, there were few sources of up-to-date photographic coverage that temporally and spatially covered warmwater reaches of the Upper Colorado River Basin. Most available maps, photographs, and videos were either confined to the main river channel and did not include adjacent floodplain and bottomland areas or were too small in scale to be useful in determining areas that would be inundated. A complete set of high-altitude, infrared photographs would have been the preferred imagery for this survey because they were scaled and easiest to interpret, but were too expensive.

A plan for complete and partial inventories of the Upper Colorado River Basin is presented in Appendix C. The videography could not be used because it was taken only for specific stream reaches and did not provide consistent information useable for the entire bottomland survey.

Over 2,000 color 35-mm slides were taken during four low-level reconnaissance photography flights over bottomland habitats in the Upper Colorado River Basin. These photographs are documented and stored in binders at the Colorado River Fishery Project offices in Vernal, Utah (Green, Yampa,

³ The Green River team consisted of David Irving, FWS, Dave Webster, NRCS, and Doyle Dow, BR. The Colorado River team consisted of Bob Burdick, FWS, and Doyle Dow.

and White rivers), and in Grand Junction, Colorado (Colorado and Gunnison rivers).

Water Discharge on Photograph Dates

Aerial photographs were taken in May 1993, 3 days before peak flow in the Green River and 2 days after peak flow in the Yampa and White rivers (Figure 3). Photographs were taken on the lowest flow day in September in the Green River and about 3 weeks after low flow in the Yampa and White rivers.

Aerial photographs were taken in June 1993, 13 days after peak flow in the Colorado River and 23 days after peak flow in the Gunnison River (Figure 3). Photographs were taken on the lowest flow day in September in the Gunnison River and about three weeks after low flow in the Colorado River.

Distribution of Bottomland Habitats


Green River. There were 132 bottomland habitat sites with a potential (historical) of 18,430 acres (7,459 ha) along the Green River (rm 0-393 [rkm 0-632]) in 1993 (Appendix D; Table D.1.). Of this total, 7,720 acres (3,124 ha; 74 sites) were inundated during the May high-water flow period and 2,438 acres (987 ha; 28 sites) in the September low-water flow period.

The largest flooded bottomland areas (Figure 4) were located in the Browns Park National Wildlife Refuge reach (rm 368-393 [rkm 592-633]; 1,371 acres [555 ha]) and between Pariette Draw and the Escalante Ranch (rm 238-310 [rkm 383-499]; 6,093 acres [2,466 ha]). The greatest concentration of flooded bottomland sites were within the Ouray National Wildlife Refuge (rm 249-265 [rkm 401-427]; 2,265 acres [917 ha]). Only 16% (12 sites) of the 74 sites flooded in May had a hydrological connection to the river, whereas 89% (24 sites) of the 28 sites that were wetted in September were still connected to the river (Appendix E; Tables E.1 and E.2). The majority of these sites had only one connection to the river. Sixteen percent (12 sites) of the May flooded sites received some water from gravity fed (irrigation) water sources and 84% (62 sites) received water from groundwater sources.

Of the 132 potential bottomland sites, 4% (6 sites) had natural levees between them and the river, whereas 11% (15 sites) had human-made levees or berms. Over 70% (93 sites) had emergent vegetation in May.

There were 30 floodplain depression bottomland habitat types and 99 floodplain terrace habitat types along the Green River. The remaining flooded habitat types consisted of two flooded gravel pits and one flooded river confluence.

Yampa River. Only 22 acres (9 ha; 1 site) of bottomland habitat were available in the Yampa River just upstream of the Yampa and Snake river confluence (rm 48 [rkm 77]) in 1993 (Appendix D; Table D.1.). It was flooded during the May high-water period, but dry in the September low-flow period



(100-300 cfs). It was connected to the river by a side channel in May, received some of its water from groundwater sources, and had no emergent vegetation (Appendix E; Tables E.3. and E.4.).

White River. A potential (historical) of 634 acres (257 ha; 2 sites) of bottomland habitat was available adjacent to the White River (rm 0-93 [rkkm 0-37.6]) in 1993 (Appendix D; Table D.1.). Of this total, 515 acres (208 ha; 2 sites) were flooded during the May high-water flow period, and 18 acres (7 ha; 1 site) were inundated during the September low-flow period.

The oxbow downstream of Highway 65 (rm 92 [rkkm 148]) was inundated from May through September, whereas the confluence of the White and Green rivers (rm 0-3 [rkkm 0-5]), with the greatest bottomland area, was dry in September. Both the oxbow and confluence sites were connected to the river in May and the oxbow site was connected to the river in September (Appendix E; Tables E.5. and E.6.). Both sites received some of their water from groundwater sources, but only the confluence site had emergent vegetation. The oxbow site was a flooded side-channel and the confluence site was a floodplain terrace.

Colorado River. There were 110 bottomland habitat sites with a potential (historical) of 12,315 acres (4,984 ha) along the Colorado River (rm 0-241 [rkkm 0-388]) in 1993 (Appendix D; Table D.2.) Of this total, 2,472 acres (1,000 ha; 101 sites) were inundated during the May and June high-water flow period and 1,135 acres (459 ha) in the September low-water flow period.

There were three distinct areas where bottomland habitats were concentrated. These were usually in large floodplain areas near agricultural centers. The first area was the stream reach between Debeque and Rifle, Colorado (rm 203-240 [rkkm 327-386]); 2,248 acres [910 ha]), the second was the stream reach in the Grand Valley between Loma and Palisade, Colorado (rm 152-185 [rkkm 245-298]; 4,564 acres [1,847 ha]), and the third area between the McGraw and Hotel Bottoms and the Cisco Boat Landing (rm 99-110 [rkkm 159-177]; 1,214 acres [491 ha]). Moab Slough (rm 62-64 [rkkm 100-103], downstream of these three areas near Moab, Utah, is the largest single bottomland area along the Colorado River encompassing approximately 875 acres (354 ha)(Figure 5).

Bottomland areas in canyon-bound stream reaches of the Colorado River were typically small in acreage. These areas included Canyonlands National Park downstream of Moab, Utah (rm 61-0 [rkkm 98-0]), Moab Canyon (rm 64-77 [rkkm 103-124]), Hittle Bottom to McGraw Bottom (rm 88-99 [rkkm 142-151]), Ruby and Horsethief canyons (rm 131-152 [rkkm 211-245]), and Debeque Canyon (rm 188-204 [rkkm 303-328]).

Seventy-one percent (71 sites) of the 100 sites flooded in May had a hydrological connection to the river, whereas 77% (43 sites) of the 56 sites still wet in September were connected to the river (Appendix E; Tables E.7. to E.13.) Although 71 sites were inundated during May and June, 26 sites had only one surface connection to the river and 15 sites had both an up- and downstream surface connection. Twenty-one percent of the May flooded sites received some water from gravity-fed (irrigation) water sources and 56%





Figure 4. Actual and potential (historic floodplain) acres (hectares) of bottomland habitat adjacent to the Green River, Colorado and Utah, 1993.

COLORADO RIVER BOTTOMLAND INVENTORY



GUNNISON RIVER BOTTOMLAND INVENTORY




Figure 5. Areas inundated during high water (May and June) and low water (September) in 1993 compared with areas that historically flooded on 240 miles of the Colorado River from the Green River confluence upstream to Rifle, Colorado and 75 miles of the Gunnison River from the Colorado River confluence upstream to the North Fork confluence. Inundated areas include floodplain depressions and terraces, side channels, canyon mouths, and artificial habitats such as gravel-pit ponds. There were 110 sites inventoried on the Colorado River and 48 sites on the Gunnison River.

received water from groundwater sources.

Along the 110 potential bottomland sites, we estimated 13.4 miles (21.6 km) of natural levees and 26.6 miles (42.9 km) of human-made dikes and levees. Of the various bottomland habitat types, 41 were floodplain terraces, 23 gravel-pit ponds, 22 side channels, and 16 canyon mouths. The remaining flooded habitat types consisted of 13 floodplain depressions and one oxbow. All 13 depressions were located between Moab Slough and the Green River confluence. Three of these depression habitat sites (rm 50.9-52.5 [rkkm 82-84], rm 33.2-35.6 [rkkm 53-57], and rm 17.5-18.7 [rkkm 28-30]), may be important off-channel habitats for razorback sucker drifting downstream if an expected or planned upstream spawning area is identified. Although these areas are wet during high water, they appear isolated from the river in photos and water enters only by groundwater infiltration and, therefore, fish do not have access to these areas. Further on-ground investigation of these areas is needed to determine water depth during runoff to monitor water duration, and to evaluate if fish can access these areas during high water.

Gunnison River. There were 48 bottomland habitat sites with a potential (historical) of 3,227 acres (1,306 ha) along the Gunnison River (rm 0-75 [rkkm 0-121]) in 1993 (Appendix D; Table D.2.). Of this total, 828 acres (335 ha; 41 sites) were inundated during the May and June high water period and 161 acres (459 ha) in the September low-water flow period.

In the Gunnison River, there was only one general area where bottomland habitats were concentrated. This was in the 17-mile stream reach in the Delta, Colorado, area that extends from Roubideau Creek upstream to Austin (rm 50-67 [rkkm 81-108]). There were 1,739 acres (705 ha) of potential floodplain habitat in this 17-mile reach. The remaining 50 miles of warmwater stream reach downstream of Roubideau Creek can be predominantly classified as canyon-bound with few bottomland habitat sites.

Eighty-five percent of the 41 sites flooded in May had a hydrological connection to the river, whereas 76% (13 sites) of the 17 sites still wet in September were connected to the river (Appendix E; Tables E.14 to E.16.). Although 41 of 48 sites were inundated during May and June, only 17 sites had one surface connection to the river and 16 sites had both up- and downstream surface connection. Thirteen of the May flooded sites received some water from a gravity-fed (irrigation) water source.

Along the 48 potential bottomland sites, we estimated 1.2 miles (2 km) of natural levees and 8.3 miles (13.4 km) of human-made dikes and levees. Of the various bottomland habitat types, 35 were floodplain terraces, six gravel-pit ponds, three oxbows, and two side channels. The remaining habitat types consisted of one floodplain depression and one canyon mouth.

Site Scoring and Prioritization

All bottomland habitat sites identified, 135 in the Green River Drainage (Appendix F; Table F.1.) and 158 in the Colorado River drainage (Appendix F;

Table F.2.), were scored and ranked according to the five-step prioritization procedure.

Green River Drainage

There were 132 bottomland habitat sites inventoried in the Green, one site in the Yampa, and two sites in the White River. The highest rank was "1" and the lowest rank "19". Bottomland habitats between Pariette Draw and Escalante Ranch (32 sites, rm 238-310 [rkm 383-499]) were the top ranked sites (1-11)(Table 2; Figure 6). There were tie-scores for several sites; hence several sites had identical ranks. Bottomland habitats in the Ouray National Wildlife Refuge (6 sites, rm 249-265 [rkm 401-427]) contained the highest ranked (1st and 2nd) sites. The Browns Park National Wildlife Refuge (8 sites, rm 368-393 [rkm 592-633]) sites ranked 12th-14th, followed by lower ranked (17th-18th) sites from Pariette Draw downstream to the Green and Colorado river confluence (92 sites, rm 0-238 [rkm 0-383]). The one Yampa

Table 2. The top 11 ranked bottomland sites adjacent to the Green River (GR), 1993. Refer to Appendix F; Tables F-1 to F-3 for a complete list of ranked sites in the Green River sub-basin.

Rank	General Area	Site Description	River: River Mile
1	Pariette Draw to Escalante Ranch	Johnson Bottom, Ouray NWR	GR: 263.0-265.0
1	Pariette Draw to Escalante Ranch	Leota Pond Complex, Ouray NWR	GR: 257.0-262.0
2	Pariette Draw to Escalante Ranch	Little Stewart Lake	GR: 295.5-297.5
2	Pariette Draw to Escalante Ranch	Brennan Bottom	GR: 262.0-266.0
2	Pariette Draw to Escalante Ranch	Wyasket Lake, Ouray NWR	GR: 253.0-257.0
2	Pariette Draw to Escalante Ranch	Sheppard Bottom, Ouray NWR	GR: 254.0-256.0
2	Pariette Draw to Escalante Ranch	Old Charlie Wash (main)/Woods Bottom, Ouray NWR	GR: 249.0-252.0
2	Pariette Draw to Escalante Ranch	Old Charlie Wash (diked)/Woods Bottom, Ouray NWR	GR: 249.0-250.0
3	Pariette Draw to Escalante Ranch	Stewart Lake	GR: 299.0-300.0
3	Pariette Draw to Escalante Ranch	Ashley Creek confluence area	GR: 297.0-298.5
4	Pariette Draw to Escalante Ranch	Escalante Ranch	GR: 302.5-309.5
4	Pariette Draw to Escalante Ranch	Meril Snow Ranch	GR: 302.0-303.0
4	Pariette Draw to Escalante Ranch	Gravel Ponds at Jensen	GR: 301.0-302.0
4	Pariette Draw to Escalante Ranch	Spring Hollow	GR: 295.0-296.0
5	Pariette Draw to Escalante Ranch	Bonanza Bridge Area	GR: 288.5-289.0
5	Pariette Draw to Escalante Ranch	Collier Draw	GR: 285.5-286.5
6	Pariette Draw to Escalante Ranch	Walker Hollow	GR: 294.0-295.0
6	Pariette Draw to Escalante Ranch	Alhandra Ferry Site	GR: 292.0-294.0
6	Pariette Draw to Escalante Ranch	Gravel Pits	GR: 292.0-293.0
7	Pariette Draw to Escalante Ranch	Mouth of Willow Creek	GR: 239.0-241.0
7	Pariette Draw to Escalante Ranch	Pariette Draw	GR: 238.0-241.0
8	Pariette Draw to Escalante Ranch	Downstream of Baeser Bend	GR: 269.0-272.0
8	Pariette Draw to Escalante Ranch	Upstream of Brennan Bottom	GR: 267.0-269.0
8	Pariette Draw to Escalante Ranch	Upstream of Brennan Bottom	GR: 266.0-267.0
8	Pariette Draw to Escalante Ranch	Ouray Ute pasture land	GR: 248.0-251.0
8	Pariette Draw to Escalante Ranch	Duchesne River confluence area	GR: 248.0-249.0
8	Pariette Draw to Escalante Ranch	White River confluence area	GR: 248.0-249.0
8	Pariette Draw to Escalante Ranch	West Branch area	GR: 243.0-247.0
8	Pariette Draw to Escalante Ranch	Tia Juana Bottom	GR: 242.0-244.0
9	Pariette Draw to Escalante Ranch	Hamacker Bottom/Baeser Bend	GR: 271.0-274.0
10	Pariette Draw to Escalante Ranch	The Stirrup	GR: 274.0-277.0
11	Pariette Draw to Escalante Ranch	Horseshoe Bend Area	GR: 277.0-284.0




Figure 6. Actual and potential (historic floodplain) acres (hectares) of the top eleven ranked bottomland habitat sites adjacent to the Green River, Utah, 1993.

River site ranked the lowest (19th), whereas the two White River sites ranked in the middle (oxbow: 15th; confluence: 10th).

Based on these rankings, there are five bottomland habitat areas in the Green River. The top ranked areas are in the Ouray National Wildlife Refuge, the next ranked areas in the Jensen to Brennan Bottom, followed by sites downstream of Ouray. The lowest ranked sites are in the Browns Park National Wildlife Refuge and the area from Desert Spring and Moon Bottom to the Green and Colorado river confluence.

Colorado River Drainage

There were 110 bottomland habitat sites inventoried in the Colorado River and 48 sites in the Gunnison River. The highest rank was "1" and the lowest "28". Unlike the Green River drainage, the top 13 ranked sites were scattered among four general areas (Table 3; Figure 7). Several bottomland sites had identical scores (rank 8th and 12th-two sites each; rank 9th and

Table 3. The top 13 ranked bottomland sites adjacent to the Colorado (CO) and Gunnison (GU) rivers, 1993. Refer to Appendix F; Tables F-4 to F-10 for a complete list of ranked sites in the Colorado River sub-basin.

Rank	General Area	Site Description	River: River Mile
1	18-mile Reach	Walter Walker State Wildlife Area	CO: 162.7-165.1
2	Delta Area	Johnson Boy's Slough	GU: 53.2- 54.2
3	15-mile Reach	Clifton Pond Area	CO: 177.7-178.2
4	Debeque Area	Debeque I-70 Slough	CO: 209.6-211.4
5	15-mile Reach	Labor Camp	CO: 182.9-183.6
6	18-mile Reach	Panorama	CO: 163.1-163.6
7	Debeque Area	Stoddard Property	CO: 209.4-210.1
8	Delta Area	Escalante State Wildlife Area (North)	GU: 50.8- 52.9
8	Delta Area	Escalante State Wildlife Area (South)	GU: 50.2- 52.4
9	Debeque Area	Etter's	CO: 204.7-206.6
9	15-mile Reach	Humphrey's	CO: 175.3-176.6
9	15-mile Reach	Griffith's	CO: 174.1-176.5
9	Debeque Area	Latham's	CO: 207.9-208.8
9	Debeque Area	"No-name"	CO: 206.2-207.5
10	18-mile Reach	Walter Walker South	CO: 164.4-166.0
10	18-mile Reach	"No-name"	CO: 169.4-169.6
10	18-mile Reach	"No-name"	CO: 161.0-162.0
10	Debeque Area	Battlement Mesa	CO: 220.7-221.7
11	15-mile Reach	"No-name"	CO: 181.7-182.2
11	Debeque Area	"No-name"	CO: 202.3-204.2
11	Debeque Area	"No-name"	CO: 201.9-202.4
11	Debeque Area	EXXON	CO: 218.9-220.1
11	Debeque Area	Wallace Creek Island	CO: 215.9-216.5
12	15-mile Reach	"No-name"	CO: 183.3-184.2
12	15-mile Reach	"Hotspot Junction"	CO: 173.9-175.1
13	15-mile Reach	Clifton Water Treatment	CO: 179.1-181.1

Bottomland Habitat Rank




Figure 7. Areas inundated during high water (May and June) and low water (September) in 1993 compared with areas that were historically flooded for the top 13 ranked bottomland sites adjacent to the Colorado and Gunnison rivers.

11th-five sites each; rank 10th-four sites). Along the Colorado River, there were 10 bottomland habitat sites within the Debeque area (rm 202.3-221.7 [rkm 326-357]), eight sites within the 15-mile reach (rm 173.9-184.2 [rkm 280-297]), and five sites within the 18-mile reach (rm 161.0-169.6 [259-273])(Table 3). There were three sites within the Delta area along the Gunnison River (rm 50.2-54.2 [rkm 81-87]).

The top ranked site was the Walter Walker State Wildlife Area in the 18-mile reach. Four other sites in the immediate vicinity ranked 6th and 10th. Johnson Boy's Slough along the Gunnison River ranked 2nd, the Clifton Pond area along the Colorado River ranked 3rd, and the Debeque I-70 Slough ranked 4th. Six other sites immediately downstream of the Debeque I-70 Slough ranked 9th and 11th.

Ecology

The wetland studies of the five initial bottomland sites conducted by Cooper and Severn (Table 4) were completed in 1993. Each site was unique and had varying levels of restoration potential. The bottomland habitat areas in Ouray National Wildlife Refuge, the Escalante State Wildlife Area, and the 29-5/8 Road gravel-pit pond have the greatest potential for razorback sucker restoration. The Escalante Ranch area and Moab Slough may require significant modification for restoration.

Hydrology

Green River Drainage

Stage vs. Discharge Relationship. Staff gauges were installed at seven different bottomland sites from mid-May (5/11) through late-June (6/28) and monitored for 34 to 48 days to obtain stage-discharge data during the high-flow period (Figures 8-14). Spring runoff in the Green, Yampa, and White rivers was moderately low in 1994 compared to moderately high flows in these rivers in 1993.

The maximum daily mean discharge in 1994 was 11,068 cfs on 20 May for the Green River at the Jensen stream gauge, 5,650 cfs on 18 May for the Yampa River at the Maybell gauge, and 2,529 cfs on 23 May for the White River at the Watson gauge. In 1993, the maximum daily mean discharge was 20,000 cfs for the Green River, 17,870 cfs for the Yampa River, and 4,570 cfs for the White River.

Stage-discharge data recorded at temporary staff-gauge sites were collected over a narrow range and at a lower magnitude of flows in 1994 compared to previous higher water years (Table 5). Flows in 1994 were lower than anticipated making it difficult to develop a reliable stage vs. discharge relationship.

The linear regression correlation coefficients (r^2) of the stage vs. discharge relationships were high (range of 0.83 to 0.97) at six of the seven

Table 4. Brief description of the five bottomland sites selected for demonstration projects in the Colorado and Green river drainages, Colorado and Utah.

River Bottomland Site	Reference	Comments/Recommendations
Colorado River Moab Slough	Cooper and Severn 1994a	Higher flows are necessary to inundate the bottomland habitat in Moab Slough. Razorback sucker larvae will also need to be introduced into the site. The area needs to receive seasonal flooding to flush out high concentrations of salt. Significant earthwork may be necessary to make the area fully functional.
29-5/8 Road Gravel Pit	Cooper and Severn 1994b	This pond could provide important fish habitat. No need to grade the pond bottom; break the dike and create a permanent access, let the pond fill and drain with natural river flows, and let fish move in and out throughout the year. Some periodic maintenance to keep the entrance or connection channel free of sediment may be necessary.
Gunnison River Escalante State Wildlife Area	Cooper and Severn 1994c	This site has as much potential for restoration as any other site in the Upper Colorado River Basin. It is dynamic and fluvially active, with fully functioning backwaters and oxbows and has areas that can be slightly modified to increase their functioning as bottomland habitat. If selenium in the Gunnison River and all irrigation and tributary flows remains high and it cannot be remediated, this may preclude use of the system in this recovery program.
Green River Ouray National Wildlife Refuge (Old Charlie Wash)	Cooper and Severn 1994d	Potentially one of the most valuable flood plain ecosystems in the entire Upper Colorado River Basin. Extensive bottomland areas occur behind natural and man-made levees adjacent to the main river channel. The greatest potential for re-coupling of the river with the flood plain occurs at the Ouray National Wildlife Refuge. Needs higher flows over a longer duration released from Flaming Gorge Dam.
Escalante Ranch area	Cooper and Severn 1994e	Contains localized concentrations of selenium, particularly in the northern and southern portions of the bottomland area. These areas may have selenium concentrations too high for fish restoration purposes. Channels may need to be constructed in some areas of the wetland to allow high water flushing flows to reduce selenium levels. This flooding may also provide an avenue for larval razorback suckers to enter the wetland.




Figure 8. River profile and stage versus discharge relationship for the Meril Snow Ranch bottomland habitat site adjacent to the Green River, Utah, 1994.




Figure 9. River profile and stage versus discharge relationship for the Little Stewart Lake bottomland habitat site adjacent to the Green River, Utah, 1994.




Figure 10. River profile and stage versus discharge relationship for the Baeser Bend bottomland habitat site adjacent to the Green River, Utah, 1994.




Figure 11. River profile and stage versus discharge relationship for the Johnson Bottom pond 2 bottomland habitat site adjacent to the Green River, Utah, 1994.




Figure 12. River profile and stage versus discharge relationship for the Leota Bottom Pond 7 bottomland habitat site adjacent to the Green River, Utah, 1994.




Figure 14. River profile and stage versus discharge relationship for the Woods Bottom (Old Charlie Wash) bottomland habitat site adjacent to the Green River, Utah, 1994.

sites. It was unfortunate, however, that high runoff flows were not available in 1994 to accurately predict the stage vs. discharge relationship. A larger data base is needed to more accurately verify the predicted stage vs. discharge relationship for the upper end of the May and June hydrograph. This is especially important when estimating the magnitude, frequency, and duration of flows necessary to flood a bottomland site.

Table 5. Range of flows and dates that stage-discharge data were collected in the Green River drainage, 1994.

Site	1994 Flows (cfs)	<u>Dates</u> MM/DD	Maximum Discharge (cfs)	
			1994	1993
Meril Snow Ranch	2,834-11,068	5/13-6/17	11,068	20,000
Little Stewart Lake	2,834-11,068	5/13-6/17	11,068	20,000
Baerer Bend	2,110-11,068	5/12-6/28	11,068	20,000
Johnson Bottom Pond 2	4,443-11,068	5/11-6/14	11,068	20,000
Leota Bottom Pond 7	3,645-11,068	5/11-6/15	11,068	20,000
Sheppard Bottom Pond 3	3,645-11,068	5/12-6/15	11,068	20,000
Woods Bottom	2,110-11,068	5/12-6/28	11,068	20,000

Channel Cross Sections. Channel cross sections at each of the seven bottomland sites were conducted in August, September, and October (Figures 8-14). The river channel widths ranged from 127 to 794 feet (39-242 m) and the bottomland sites ranged from 950 to 4,381 feet (290-1,335 m). The widest wetted cross section was at Johnson Bottom (4,381 feet [1,335 m]) and the narrowest cross section was at Baerer Bend (950 feet [290 m]) on the Green River.

Using the stage vs. discharge and flow-frequency data (Appendix I; Figure I.1.), the flow necessary to flood the particular bottomland (floodplain depression) for all seven sites was estimated (Table 6; Figures 8-14). For example, at the Meril Snow Ranch site (Green River rm 303.0 [rkm 487.6]) we estimated the floodplain depression would flood at about 14,000-15,000 cfs; these flows that might be experienced about one year out of three to four (since 1966). However, this site requires validation of the stage vs. discharge relationship because the correlation coefficient was the lowest of the seven sites ($r^2=0.64$). Re-establishing a staff gauge and obtaining another data set during runoff at all seven sites is necessary to refine the

stage vs. discharge relationship and obtain more reliable estimates of overbank flooding.

Table 6. Estimated overbank flooding and frequency of flooding at the seven bottomland habitat sites in the Green River (GR). Overbank flooding was determined from staff vs. discharge relationships and river profiles conducted for each site in 1994. Flow percentile data for the post-water development period were taken from USGS stream gauging stations nearest the site.

Site River Mile	Estimated Overbank/ Side Channel Flooding (cfs)	Percent of Time ^a Flood Flows Are Available	USGS Stream Gauge/ Time Period Evaluated
Meril Snow Ranch GR 303.0 ^b	14,000-15,000	25- 35%	Jensen ^b 1966-1993
Little Stewart Lake GR 297.0	18,000-19,000	10- 15%	Jensen 1966-1993
Baerer Bend GR 271.5	19,000-20,000	5- 10%	Jensen 1966-1933
Johnson Bottom Pond 2 GR 262.0	12,000-13,000	40- 50%	Jensen 1966-1933
Leota Bottom Pond 7 GR 258.0	20,000-22,000	1- 5%	Jensen 1966-1993
Sheppard Bottom Pond 3 GR 251.0	7,000- 8,000	65- 70%	Jensen 1966-1993
Woods Bottom GR 252.0	2,000- 4,000	95-100%	Jensen 1966-1993

^a For example 25-35% of the time = 2.5 to 3.5 years out of 10.

^b USGS gauge at Jensen, Utah, rm 317 (rkm 510).

The Little Stewart Lake, Baerer Bend, and Leota Pond 7 bottomland sites would flood between 18,000-22,000 cfs; one year out of ten. The Little Stewart Lake site has an old inlet ditch just upstream that could flood between 6,000-7,000 cfs every other year. Removal of about five feet of bank at the Baerer Bend site would allow flooding between 10,000-11,000 cfs every other year. The Johnson Pond 2 bottomland site would flood into the inlet ditch between 12,000-13,000, two years out of five. The Sheppard Bottom pond 3 site would flood into the inlet ditch between 7,000-8,000 cfs, seven years out of ten. The Woods Bottom site would flood into the inlet ditch between 2,000-4,000 cfs, nine out of 10 years.

Colorado River Drainage


Stage vs. Discharge Relationship. Nine staff gauges were installed at nine different bottomland sites during mid-May and monitored for approximately 45 days to obtain stage vs. discharge information for flows during the high-flow period (Figures 15-22). Information for one site (Pike's, Colorado River rm 179.1 [rkm 288]) was not useable because livestock repeatedly disturbed the position of the camera; at two other sites (Griffith's, Colorado River rm 176.0 [rkm 283]) and Clifton Sanitation No. 1., Colorado River rm 178.5 rkm 287]) the period of time information was collected was prematurely terminated because the cameras were stolen. Spring runoff in the Colorado and Gunnison rivers was moderately low in 1994 compared to 1993 which was considered a moderately high year. The maximum daily mean discharge recorded during 1994 for the Colorado River at the Stateline USGS gauge was 13,700 cfs on 2 June, whereas in the Gunnison River the maximum daily mean discharge at the Whitewater gauge was recorded at 6,040 cfs on 23 May. In 1993, the maximum daily mean discharge recorded for the Colorado River at the Stateline gauge was 44,000 cfs; in the Gunnison River the maximum daily mean discharge was 20,500 cfs.

Empirical stage data were collected over a narrow range and at a lower magnitude of flows in 1994 compared to previous higher water years (Table 7). Flows in 1994 were lower than anticipated to obtain a reliable stage vs. discharge relationship.

Table 7. Range of flows and dates that stage-discharge data were collected in the Colorado and Gunnison river drainages, 1994.

Site	Flows (cfs)	Date MM/DD	Maximum Discharge (cfs)	
			1994	1993
Johnson Boy's Slough	1,441- 6,040	5/16-7/ 7	6,040	20,500
Confluence Park	2,708- 5,054	5/ 9-6/ 4	5,000	15,400
Adobe Creek	7,540-13,700	5/12-6/13	13,700	44,000
30-29 Road	5,511- 8,619	5/16-6/ 2	8,930	25,900
Griffith'	3,974- 9,022	5/12-6/14	9,022	25,900
Clifton San. No. 1	5,698- 7,068	5/13-5/31	8,930	25,900
Pike's	No Data	---	8,930	25,900
EXXON	4,424-11,000	5/17-6/ 3	11,000	22,200
Battlement Mesa	6,419-11,000	5/17-6/ 8	11,000	22,200

The linear regression correlation coefficients (r^2) were high (range of 0.84 to 0.98) for stage vs. discharge relationships at five of the eight sites. However, it was unfortunate that high runoff flows did not occur in 1994 to accurately predict the stage vs. discharge relationship. Additional empirical data are necessary to more accurately verify the predicted stage vs. discharge relationship for the upper end of the May and June hydrograph. This information is even more important for estimating the magnitude, frequency,



Rank 4 Eqn 4 $y=a+bx^2$
 $r^2=0.33424166$ DF Adj $r^2=0.28985777$ FitStdErr=1557.474 Fstat=15.563442
 $a=-3016279.5$
 $b=3.0659516$





Figure 15. River profile (top) and stage vs. discharge relationship (bottom) for Adobe Creek bottomland habitat site adjacent to the Colorado River, river mile 159.9. Note: Q=discharge in cfs; W.S.El=relative water surface elevation.



Rank 42 Eqn 4 $y=a+bx^2$

$r^2=0.54354978$ DF Adj $r^2=0.51311976$ FitStdErr=503.56338 Fstat=36.915401

$a=-1413777.2$

$b=1.4329385$





Figure 16. River profile (top) and stage vs. discharge relationship (bottom) for 30-29 Road bottomland habitat site adjacent to the Colorado River, river mile 174.4. Note: Q=discharge in cfs; W.S.El=relative water surface elevation.



Rank 16 Eqn 4 $y=a+bx^2$

$r^2=0.84935748$ DF Adj $r^2=0.8446499$ FitStdErr=480 47104 Fstat=366.48508

$a=-1272981.1$

$b=1.2871999$





Figure 17. River profile (top) and stage vs. discharge relationship (bottom) for Griffith's bottomland habitat site adjacent to the Colorado River, river mile 176.0. Note: Q=discharge in cfs; W.S.El=relative water surface elevation.



Rank 18 Eqn 4 $y=a+bx^2$

$r^2=0.55656107$ DF Adj $r^2=0.52968598$ FitStdErr=363 17851 Fstat=42.673466

$a=-1274946.5$

$b=1.2950181$





Figure 18. River profile (top) and stage vs. discharge relationship (bottom) for Clifton Sanitation No. 1 bottomland habitat site adjacent to the Colorado River, river mile 178.5. Note: Q=discharge in cfs; W.S.El=relative water surface elevation.



Rank 23 Eqn 4 $y=a+bx^2$
 $r^2=0.98484665$ DF Adj $r^2=0.98406955$ FitStdErr=191 73785 Fstat=2599 6798
 $a=-1108156.2$
 $b=1.121198$





Figure 19. River profile (top) and stage vs. discharge relationship (bottom) for the EXXON bottomland habitat site adjacent to the Colorado River, river mile 220.0. Note: Q=discharge in cfs; W.S.El=relative water surface elevation.



Rank 61 Eqn 4 $y=a+bx^2$

$r^2=0.88133364$ DF Adj $r^2=0.87554504$ FitStdErr=304 18245 Fstat=311.9335

$a=-1458783.1$

$b=1.4811788$





Figure 20. River profile (top) and stage vs. discharge relationship (bottom) for the Battlement Mesa bottomland habitat site adjacent to the Colorado River, river mile 221.3. Note: Q=discharge in cfs; W.S.El=relative water surface elevation.



Rank 43 Eqn 4 $y=a+bx^2$

$r^2=0.93406376$ DF Adj $r^2=0.93212446$ FitStdErr=344.12458 Fstat=977.46555

$a=-1341993.8$

$b=1.3492054$





Figure 21. River profile (top) and stage vs. discharge relationship (bottom) for Johnson Boy's Slough bottomland habitat site adjacent to the Gunnison River, river mile 53.6. Note: Q=discharge in cfs; W.S.El=relative water surface elevation.



Rank 4 Eqn 4 $y=a+bx^2$

$r^2=0.97717443$ DF Adj $r^2=0.97638734$ FitStdErr=88.558137 Fstat=2525.8201

$a=-763679.21$

$b=0.77353342$




Figure 22. River profile (top) and stage vs. discharge relationship (bottom) for Confluence Park bottomland habitat site adjacent to the Gunnison River, river mile 57.1. Note: Q=discharge in cfs; W.S.El=relative water surface elevation.

and duration of flows necessary to flood a bottomland site. Therefore, the flows that were estimated to flood each of these bottomland sites should be considered provisional in this report.

Channel Cross Sections. Channel cross sections at each of the nine bottomland sites were conducted in August and September (Figures 15-23). The widest wetted cross section was at Adobe Creek (670 feet [204 m]); the narrowest cross section was at Confluence Park on the Gunnison River (200 feet [61 m]). Using the stage-discharge and flow-frequency data (Appendix I; Figures I.2-I.6), we estimated the flow necessary to flood the particular




Figure 23. River profile for Pike's bottomland habitat site adjacent to the Colorado River, river mile 179.1. Note: stage vs. discharge relationship not available for this site.

bottomland (tertiary side channel, depression, terrace, oxbow, etc.) for six of the nine sites (Table 8; Figures 15-22). For example, at Adobe Creek (Colorado River rm 159.9 [rkm 257]) we estimated the tertiary side channel would flood at about 19,000-20,000 cfs, flows that might be experienced about one year out of every three (since 1952). However, validation of the stage

Table 8. Estimated overbank flooding and frequency of flooding at the nine bottomland habitat sites in the Colorado and Gunnison rivers. Overbank flooding was determined from staff vs. discharge relationships and river profiles conducted for each site in 1994. Flow percentile data for the post-water development period were taken from USGS stream gauging stations nearest the site.

Site River Mile	Estimated Overbank/ Side Channel Flooding (cfs)	Percent of Time ^a Flood Flows Are Available	USGS Stream Gauge/ Time Period Evaluated
Johnson Boy's Slough GU 53.7 ^b	No Estimate	---	---
Confluence Park GU 57.1	8,000- 9,000	15-20%	GU @ Delta/ 1967-1993
Adobe Creek CO 159.9	19,000-20,000	20-25%	CO @ Stateline/ 1952-1993
30-29 Road CO 174.4	16,000-19,000	10%	15-mile ^c
Griffith's CO 176.0	9,000-10,000	50%	15-mile
Clifton Sanitation No. 1 CO 178.5	Reduce dike elevation to 996.0 9,600	50%	15-mile
Pike's CO 179.1	No Estimate	---	---
EXXON CO 220.0	No Estimate	---	---
Battlement Mesa CO 221.3	10,500-12,000	44-54%	CO @ Cameo/ 1952-1993

^a For example 15-20% of the time = 1.5 to 2 years out of 10

^b GU=Gunnison River; CO=Colorado River

^c The USGS stream gauge for the 15-mile reach was installed in 1991 and only three years of data are available. Therefore there is an insufficient data base to conduct a direct flow exceedence analysis for this reach. Another approach was used. 15-mile reach mean daily flow exceedence for June was determined by adding the mean daily flows at the Cameo and Plateau Creek USGS stream gauges for June for the post-water development period, 1970-1993. Diversions upstream of the 15-mile reach and downstream of the Cameo and Plateau Creek gauges were subtracted. The flow value used for the diversions for June was 1,679 cfs (2,260 minus 581).

vs. discharge relationship at this site is needed because the correlation coefficient was the lowest of the eight sites ($r^2=0.29$). Re-establishing a staff gauge and obtaining another data set during runoff at this site and at three other sites, Pike's on the Colorado River and Johnson Boy's Slough and Confluence Park on the Gunnison River, is necessary to refine the stage vs. discharge relationship and obtain more reliable estimates of overbank flooding. This should be conducted during runoff of a water year higher than that experienced in 1994.

At Confluence Park on the Gunnison River, inundation of a side channel was estimated to occur between 8,000-9,000 cfs, which now occurs about 15-20% (one and one-half to two years out of 10) since 1977. It is uncertain what minimum flow is required to inundate this site. It might be slightly higher than the estimated 8,000-9,000 cfs flow level because color aerial photographs (1:12000) taken 2 June 1993 confirmed that the Confluence Park site was inundated on this date when the mean daily streamflow at the Delta USGS gauge was 11,100 cfs.

To inundate the 30-29 Road Site, 16,000-19,000 cfs would be required, which presently occurs about once every 10 years (since 1970). Griffith's, a floodplain terrace with side channels, appears to flood more often (about once every two years) than any of the other six sites. Flows necessary to flood the side channel at this site were estimated to be about 9,000-10,000 cfs. The Battlement Mesa site presently floods a flow-through side channel about four to five times every 10 years (Table 8). At the Clifton Sanitation No. 1 site removing a portion of the human-made dike by reducing the present elevation by about four feet would reconnect a former side channel to the river more frequently (once every two years) during runoff. Dikes exist at Confluence Park, Johnson Boy's Slough, Adobe Creek, 29-5/8 Road, and Pike's. We did not estimate the flows necessary to flood these sites with complete or partial removal of dikes. However, the flows necessary to inundate bottomland habitats at each of these sites can be predicted using stage-discharge data (Figures 15-22). Given the present hydrologic regime, current water availability, and future compact water development in the Upper Colorado River Basin, inundating bottomland sites might be best accomplished by removing portions of- or entire human-made dikes or natural obstructions (e.g., levees) to increase flooding frequency. Those sites that flood more frequently under the existing hydrologic regime will also flood more frequently in the future, and will provide more benefits to endangered fish.

Land Ownership Status

Green River Drainage

Most (75%) of the bottomlands in the Browns Park National Wildlife Refuge area is Federal ownership (Appendix E; Tables E.1. and E.2.; Figure 24). Forty-two percent of the land between Escalante Ranch and Brennan Bottom is in private ownership. All of the Ouray National Wildlife Refuge bottomland sites are federally owned. Seventy-one percent of the bottomland sites are on tribal lands between Ouray and Pariette Draw. The majority of bottomland habitat between Desert Spring and the Green and Colorado river confluence is




Figure 24. Land ownership status in the Browns Park National Wildlife Refuge bottomland habitat area (rm 368-393), Escalante ranch to Brennan Bottom area (rm 262-310), Ouray National Wildlife Refuge area (rm 249-265), Ouray to Pariette Draw area (rm 238-251), and Desert Spring to Green and Colorado river confluence area (rm 0-249), Green River, Colorado and Utah, 1993.

federal land. The one Yampa River site was privately owned (Appendix E; Tables E.3. and E.4.), whereas the White River oxbow site was privately owned and the confluence site on tribal reservation land (Appendix E; Tables E.5. and E.6.).

Only five of the 12 private land owners were contacted in 1993. The owners of three of these bottomland sites, Baeser Bend, Brennan Bottom, and Pariette Draw, did not have any interest in selling their land or entertaining wildlife easements. The Meril Snow Ranch owner was not interested in selling his land but was receptive to development of a wildlife easement. All of the Little Stewart Lake property is privately owned, but approximately one-half presently has a wildlife easement by FWS. The easement is administered by the Ouray National Wildlife Refuge and allows FWS access and complete management of bottomland resources. The private landowners of the non-easement portion are not interested in selling their land or developing a wildlife easement.

None of the tribal-owned bottomland sites (Duchesne and White river confluence areas, West Branch, and Tia Juana Bottom) were available for land acquisition or wildlife easement. The Woods Bottom (Old Charlie Wash) bottomland site is owned by the tribe, but is currently being leased by FWS.

Colorado River Drainage

Of the 110 bottomland sites identified and inventoried along the 241 miles of the Colorado River, 51% or about 6,355 acres (2,572 ha) are in private ownership (Appendix H; Table H.1.; Figure 25). Of the 48 bottomland sites along the 75 miles of the Gunnison River, 70% or about 2,221 acres (899 ha) are in private ownership (Appendix H; Table H.1.; Figure 26). Fifteen percent (1,826 acres [739 ha]) of the bottomland sites along the Colorado River is in public (state and federal) ownership and 22% (697 acres [282 ha]) along the Gunnison River is in public (state and federal) ownership.

The greatest amount of bottomland habitat along the Colorado River owned by the federal government (83%) is within Canyonlands National Park (Figure 26). The federal government does not own any bottomland area between Rifle and Loma (Figure 26). The federal government owns or controls only 8% of the bottomland habitat along the Gunnison River and that lies within a canyon-bound area between Roubideau Creek (rm 50 [rkm 81]) and the Colorado River confluence.

It is apparent from the vast area that is owned or controlled by private landowners, willingness from these landowners to cooperate will have an important impact if the Floodplain Habitat Restoration Program is to succeed. Innovative and creative strategies and approaches to establish agreements, easements, or acquire private property may be necessary for the program to proceed.

Johnson Boy's Slough along the Gunnison River is privately and jointly owned by Kenneth and Wendell Johnson. Initially, they have been very cooperative in allowing access. Their property (about 500 acres [202 ha]), of which a majority is upland acreage and not necessarily bottomland habitat, is








Figure 25. Land ownership status of bottomland sites inventoried in five stream reaches adjacent to the Colorado River, Colorado and Utah, extending from Rifle, Colorado (river mile 241) downstream to the Green River confluence (river mile 0). Note: Land ownership by area (hectares) and percentages in Appendix G; Table G.1.

**North Fork Confluence-
Austin
RM 75-67**



**Austin-
Railroad Bridge
RM 67-50**



**Railroad Bridge-
Colorado R. Confluence
RM 50-0.7**





Figure 26. Land ownership status of bottomland sites inventoried in three stream reaches adjacent to the Gunnison River, Colorado extending from the North Fork River confluence (river mile 75) downstream to the Colorado River confluence (river mile 0.7). Note: land ownership by area (hectares) and percentages in Appendix G; Table G.1.

currently for sale. The former oxbow and side channel adjacent to the river have been identified as potential bottomland habitat. Confluence Park, the other site along the Gunnison River 4 miles upstream from Johnson Boy's Slough, is owned by five entities which include, 1) The City of Delta, 2) Fred Wild, 3) Ray Toole, 4) Delta Environment, Inc. (John Parker), and 5) Gerald Davidson. The first four landowners were receptive to allow further screening of their sites. The Davidson's were not interested in the program.

On the Colorado River the Adobe Creek site is owned by Mr. Oby Blanchard. He does not wish to sell his land but has allowed screening to continue and is receptive to further development at this site. The 30-29 Road Site is primarily owned by the State of Colorado, Department of Natural Resources, Division of Parks and Outdoor Recreation. They are willing participants under the Recovery Program. Acquisition is the only option available at the site owned by Mr. L. O. Griffith. This land is also being considered for purchase as mitigation under the Grand Valley Salinity Control Project (Personal communication, Pat Arbeiter, 1994). The site at rm 178.5 (rkm 287) is controlled by Clifton Sanitation No. 1. Their only option is to enter into an easement with the Recovery Program. Harold Pike is interested in selling a majority (about 18 acres [7 ha]) of his property which includes the pasture that is a depression adjacent to the river. This depression which is a former side channel is inundated during runoff but is currently protected at the downstream end by a dike that is comprised of loosely-placed cement fill.

The two bottomland habitat sites on the Colorado River between Rifle and Debeque are owned by corporations. Bill Wilde representing Battlement Mesa Partners indicated the bottomland site identified at rm 221.3 (rkm 356) will be not be developed and "left as is" for a greenbelt area. The other site immediately downstream at rm 220 (rkm 354) is owned by EXXON, Houston, Texas. EXXON's land manager has indicated that this bottomland is currently for sale.

RECOMMENDATIONS

1. Validate Overbank Flows. Flows that were estimated to flood each bottomland site are considered provisional. Staff gauges should be re-deployed at each of the seven bottomland sites in the Green River and at Adobe Creek and Pike's on the Colorado River and Johnson Boy's Slough and Confluence Park on the Gunnison River to obtain additional empirical data to verify the stage vs. discharge relationships. This is necessary so that the magnitude, frequency, and duration of flows necessary to flood these bottomland sites can be reliably estimated. High runoff flows did not occur in 1994 and the stage vs. discharge relationship could not be accurately predicted. Runoff flows in 1995 should be higher than 1994. Snowpack estimates as of 1 May 1995 indicate that the three major stream drainages in the Green River drainage are near 100% of normal (Green River: 201%; Yampa and White rivers: 120%); in the Upper Colorado River drainage both the Colorado (132%) and Gunnison River (161%) are over 100% (Personal communication, NRCS, Denver, Colorado, and Salt Lake City, Utah).

This work could be accomplished prior to and during runoff in 1995.

2. Complete Screening. Contaminant and geomorphological screening was not conducted at all of the initial 19 sites because of lack of time. Screening of bottomland sites such as Confluence Park and Johnson Boy's Slough on the Gunnison River, and Pike's (upper end of Clifton Sanitation No. 1 site), and Adobe Creek on the Colorado River should be conducted to determine if these sites are suitable and have potential for bottomland habitat restoration.
3. Additional Site Visits. Three bottomland sites along the Colorado River that were identified from photographic imagery (35-mm color slides) as depressions located downstream of Moab, Utah, deserve further on-ground investigation to determine if they have restoration potential. These sites (rm 50.9-52.5 [rkm 82-84], rm 33.2-35.6 [rkm 53-57], and rm 17.5-18.7 [rkm 28-30]), which are each less than 25 acres (10 ha), are currently heavily guarded by dense tamarisk stands and are only wetted during runoff via groundwater infiltration. These sites might provide off-channel habitats important for larval razorback sucker drifting downstream if an expected or planned upstream spawning area is identified. Since most of these sites have well established natural levees that have formed due to both reduced high flow frequencies and proliferation of tamarisk, some physical manipulation (e.g., reducing the height of the levee by excavation) of these sites would probably be required to allow fish and water access to these sites. Unfortunately, these sites are in canyon-bound areas and do pose logistical problems because they are remote and difficult to access.
4. Obtain Color Aerial Photographs. Obtain a complete set of current color aerial photographs of the entire Upper Colorado River Basin. This would include the Green River from the Colorado River confluence upstream to Browns Park, the Yampa River from the Green River confluence upstream to Cross Mountain Canyon, and the White River from the Green River confluence upstream to Taylor Draw Dam. For the Upper Colorado River, the Green River confluence upstream to Rifle and the Gunnison River from the Colorado River confluence upstream to the North Fork confluence. Photos should be at a scale between 1:6000 and 1:12000 and taken to coincide with runoff during May and June. Color photos should be taken during water years when high runoff flows are anticipated to identify those bottomland sites that may have potential for habitat restoration.
5. Re-rank Bottomland Sites. Four criteria were initially chosen that could be used with available photographic imagery and the best biological information to date to prioritize bottomland sites and to select a few (15-20) bottomland sites so that staff gauges could be installed prior to the 1994 runoff to obtain preliminary hydrology information at each site. These selection criteria could be revised, substituted, or supplemented as sites are

further screened and additional information comes forth. However, if alternative or additional criteria are developed, all bottomland sites listed in this inventory would have to be re-scored and re-ranked to generate a new prioritization list.

LITERATURE CITED

- Brinson, M.M. 1992. A hydrogeomorphic classification for wetlands. Technical Report WRP-DE-4. Biology Department, East Carolina University, Greenville, North Carolina. Prepared for Office, Chief Engineers, U.S. Army, Washington, D.C.
- Burdick, B.D. 1992. A plan to evaluate stocking to augment or restore razorback sucker in the Upper Colorado River. U.S. Department of Interior, Fish and Wildlife Service, Grand Junction, Colorado. 56 pp.
- Burdick, B.D. 1994a. Conceptual management plan for habitat enhancement in flooded bottomlands: Escalante State Wildlife Area, Gunnison River downstream of Delta, Colorado. Recovery implementation program for endangered fish species in the Upper Colorado River Basin. U.S. Department of the Interior, Fish and Wildlife Service, Region 6, Denver, Colorado.
- Burdick, B.D. 1994b. Conceptual management plan for habitat enhancement in flooded bottomlands: Gravel Pit at 29-5/8 Road, Colorado River near Grand Junction, Colorado. Recovery implementation program for endangered fish species in the Upper Colorado River Basin. U.S. Department of the Interior, Fish and Wildlife Service, Region 6, Denver, Colorado.
- Collins, K. 1994. Conceptual management plan for habitat enhancement in flooded bottomlands: Scott M. Matheson wetland preserve, Moab, Utah. Recovery implementation program for endangered fish species in the Upper Colorado River Basin. U.S. Department of the Interior, Fish and Wildlife Service, Region 6, Denver, Colorado.
- Cooper, D.J. and C. Severn. 1994a. Ecological characteristics of wetlands at the Moab Slough, Moab, Utah. Recovery implementation program, Upper Colorado River Basin. U.S. Department of the Interior, Fish and Wildlife Service, Denver, Colorado. 121 pp.
- Cooper, D.J. and C. Severn. 1994b. Evaluation of the 29-5/8 Mile Pond near Grand Junction, Colorado. Recovery implementation program, Upper Colorado River Basin. U.S. Department of the Interior, Fish and Wildlife Service, Denver, Colorado. 5 pp.
- Cooper, D.J. and C. Severn. 1994c. Wetlands of the Escalante State Wildlife Area on the Gunnison River, near Delta, Colorado: Hydrology, water chemistry, vegetation, invertebrate communities, and restoration potential. Recovery implementation program, Upper Colorado River Basin. U.S. Department of the Interior, Fish and Wildlife Service, Denver, Colorado. 94 pp.
- Cooper, D.J. and C. Severn. 1994d. Wetlands of the Escalante Ranch area, Utah: Hydrology, water chemistry, vegetation, invertebrate communities, and restoration potential. Recovery implementation program, Upper Colorado River Basin. U.S. Department of the Interior, Fish and Wildlife Service, Denver, Colorado. 97 pp.

- Cooper, D.J. and C. Severn. 1994e. Wetlands of the Ouray National Wildlife Refuge, Utah: Hydrology, water chemistry, vegetation, invertebrate communities, and restoration potential. Recovery implementation program, Upper Colorado River Basin. U.S. Department of the Interior, Fish and Wildlife Service, Denver, Colorado. 79 pp.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. Office of Biological Services, U.S. Fish and Wildlife Service, U.S. Department of the Interior, Washington, D.C.
- Grabowski, S.J. and S.D. Hiebert. 1989. Some aspects of tropic interactions in selected backwaters and the main channel of the Green River, Utah: 1987-1988. U.S. Bureau of Reclamation, Research and Laboratory Services Division, Environmental Sciences Section, Denver, Colorado. 130 pp. + an appendix of 155 pp.
- Graf, W.L. 1978. Fluvial adjustments to the spread of tamarisk on the Colorado Plateau region. Bulletin of the Geological Society of America 89: 1491-1501.
- Lyon, J.G. 1993. Practical Handbook for Wetland Identification and Delineation. Lewis Publishers. Florida, USA.
- Mabey, L.W. and D.K. Shiozawa. 1993. Planktonic and benthic micro-crustaceans from floodplain and river habitats of the Ouray Refuge on the Green River, Utah. Department of Zoology, Brigham Young University, Provo, Utah. 31 pp.
- Marsh, P.C. and D.R. Langhorst. 1988. Feeding and fate of wild larval razorback sucker. Environmental Biology of Fishes 21: 59-67.
- McAda, C.W. 1977. Aspects of the life history of three catostomids native to the Upper Colorado River basin. M. S. Thesis, Utah State University, Logan, Utah. 117 pp.
- Minckley, W.L., P.C. Marsh, J.E. Brooks, J.E. Johnson, and B.L. Jensen. 1991. Management toward recovery of the razorback sucker. Pages 303-358 in W.L. Minckley and J.E. Deacon eds. Battle against extinction. University of Arizona Press, Tucson, AR.
- Modde, T. and D. Irving. 1994. Conceptual management plan for habitat habitat enhancement in flooded bottomlands: Old Charlie Wash. Recovery implementation program for endangered fish species in the Upper Colorado River Basin. U.S. Department of the Interior, Fish and Wildlife Service, Region 6, Denver, Colorado.
- Osmundson, D.B. and L.R. Kaeding. 1991. Recommendations for flows in the 15-mile reach during October-June for maintenance and enhancement of endangered fish populations in the Upper Colorado River. Colorado River Fishery Project, U.S. Department of the Interior, Fish and Wildlife Service, Grand Junction, Colorado. 82 pp.

- Papoulias, D. and W.L. Minckley. 1990. Food limited survival of larval razorback sucker, *Xyrauchen texanus*, in the laboratory. *Environmental Biology of Fishes* 29: 73-78.
- Quarterone, F. 1993. Historical accounts of Upper Colorado River basin endangered fish. Colorado Division of Wildlife, Denver, 66 pages.
- Richardson, C. 1994. Conceptual management plan for habitat enhancement in flooded bottomlands: Escalante Ranch, Jensen Utah. Recovery implementation program for endangered fish species in the Upper Colorado River Basin. U.S. Department of the Interior, Fish and Wildlife Service, Region 6, Denver, Colorado.
- The Nature Conservancy. 1992. Proposal for developing a riparian wetland community type classification for the White and Colorado River watersheds, within the Upper Colorado River Basin in Colorado. Submitted to: U.S. Environmental Protection Agency, Denver, Colorado.
- Tyus, H.M. and C.A. Karp. 1989. Habitat use and streamflow needs of rare and endangered fishes, Yampa River, Colorado. U.S. Fish and Wildlife Service, Biological Report 89(14). 27 pp.
- U.S. Fish and Wildlife Service. 1987. Recovery implementation program for endangered fish species in the Upper Colorado River Basin. U.S. Department of the Interior, Fish and Wildlife Service, Region 6, Denver, Colorado. 6 Sections, Various pagination.
- Wydoski, R.S. and E.D. Wick. 1994. Enhancement strategies for flooded bottomlands and other habitats in the recovery of endangered fishes in the Upper Colorado River basin. Draft report. U. S. Department of Interior, Fish and Wildlife Service and National Park Service, Denver, Colorado. 80 pp + Appendices.

APPENDIX A
The Floodplain Habitat Restoration Program,
Commonly Asked Questions, and
Right-of-Entry Permit Form



RECOVERY PROGRAM FOR THE ENDANGERED FISHES OF THE UPPER COLORADO

John Hamill,
Program Director

Ralph Morgenweck,
Implementation
Committee Chairman

U.S. Fish and Wildlife Service • P.O. Box 25486 • Denver Federal Center • Denver, Colorado 80225 • (303) 236-2985

The Flood Plain Habitat Restoration Program

The Flood Plain Habitat Restoration Program is a component of the Recovery Program for Endangered Fish of the Upper Colorado River Basin. The purpose of the Recovery Program is to recover endangered fishes while allowing water development to continue (see "Swimming Upstream" brochure). The Flood Plain Habitat Restoration Program seeks to restore riverside habitats to benefit endangered fishes of the Green, Colorado, and Gunnison rivers within Utah and Colorado. Biologists believe that young razorback suckers, and possibly bonytail chubs, rely on annual flooding to survive and grow. Habitat for these native fish is disappearing, however, because dams and levees have reduced flooding. As a result, species of native fishes that rely on flooding have declined, and some are close to extinction.

As an owner of riverfront property, you can help to prevent these fishes from becoming extinct. With your permission, our scientists would like to collect data on your property to determine its potential for providing habitat for endangered fishes. If your land is suitable for restoring fish habitat, we would like to work cooperatively with you to manage a portion of your property for the endangered fishes. This may require some initial excavation and construction, followed by some spring and summer flooding. Although the program is entirely voluntary, money is available for these cooperative projects. If you choose not to participate, then we will continue to search elsewhere for habitats that can be managed for endangered fishes.

If you desire additional information, please contact:

Patrick C. Nelson
Flood Plain Habitat Restoration Program Coordinator
U.S. Fish and Wildlife Service
P.O. Box 25486, Denver Federal Center
Denver, Colorado 80225
Phone (303) 236-2985 Extension 226

Commonly-Asked Questions

Why do you want to collect data on my property?

Scientists from the Flood Plain Habitat Restoration Program are evaluating areas along the river which may have potential to assist in recovery of endangered fishes. Your property has been identified as one that may provide habitat for the endangered fishes.

If I give permission for scientists to collect data, how will I be affected, and how will my property be affected?

The scientists will take every precaution to avoid disturbing you, your family, and your property. We will leave it up to you to decide when scientists are allowed to collect data on your property, which access routes they can take, and whether or not they should call you before they visit your property. The scientists will take some photos and measurements, install (and later remove) a temporary gauge, and collect water, sediment, fish, and vegetation samples. There will be no noticeable evidence of this work.

How many scientists will be involved, and for how long?

There will be several scientists involved, as many as five. Some of the scientists may be accompanied by one or more assistants. Each scientist will require from one to four days to collect the necessary data. Scientists will call you before they visit your property, if you choose.

What if the scientists leave a gate open and my livestock get out, or if their truck gets stuck and they dig up my property trying to get out, or if they damage my property in some other way? Will I be paid for damages?

If your property is damaged by the government or its employees, you will be fairly compensated for the damages.

What if one of the scientists is injured while working on my property? Will I be held liable?

No. Government employees that are injured during the performance of their duties are covered by workman's compensation. You will not be held liable.

If I cooperate, what is in it for me?

If the scientists determine that your land has potential for use as endangered species' habitat, then you will be invited to participate in a cooperative agreement. A real estate specialist will visit with you to develop an agreement that is acceptable to you. If you accept, you will be compensated for your participation. Also, you will have the satisfaction of knowing that you are contributing towards recovery of endangered species.

How much compensation will I receive?

The amount of compensation you receive will depend on the size and value of your property and the type of agreement developed. A real estate specialist will be available to explain some of the options to you.

Can I refuse to participate?

Yes. If a property owner chooses not to participate, then we will look elsewhere for riverside habitats to manage for endangered species. You are under no obligation to participate.

We wish to extend our thanks to you for your consideration and participation in this very important work. If you have additional questions, you may write or call:

Patrick C. Nelson
Flood Plain Habitat Restoration Coordinator
U.S. Fish and Wildlife Service
P.O. Box 25486, Denver Federal Center
Denver, Colorado 80225
Phone (303) 236-2985 Extension 226

FLOOD PLAIN HABITAT RESTORATION PROGRAM

Right-of-Entry Permit

_____ of _____ COUNTY, STATE
OF UTAH, does hereby grant the Bureau of Reclamation and its authorized agents
or assigns permission to enter upon the following described property:

For the purpose of conducting investigations necessary to determine the suitability of the property for the Flood Plain Habitat Restoration Program. These investigations may include, but are not limited to, the following activities: surveying, collecting data or samples of soil, water, vegetation, sediment, and fish, and the installation of temporary gauges.

This agreement shall be in effect upon execution thereof, and shall remain in effect until _____.

The Bureau of Reclamation agrees to:

1. Pay for all damages to the land, crops, livestock, and personal property situated thereon caused through exercise of the rights granted herein based upon an appraisal made by the Bureau of Reclamation, or by mutual agreement between the parties hereto.
2. Keep all gates closed and not interfere with any livestock or irrigation practices.
3. Restore the land to as nearly its original condition as practicable, upon completion of the described work.

It is understood and agreed between the parties that the owner(s) will not be held responsible for any personal injury to employees of the United States, its agents, assigns, and contractors, nor for any damage to their personal property.

Landowner _____ Date ACCEPTED:
United States Bureau of Reclamation

Landowner _____ Date BY: _____ Date

APPENDIX B
Existing and needed inventory maps and photos

Table B.1. Existing and needed inventory maps, aerial photographs, infrared photographs, and videos of bottomland habitat in the Green River Basin, Colorado and Utah, 1993.

DRAINAGE	EXISTING INVENTORY	INVENTORY NEEDS
Green R.	Colored infrared aerial photographs, shot at 1 in. = 6,000 ft. from Flaming Gorge Dam to the Green/Colorado River confluence in the fall of 1974 by the U.S. Army Corps of Engineers, SLC, Ut. Colored aerial videography, shot at various altitudes from Flaming Gorge dam to Jensen and them at Ouray during the May and June 1992 by Argonne National Laboratories, Bowling Brook, Ill. Various maps from the early 1900's to present by the U.S. Bureau of Reclamation, SLC, Ut.	Colored infrared aerial photographs, shot at 1 in. = 10,000 ft. from Flaming Gorge Dam to the Green/Colorado River confluence in at high (May-June) and low (July/August) discharge. The existing aerial photographs do not provide the needed information because they were shot at 1:6,000 only include the main river channel and not the adjacent bottomland areas.
Yampa R.	Colored infrared aerial photographs, shot at 1 in. = 6,000 ft. of the entire river in late 1970's or early 1980's by the U.S. Army Corps of Engineers, Grand Junction, Co. Various maps from the early 1900's to present by the U.S. Bureau of Reclamation, Denver, Co.	Colored infrared aerial photographs, shot at 1 in. = 10,000 ft. from Flaming Gorge Dam to the Green/Colorado River confluence in at high (May-June) and low (July/August) discharge. The existing aerial photographs do not provide the needed information because they were shot at 1:6,000 only include the main river channel and not the adjacent bottomland areas.
White R.	Colored infrared aerial photographs, shot at 1 in. = 6,000 ft. of the entire river in late 1970's or early 1980's by the U.S. Army Corps of Engineers, Grand Junction, Co. Various maps from the early 1900's to present by the U.S. Bureau of Reclamation, Denver, Co.	Colored infrared aerial photographs, shot at 1 in. = 10,000 ft. from Flaming Gorge Dam to the Green/Colorado River confluence in at high (May-June) and low (July/August) discharge. The existing aerial photographs do not provide the needed information because they were shot at 1:6,000 only include the main river channel and not the adjacent bottomland areas.

APPENDIX C

Complete and partial inventory plan for the
Green and Colorado rivers, 1993

Table C.1. Plan for complete and partial infrared photograph inventories for the Green River, Colorado and Utah, 1993.

PLAN A: Complete Inventory	PLAN B: Partial Inventory
<u>Green River:</u> Take colored infrared aerial photographs of the Green River from Flaming Gorge Dam to the Green/Colorado River confluence (410 rm [659.8 rkm]). Shoot at 1 in. (2.5 cm) = 10,000 ft. (3,048 m) during high (May-June) and low (July-August) discharge. The cost to shoot this area would be about \$28,700 (\$70/mile [\$43.5/km]). <u>Yampa River:</u> Take colored infrared aerial photographs of the Yampa River from Cross Mountain Canyon to the Yampa/Green River confluence (47 rm [75.6 rkm]). Shoot at 1 in. (2.5 cm) = 10,000 ft. (3,048 m) during high (May-June) and low (July-August) discharge. The cost to shoot this area would be about \$3,290 (\$70/mile [\$43.5/km]). <u>White River:</u> Take colored infrared aerial photographs of the White River from Taylor Draw Dam to the White/Green River confluence (104 rm [167.4 rkm]). Shoot at 1 in. (2.5 cm) = 10,000 ft. (3,048 m) during high (May-June) and low (July-August) discharge. The cost to shoot this area would be about \$7,280 (\$70/mile [\$43.5/km]).	<u>Green River:</u> Take colored infrared aerial photographs of the Green River at specific bottomland reaches. Shoot at 1 in. (2.5 cm) = 10,000 ft. (3,048 m) during high (May-June) and low (July-August) discharge. The cost is \$70/mile (\$43.5/km). Brown's Park (rm 396 [rkm 637.3]) - Island Park (rm 333 [rkm 535.9]), a distance of 63 rm (101.4 rkm) and a cost of about \$4,410. Split Mountain (rm 320 [rkm 515.0]) - Horseshoe Bend (rm 280 [rkm 450.6]), a distance of 40 rm (64.4 rkm) and a cost of about \$2,800. Split Mountain (rm 320 [rkm 515.0]) - Sand Wash (rm 216 [rkm 347.6]), a distance of 104 rm (167.4 rkm) and a cost of about \$7,280. Horseshoe Bend (rm 280 [rkm 450.6]) - Ouray Refuge (rm 258 [rkm 415.2]), a distance of 22 rm (35.4 rkm) and a cost of about \$1,540. Ouray Refuge (rm 258 [rkm 415.2]) - Sand Wash (rm 216 [rkm 347.6]), a distance of 42 rm (67.6 rkm) and a cost of about \$2,940. Green River (rm 120 [rkm 193.1]) - Green/Colorado River Confluence (rm 0), a distance of 120 rm (193.1 rkm) and a cost of about \$8,400. <u>Yampa River:</u> The same as Plan A. <u>White River:</u> the same as Plan A.

APPENDIX D

**Bottomland habitat sites, river miles,
historical and actual flooded acreages**

Table D.1. Bottomland habitat sites, river mile, and actual and potential inundated acres (hectares), Green River, Colorado and Utah, May and September 1993.

Site Description	River Mile	05/25/93		09/28/93		Potential Area	
		Acres	Hectares	Acres	Hectares	Acres	Hectares
GREEN RIVER							
Sears Cr.-Allen Ranch, BPNWR	388-392.5	215.8	87.3	183.2	74.1	215.8	87.3
Crouse Creek BPNWR	381.5-382.5	70.0	28.3	39.3	15.9	70.0	28.3
Butch Cassidy lake BPNWR	379.5-380.5	90.8	36.7	0.0	0.0	90.8	36.7
Hog Lake BPNWR	376.5-378.0	213.0	86.2	59.7	24.2	350.0	141.6
Warren Bottom, BPNWR	375.5-377.0	316.4	128.0	203.2	82.2	320.0	129.5
Spitzie Bottom, Browns Park NWR	375.0-376.0	101.5	41.1	116.5	47.1	172.0	69.6
J S Hoy Bottom, Browns Park NWR	372.0-374.0	323.4	130.9	137.2	55.5	323.4	130.9
Grimes Bottom BPNWR	367.5-369.5	40.0	16.2	34.2	13.8	50.0	20.2
Escalante Ranch	302.5-309.5	148.5	60.1	51.9	21.0	750.1	303.6
Meril Snow Ranch	302.0-303.0	81.9	33.1	1.3	0.5	136.4	55.2
Gravel Ponds at Jensen	301.0-302.0	10.2	4.1	9.9	4.0	10.2	4.1
Stewart Lake	299.0-300.0	569.0	230.3	160.0	64.7	663.7	268.6
Ashley Creek confluence area	297.0-298.5	92.4	37.4	0.0	0.0	161.6	65.4
Little Stewart Lake	295.5-297.5	131.5	53.2	143.3	58.0	263.7	106.7
Spring Hollow	295.0-296.0	19.0	7.7	9.4	3.8	200.6	81.2
Walker Hollow	294.0-295.0	22.2	9.0	13.1	5.3	62.0	25.1
Alhandra Ferry Site	292.0-294.0	18.0	7.3	1.9	0.8	205.2	83.0
Gravel Pits	292.0-293.0	4.7	1.9	1.2	0.5	4.7	1.9
Bonanza Bridge Area	288.5-289.0	12.3	5.0	0.0	0.0	52.5	21.2
Collier Draw	285.5-286.5	18.7	7.6	0.0	0.0	44.0	17.8
Horseshoe Bend Area	277.0-284.0	85.0	34.4	14.1	5.7	239.0	96.7
The Stirrup	274.0-277.0	8.3	3.4	0.0	0.0	10.8	4.4
Hamacker Bottom/Baeser Bend	271.0-274.0	315.9	127.8	4.0	1.6	549.3	222.3
Downstream of Baeser Bend	269.0-272.0	42.9	17.4	0.0	0.0	68.7	27.8
Upstream of Brennan Bottom	267.0-269.0	36.9	14.9	32.4	13.1	36.9	14.9
Upstream of Brennan Bottom	266.0-267.0	8.8	3.6	0.0	0.0	32.0	12.9
Brennan Bottom	262.0-266.0	99.9	40.4	0.0	0.0	374.6	151.6
Johnson Bottom, Ouray NWR	263.0-265.0	257.8	104.3	207.6	84.0	257.8	104.3
Leota Pond Complex, Ouray NWR	257.0-262.0	865.1	350.1	99.6	40.3	1,351.1	546.8
Wyasket Lake, Ouray NWR	253.0-257.0	533.1	215.7	778.6	315.1	888.9	359.7
Sheppard Bottom, Ouray NWR	254.0-256.0	245.6	99.4	0.0	0.0	720.0	291.4
Main Old Charlie (Woods) Bottom, Ouray NWR	249.0-252.0	198.4	80.3	13.8	5.6	302.2	122.3
Diked Old Charlie (Woods) Bottom, Ouray NWR	249.0-250.0	164.7	66.7	12.7	5.1	277.3	112.2
Ouray Ute pasture land	248.0-251.0	431.7	174.7	0.0	0.0	731.2	295.9
Duchesne River confluence area	248.0-249.0	134.3	54.3	3.0	1.2	250.4	101.3
White River confluence area	248.0-249.0	496.4	200.9	0.0	0.0	614.7	248.8
West Branch area	243.0-247.0	615.1	248.9	37.4	15.1	880.9	356.5
Tia Juana Bottom	242.0-244.0	319.3	129.2	64.6	26.1	625.1	253.0
Mouth of Willow Creek	239.0-241.0	0.0	0.0	0.0	0.0	156.3	63.3
Pariette Draw	238.0-241.0	105.8	42.8	0.0	0.0	343.1	138.8
Desert Spring/Moon Bottom	227.0-228.0	10.0	4.0	0.0	0.0	10.0	4.0
Fourmile Wash/King Bottom	226.0-227.0	0.0	0.0	0.0	0.0	11.7	4.7
Indian Pasture/Hydes Bottom	224.0-225.0	30.8	12.5	0.0	0.0	65.0	26.3
Rays and Long Bottom	220.0-222.0	25.0	10.1	0.0	0.0	45.4	18.4
Between Long and Boat Bottom	219.0-220.0	5.0	2.0	0.0	0.0	11.4	4.6
Nine Mile Creek	213.0-214.0	6.6	2.7	0.0	0.0	116.1	47.0
Nutter Hole	211.0-213.0	0.0	0.0	0.0	0.0	56.1	22.7
Duches Hole	209.0-211.0	0.0	0.0	0.0	0.0	90.0	36.4
Tabyago Canyon/Little Horse	206.0-208.0	0.0	0.0	0.0	0.0	22.9	9.3
Little Horse Bottom	204.0-207.0	0.0	0.0	0.0	0.0	85.2	34.5
Downstream Maverick Canyon	203.0-204.0	0.0	0.0	0.0	0.0	147.6	59.7
Gold Hole	201.0-202.0	29.1	11.8	0.0	0.0	46.7	18.9
Rock House Bottom	200.0-201.0	2.2	0.9	0.0	0.0	80.7	32.7
Little Rock Canyon	199.0-200.0	8.2	3.3	0.0	0.0	61.6	24.9
Stampede Flat	198.0-199.0	0.0	0.0	0.0	0.0	64.6	26.1
Hoodoo Forms	197.0-198.0	0.0	0.0	0.0	0.0	184.7	74.7
Between Snap-Three Canyons	170.0-171.0	0.0	0.0	0.0	0.0	20.6	8.3
Chandler Falls Canyon	165.0-167.0	0.0	0.0	0.0	0.0	33.4	13.5
Trail and Bull Canyon Rapids	164.5-165.5	0.0	0.0	0.0	0.0	23.6	9.6
Joe Hutch Canyon/Rapids	160.0-161.0	0.0	0.0	0.0	0.0	16.8	6.8
Florence Creek/Three Fords	157.0-158.0	0.0	0.0	0.0	0.0	11.0	4.5
Downstream of Three Fords	153.5-155.0	0.0	0.0	0.0	0.0	16.3	6.6
Last Chance Canyon/Rapids	152.0-153.0	0.0	0.0	0.0	0.0	51.2	20.7
Range Creek Rapids	150.5-152.0	0.0	0.0	0.0	0.0	10.7	4.3
Little Big Horn Mesa	143.0-144.0	0.0	0.0	0.0	0.0	14.0	5.7
Price River confluence	137.0-138.5	0.0	0.0	0.0	0.0	23.7	9.6
Downstream of Price River	136.5-137.5	0.0	0.0	0.0	0.0	14.1	5.7
Short Canyon/Rapids	131.5-132.5	0.0	0.0	0.0	0.0	4.3	1.7
Willow Bend-Tusher Rapid	128.5-129.5	0.0	0.0	0.0	0.0	152.2	61.6
Upriver of Green River Utah	121.5-125.5	0.0	0.0	0.0	0.0	513.9	208.0
Little Grand Wash	114.5-115.5	0.0	0.0	0.0	0.0	23.0	9.3
Downstream of Grand Wash	113.0-114.0	0.0	0.0	0.0	0.0	62.2	25.2
Fivemile Wash/Little Valley	111.0-112.0	0.0	0.0	0.0	0.0	15.0	6.1
Downstream of Ninemile Wash	109.0-111.0	0.0	0.0	0.0	0.0	20.5	8.3
Anvil Bottom	101.5-102.5	0.9	0.4	0.0	0.0	9.7	3.9
Upstream of San Rafael River	98.0-99.0	0.0	0.0	0.0	0.0	26.4	10.7
Sand Rafael River confluence	96.5-97.5	1.0	0.4	0.0	0.0	44.0	17.8
White and Red Wash	95.0-96.5	3.9	1.6	0.0	0.0	16.5	6.7
Between Red Wash-Bull Bottom	94.0-95.5	0.0	0.0	0.0	0.0	38.7	15.7
Bull Bottom-Labyrinth Canyon	92.0-93.0	0.0	0.0	0.0	0.0	33.4	13.5
Labyrinth Canyon confluence	91.5-92.5	0.0	0.0	0.0	0.0	36.3	14.7
Three Canyon/Trin-Alcove Bend	88.5-91.5	0.0	0.0	0.0	0.0	65.2	26.4
Junes Bottom	87.0-88.0	0.0	0.0	0.0	0.0	35.9	14.5
Junes Bottom-Bull Hollow	85.0-86.0	0.0	0.0	0.0	0.0	36.5	14.8
Bull Hollow	84.0-85.0	1.7	0.7	0.0	0.0	23.4	9.5
Bull Hollow-Tenmile Canyon	81.0-83.0	0.0	0.0	0.0	0.0	37.8	15.3
Tenmile Canyon/Bottom	79.0-82.0	10.2	4.1	0.0	0.0	89.7	36.3
Keg Springs Canyon	75.0-78.5	0.0	0.0	0.0	0.0	48.1	19.5
Hey Joe Canyon	74.5-76.0	0.6	0.2	0.0	0.0	25.0	10.1
Hey Joe Canyon-Spring Canyon	74.0-75.0	0.0	0.0	0.0	0.0	33.6	13.6

Table D.1. Continued.

Site Description	River Mile	05/25/93		09/28/93		Potential Area	
		Actual Area Acres	Hectares	Actual Area Acres	Hectares	Acres	Hectares
<u>GREEN RIVER</u>							
Spring Canyon Point	69.5-74.0	0.0	0.0	0.0	0.0	89.3	36.1
Bowknot Bend	62.0-70.0	13.6	5.5	0.0	0.0	261.1	105.7
Twomile Canyon-Deadman Point	60.0-62.0	4.1	1.7	0.0	0.0	36.4	14.7
Deadman Point-Horseshoe Canyon	59.0-60.0	0.0	0.0	0.0	0.0	48.3	19.5
Downstream of Horseshoe Canyon	58.0-59.0	0.0	0.0	0.0	0.0	50.2	20.3
Cottonwood Bottom	55.0-56.5	1.8	0.7	0.0	0.0	64.3	26.0
Cottonwood-Mineral Bottoms	54.0-55.0	0.0	0.0	0.0	0.0	43.6	17.6
Mineral-Tidwell Bottoms	53.0-54.0	0.0	0.0	0.0	0.0	74.3	30.1
Tidwell Bottom	51.0-52.5	0.0	0.0	0.0	0.0	162.9	65.9
Horsethief Bottom	50.0-51.5	0.0	0.0	0.0	0.0	32.1	13.0
Woodruff Bottom	49.0-50.0	0.0	0.0	0.0	0.0	81.1	32.8
Point Bottom	47.5-49.5	0.0	0.0	0.0	0.0	32.7	13.2
Saddle Horse Bottom	45.5-47.5	0.0	0.0	0.0	0.0	14.0	5.7
Horsethief Canyon	45.0-46.0	0.0	0.0	0.0	0.0	69.6	28.2
Horsethief-Upheaval Bottom	44.0-45.5	0.0	0.0	0.0	0.0	104.7	42.4
Upheaval Bottom/Canyon	43.5-44.5	1.4	0.6	0.0	0.0	78.3	31.7
Hardscrabble Bottom	42.5-43.5	0.0	0.0	0.0	0.0	30.7	12.4
Hardscrabble-Fort Bottoms	38.5-42.5	1.9	0.8	0.0	0.0	156.2	63.2
Potato Bottom	35.5-38.0	0.0	0.0	0.0	0.0	76.8	31.1
Potato-Beaver Bottoms	34.5-35.5	0.0	0.0	0.0	0.0	44.2	17.9
Beaver-Queen Ann Bottoms	33.5-34.5	5.1	2.1	0.0	0.0	49.1	19.9
Queen Ann Bottom	33.0-34.0	2.4	1.0	0.0	0.0	68.8	27.8
Queen Ann-Anderson Bottoms	32.0-33.0	0.0	0.0	0.0	0.0	15.0	6.1
Anderson Bottom	30.0-31.5	2.1	0.8	4.8	1.9	23.3	9.4
Unknown and Valentine Bottoms	29.0-32.0	0.0	0.0	0.0	0.0	112.2	45.4
Valentine Bottom	27.0-29.0	1.3	0.5	0.0	0.0	117.8	47.7
Stillwater Canyon-Sphinx	26.0-28.5	17.7	7.2	0.0	0.0	96.3	39.0
Downstream Valentine Bottom	25.5-26.5	45.5	18.4	0.0	0.0	67.6	27.4
Valentine-Tuxedo Bottoms	24.0-25.5	0.0	0.0	0.0	0.0	96.9	39.2
Tuxedo Bottom-Turks head	19.5-24.0	4.0	1.6	0.0	0.0	235.9	95.5
Deadhorse Canyon confluence	18.5-20.0	1.6	0.6	0.0	0.0	26.3	10.6
Downstream Deadhorse Canyon	17.0-19.0	2.6	1.1	0.0	0.0	57.8	23.4
Deadhorse-Horse Canyons	16.0-17.0	0.0	0.0	0.0	0.0	21.3	8.6
Horse Canyon	13.5-14.0	7.4	3.0	0.0	0.0	51.7	20.9
Downstream of Horse Canyon	12.5-13.5	0.0	0.0	0.0	0.0	32.6	13.2
Horse-Jasper Canyons	10.5-12.5	2.3	0.9	0.0	0.0	60.4	24.4
Jasper Canyon	8.0-10.5	1.8	0.7	0.0	0.0	55.8	22.6
Jasper Canyon-Short Canyon	5.0-8.0	0.6	0.2	0.0	0.0	33.5	13.6
Short Canyon-Colorado River	2.0-3.5	2.0	0.8	0.0	0.0	13.8	5.6
Upstream of Colorado River	1.0-2.0	0.6	0.2	0.0	0.0	14.8	6.0
Near Colorado River confluence	0.5-1.5	0.5	0.2	0.0	0.0	29.0	11.7
Colorado River confluence	0.0-0.5	0.0	0.0	0.0	0.0	15.6	6.3
sample statistics for all sites	n	132.0	132.0	132.0	132.0	132.0	132.0
	min	0.0	0.0	0.0	0.0	4.3	1.7
	mean	58.5	23.7	18.5	7.5	139.6	56.5
	max	865.1	350.1	778.6	315.1	1,351.1	546.8
	sum	7,719.8	3,124.1	2,437.9	986.6	18,429.7	7,458.2
sample statistics for inundated sites	n	74.0	74.0	28.0	28.0	132.0	132.0
	min	0.5	0.2	1.2	0.5	4.3	1.7
	mean	104.3	42.2	87.1	35.2	139.6	56.5
	max	865.1	350.1	778.6	315.1	1,351.1	546.8
	sum	7,719.8	3,124.1	2,437.9	986.6	18,429.7	7,458.2
<u>YAMPA RIVER</u>							
Upstream Snake River confluence	48.0-48.5	21.9	8.9	0.0	0.0	21.0	8.9
<u>WHITE RIVER</u>							
Oxbow downstream Highway 65	91.0-93.0	18.9	7.6	17.6	7.1	18.9	7.6
White/Green river confluence	0.0-2.5	496.4	200.9	0.0	0.0	614.7	248.8
	n	2.0	2.0	2.0	2.0	2.0	2.0
	min	18.9	7.6	0.0	0.0	18.9	7.6
	mean	257.7	104.3	8.8	3.6	316.8	128.2
	max	496.4	200.9	17.6	7.1	614.7	248.8
	sum	515.3	208.5	17.6	7.1	633.6	256.4

Table D.2. Bottomland habitat sites, river mile, actual (May [runoff] and September [post-runoff] 1993) inundated area (hectares), and historical inundated area (hectares), Colorado and Gunnison rivers, Colorado and Utah, May and September 1993.

Site Description	River Mile	Actual Inundated Area (ha)		Historical Inundated Area (ha)
		Runoff (May 1993)	Post-Runoff (September 1993)	
<u>Colorado River</u>				
Rifle West	238.0-241.1	12.2	12.1	150.7
Rifle I-70 West Interchange	236.5-238.0	5.3	5.3	126.4
CDOW/Gentry Property	234.2-235.8	10.4	10.2	66.8
"Trash-can Pond"	230.2-230.8	5.3	1.8	5.8
Mahaffey/Lemon	228.2-229.8	6.5	2.3	54.4
Mahaffey's	228.5-229.5	2.5	2.2	30.5
Hoaglund's	227.1-227.7	1.2	< 0.1	16.4
Dere/Ortiz	227.5-228.5	6.1	6.0	39.3
Parachute East	223.0-223.4	2.1	0.7	13.7
Knight's	224.5-225.4	4.4	0	28.2
Parachute Bridge	223.0-223.4	2.1	0.7	13.7
"No-name"	222.7-223.1	0.4	0.6	8.6
"No-name"	221.1-222.9	11.5	11.3	23.4
Battlement Mesa	220.7-221.7	10.2	3.9	34.4
EXXON	218.9-220.1	3.1	0	34.9
Doyle Property	218.4-219.5	0	0	8.7
"No-name"	217.2-218.0	1.4	0	22.3
Una North	216.4-217.4	6.4	6.3	40.7
Wallace Creek Island	215.9-216.5	3.4	0	13.3
Stoddard Property	209.4-210.1	8.1	8.1	23.2
Debeque I-70 Slough	209.6-211.4	6.7	3.6	40.4
Latham's	207.9-208.8	6.2	1.6	26.1
"No-name"	206.2-207.5	2.1	0	22.9
Etter's	204.7-206.6	22.8	22.8	65.1
"No-name"	202.3-204.2	6.9	3.3	86.4
"No-name"	201.9-202.4	0.2	0	6.1
Long Point Bottom	198.0-198.9	2.3	0.2	10.6
Beavertail	194.3-195.6	2.0	1.8	12.6
Island Acres	191.1-192.1	11.4	11.4	28.1
Cameo	190.0-191.0	1.2	1.2	24.6
Palisade	184.2-185.0	2.5	2.1	22.0
"No-name"	183.3-184.3	3.8	1.3	17.2
Labor Camp	182.9-183.6	8.3	2.7	19.6
"No-name"	181.6-182.6	0.6	0	43.3
"No-name"	181.7-182.2	2.8	2.2	6.9
"No-name"	180.4-181.5	4.2	1.4	21.2
Clifton Water Treatment	179.1-181.1	20.7	17.8	122.5
Clifton Pond	177.7-178.2	21.7	15.6	81.7
Corn Lake	176.6-177.7	14.5	8.3	38.6
Humphrey's	175.3-176.7	47.9	25.0	86.2
Griffith's	174.1-176.5	24.2	2.6	52.7
"Hotspot Junction" (30-29 Road)	173.9-175.1	11.3	9.4	97.9
"No-name"	173.4-173.8	1.4	0.6	10.6
"No-name"	173.0-173.6	0	0	7.8
Mill Tailing Site	172.5-173.4	1.9	< 0.1	40.6
Watson Island	171.0-172.1	9.3	4.6	20.6
"No-Name"	171.3-171.8	3.0	0.4	5.9
"No-name"	170.4-171.0	11.2	0	16.4
Connected Lakes Area	168.7-170.3	77.2	59.3	220.6

Table D.2. Continued.

Site Description	River Mile	Actual Inundated Area (ha)		Historical Inundated Area (ha)
		Runoff (May 1993)	Post-Runoff (September 1993)	
<u>Colorado River</u>				
Walter Walker South	164.4-166.0	28.7	3.0	60.3
"No-name"	166.4-169.7	30.7	20.3	117.0
Appleton Drain East	165.1-166.4	19.5	17.0	61.1
Walter Walker SWA	162.7-165.1	17.6	4.7	144.5
Panorama	163.1-163.6	3.1	2.8	20.6
"No-name"	161.0-162.7	2.5	0.9	20.6
DuPont Island	161.0-162.0	16.8	3.6	73.9
DuPont's	159.1-161.9	37.5	25.2	86.6
Paul Smith's	158.0-159.1	13.8	2.4	53.4
Fruita Sewage Ponds	156.6-158.0	16.6	16.5	40.9
Fruita 340 Bridge	157.5-158.3	0.8	0.4	17.0
Snook's Bottom	155.9-157.1	1.4	0.6	38.0
Horsethief SWA	151.9-154.7	4.9	1.5	180.9
Spann's	151.4-152.4	5.2	4.7	34.0
"No-name"	150.8-151.5	0	0	27.5
Crow Bottom	143.9-146.5	0	0	96.0
"No-name"	145.5-146.6	0	0	39.7
Vulture Bottom	139.7-142.1	0	0	61.5
"No-name" Island	137.2-138.0	0	0	23.2
Black Rocks	136.7-137.1	0.6	0	17.8
Knowles Canyon	133.6-135.0	1.0	0	52.0
Joufflas Bottom	129.8-133.9	0	0	99.6
Wildass Canyon Ranch	127.3-131.3	14.6	1.9	189.0
"No-name"	125.8-126.8	0.8	0	54.0
Elizondo Ranches	126.5-127.8	4.0	1.7	87.1
Westwater Wash	124.8-126.0	16.8	0.2	84.0
Cisco Landing Area	107.6-111.6	4.0	0	267.0
Fish Ford Area	103.0-105.9	10.6	0	100.5
McGraw/Hotel Bottom	98.1-101.0	24.7	?	123.7
White Ranch	77.5-78.4	0	0	26.1
Courthouse Wash	63.8	1.6	0	2.8
Moab Slough	61.5-64.0	212.0	81.0	354.0
Kane Spring	58.2	1.6	0	2.8
Billboard (Lake Bottom)	50.9-52.5	9.9	0	36.0
Jackson Bottom	47.4-49.0	13.8	0	34.0
"No-name"	44.0-44.7	11.4	0	23.0
"No-name"	42.8-43.7	4.8	0	22.0
"No-name"	41.1-42.1	10.7	0	22.3
"No-name"	40.2-41.3	0.4	0	21.0
"No-name"	39.5-40.6	1.2	0	17.4
"No-name" Canyon Mouth	38.8	0.4	0	1.2
"No-name"	38.6-39.2	0.6	0	10.5
Goose Neck	33.2-35.6	10.1	0	57.5
Shafer Canyon Mouth	34.8	1.2	0	5.8
"No-name" Canyon Mouth	32.1	0.4	0	1.6
"No-name" Canyon Mouth	31.6	0.2	0	1.6
"No-name"	30.8-31.9	1.9	0	27.9
Little Bridge Canyon	30.0	0.8	0	1.8
Lockhart Canyon	26.5	0.6	0	5.8
"No-name"	25.3-26.5	7.7	0	31.0

Table D.2. Continued.

Site Description	River Mile	Actual Inundated Area (ha)		Historical Inundated Area (ha)
		Runoff (May 1993)	Post-Runoff (September 1993)	
<u>Colorado River</u>				
Lathrop Canyon	23.5	0.4	0	4.5
Buck Canyon	22.7	0.6	0	4.5
Gooseberry Canyon	21.7	0.4	0	2.8
Dogleg Canyon	21.2	0.4	0	3.2
Sheep Bottom	17.5-18.7	0.8	0	27.3
Indian Creek Canyon Mouth	16.5	0.6	0	4.5
Monument Creek Mouth	15.3	0.2	0	5.3
"No-name" Canyon Mouth	11.9	< 0.1	0	3.2
"No-name" Canyon Mouth	10.1	0.2	0	0.4
Salt Creek Canyon Mouth	3.4	< 0.1	0	2.9
Elephant Creek Canyon Mouth	3.0	< 0.1	0	1.7
sample statistics	n	110	109	110
	min	0	0	0.4
	max	212.0	81.0	354.0
	mean	9.1	4.2	45.0
	sum	1,000.3	459.3	4,948.4
<u>Gunnison River</u>				
"No-name"	74.3-74.7	0	0	24.0
Lawhead Gulch Bottom	70.1-71.6	0.4	0	13.8
Ferganchick's	69.5-70.5	0	0	16.7
"No-name"	68.3-69.4	0	0	12.1
"No-name"	67.5-68.2	0	0	3.8
Austin County Bridge	65.7-66.4	0	0	17.2
Austin Hwy 92 Bridge	65.0-65.5	4.1	0	13.5
"No-name"	63.3-65.3	0.8	0	52.2
"No-name"	63.0-64.6	0	0	17.7
Colorado Hwy 65 Bridge	62.6-63.2	9.9	0	14.4
Tongue Creek	61.5-62.5	16.2	0	35.6
Hutchin's	60.1-60.8	4.8	1.6	17.3
North Delta	57.6-59.5	14.0	0	49.3
South Delta	57.6-58.8	10.0	9.9	39.9
Confluence Park	56.7-57.7	8.4	6.7	40.1
Uncompahgre R. Confluence	56.3-57.0	8.1	9.3	34.4
Delta City Sewage Plant	55.6-56.5	14.7	3.6	38.0
"No-name"	54.6-55.5	1.9	1.5	27.9
"No-name"	54.2-55.1	9.8	2.2	28.2
"No-name"	53.1-54.1	1.3	0	21.7
Johnson Boy's Slough	53.2-54.2	9.9	4.1	63.1
Escalante SWA North	50.8-52.9	28.7	6.0	110.5
Escalante SWA South	50.2-52.4	48.6	4.5	82.6
"Blue Duck" Bottom	49.5-50.4	6.2	0	21.7
"No-name"	49.4-49.7	3.5	0	11.2
"No-name"	48.6-49.3	3.5	0	14.1
"No-name"	46.6-47.5	4.9	0	14.2
Escalante Ranches	41.7-42.8	2.5	0	15.1
"No-name"	39.9-41.0	16.4	0	33.8
Dominguez	36.4-36.9	2.8	1.3	11.0

Table D.2. Continued.

Site Description	River Mile	Actual Inundated Area (ha)		Historical Inundated Area (ha)
		Runoff (May 1993)	Post-Runoff (September 1993)	
<u>Gunnison River</u>				
Broughton	35.6-36.0	9.4	0	25.0
McKendricks's	34.8-36.3	3.5	0	44.9
Peeple's South (VIP Camp)	34.0-35.2	2.9	0	23.1
Peeple's Island	34.6-34.9	5.7	2.2	8.6
Sand Flat	27.3-28.7	0.4	0	14.5
Tunnel Point	26.3-26.9	2.8	0	6.6
Dads Flat	24.7-26.3	0.2	0	38.9
Deer Run	23.0-23.6	1.5	0.2	9.9
"No-name"	22.1-22.7	0	0	10.0
"No-name"	21.3-21.8	5.5	3.4	9.3
"No-name"	18.4-19.2	3.5	0	4.4
Kannah Creek	18.0-18.6	1.2	0.4	13.0
Whitewater Blding Materials	13.3-16.0	10.3	0	70.0
Duck Blind	12.7-13.8	19.6	0	32.0
Bangs Canyon	11.7-13.0	13.6	7.3	38.5
Schroeder's	4.5-6.1	10.3	0.8	26.1
Mule Farm	3.1-3.9	12.2	0	31.0
Redlands	2.2-3.0	1.3	0	4.8
sample statistics	n	48	48	48
	min	0	0	3.8
	max	48.6	9.9	110.5
	mean	7.0	1.4	27.2
	sum	335.3	65.0	1,305.7

APPENDIX E
Parameters used to categorize and rank
floodplain restoration sites

Table E.1. The 10 parameters from 35-mm colored slide photo-imagery used to categorize and rank bottomlands important to endangered fishes in the Green River, Colorado and Utah (Part 1).

Basin: Upper Colorado River
Sub-basin: Green River
River: Green River

Runoff (May 1993)										Post-Runoff (September 1993)											
Total					Hydrological					Total					Hydrological						
Potential	Inundated	Area	Acres	(ha)	Inundated	(1/N)	Acres	(ha)	Up/Down	Connection	Potential	Inundated	Area	Acres	(ha)	Inundated	(1/N)	Acres	(ha)	Up/Down	Connection
							(Y/N)		No.									(Y/N)		No.	
388.0-392.5	T11N	R24E	13	215.8	Y	215.8	Y	2	Both	Gravity ^a	215.8	Y	215.8	Y	2	Both	Gravity	183.2	Y	2	Both
381.5-382.5	T10N	R25E	2	39.3	Y	70.0	Y	2	Both	Gravity	70.0	Y	70.0	Y	2	Both	Gravity	39.3	Y	2	Both
379.5-380.5	T10N	R104W	13	90.8	Y	90.8	Y	1	Up	Gravity	90.8	Y	90.8	Y	1	Up	Gravity	0.0	N	1	Up
376.5-378.0	T10N	R103W	9	350.0	Y	213.0	N	0	None	Ground Water ^a	350.0	Y	350.0	Y	0	None	Ground Water	59.7	Y	0	None
375.5-377.0	T10N	R103W	16	478.4	Y	116.5	Y	1	None	Ground Water	478.4	Y	478.4	Y	1	None	Ground Water	203.2	Y	1	None
374.5-376.0	T10N	R103W	36	511.5	Y	116.5	Y	1	None	Ground Water	511.5	Y	511.5	Y	1	None	Ground Water	137.2	Y	1	None
372.0-374.0	T10N	R103W	56	323.4	Y	223.4	Y	1	Up	Gravity	323.4	Y	323.4	Y	1	Up	Gravity	34.2	Y	1	Up
367.5-369.5	T10N	R103W	6	50.0	Y	40.0	Y	1	Up	Gravity	50.0	Y	50.0	Y	1	Up	Gravity	51.9	Y	1	Up
362.5-369.5	T5S	R23E	4	750.1	Y	148.5	N	0	None	Ground Water	750.1	Y	750.1	Y	0	None	Ground Water	136.4	Y	0	None
362.0-363.0	T5S	R23E	16	81.9	Y	81.9	N	0	None	Ground Water	81.9	Y	81.9	Y	0	None	Ground Water	9.9	Y	0	None
361.0-362.0	T5S	R23E	21	10.2	Y	10.2	N	0	None	Ground Water	10.2	Y	10.2	Y	0	None	Ground Water	160.0	Y	0	None
299.0-300.0	T5S	R23E	29	663.7	Y	569.0	N	0	None	Ground Water	663.7	Y	663.7	Y	0	None	Ground Water	563.7	Y	0	None
296.5-297.5	T5S	R23E	31	163.9	Y	131.5	N	0	None	Ground Water	163.9	Y	163.9	Y	0	None	Ground Water	143.3	Y	0	None
296.5-296.0	T6S	R23E	11	200.6	Y	19.0	N	0	None	Ground Water	200.6	Y	200.6	Y	0	None	Ground Water	9.4	Y	0	None
294.0-295.0	T6S	R23E	14	22.2	Y	22.2	N	0	None	Ground Water	22.2	Y	22.2	Y	0	None	Ground Water	13.1	Y	0	None
292.0-294.0	T6S	R23E	15	60.5	Y	18.0	N	0	None	Ground Water	60.5	Y	60.5	Y	0	None	Ground Water	1.9	Y	0	None
292.0-293.0	T6S	R23E	16	4.7	Y	4.7	N	0	None	Ground Water	4.7	Y	4.7	Y	0	None	Ground Water	1.2	Y	0	None
288.5-289.0	T6S	R23E	7	52.5	Y	12.3	N	0	None	Ground Water	52.5	Y	52.5	Y	0	None	Ground Water	0.0	N	0	None
287.5-286.5	T6S	R23E	14	94.0	Y	88.0	N	0	None	Ground Water	94.0	Y	94.0	Y	0	None	Ground Water	0.0	N	0	None
287.5-286.0	T6S	R23E	14	94.0	Y	88.0	N	0	None	Ground Water	94.0	Y	94.0	Y	0	None	Ground Water	0.0	N	0	None
274.0-277.0	T6S	R21E	22	10.8	Y	8.3	N	0	None	Ground Water	10.8	Y	10.8	Y	0	None	Ground Water	0.0	N	0	None
271.0-274.0	T6S	R21E	31	549.3	Y	315.9	N	0	None	Ground Water	549.3	Y	549.3	Y	0	None	Ground Water	4.0	Y	0	None
269.0-272.0	T7S	R21E	33	68.7	Y	42.9	N	0	None	Ground Water	68.7	Y	68.7	Y	0	None	Ground Water	0.0	N	0	None
269.0-270.0	T7S	R21E	17	36.9	Y	36.9	N	0	None	Ground Water	36.9	Y	36.9	Y	0	None	Ground Water	32.4	Y	0	None
267.0-269.0	T7S	R21E	17	36.9	Y	36.9	N	0	None	Ground Water	36.9	Y	36.9	Y	0	None	Ground Water	0.0	N	0	None
266.0-267.0	T7S	R21E	19	32.0	Y	8.8	N	0	None	Ground Water	32.0	Y	32.0	Y	0	None	Ground Water	0.0	N	0	None
262.0-266.0	T7S	R21E	19	374.6	Y	29.9	N	0	None	Ground Water	374.6	Y	374.6	Y	0	None	Ground Water	0.0	N	0	None
257.0-265.0	T7S	R21E	32	135.1	Y	66.5	N	0	Both	Gravity	135.1	Y	135.1	Y	0	Both	Gravity	205.0	Y	0	Both
257.0-265.0	T7S	R21E	32	135.1	Y	66.5	N	0	Both	Gravity	135.1	Y	135.1	Y	0	Both	Gravity	99.6	Y	2	Both
253.0-257.0	T8S	R20E	13	846.0	Y	533.1	Y	1	Up	Gravity	888.9	Y	888.9	Y	1	Up	Gravity	778.6	Y	1	Up
254.0-256.0	T8S	R20E	32	720.0	Y	245.6	Y	1	Up	Gravity	720.0	Y	720.0	Y	1	Up	Gravity	0.0	N	1	Up
249.0-252.0	T8S	R20E	27	302.2	Y	198.4	Y	2	Both	Gravity	302.2	Y	302.2	Y	2	Both	Gravity	13.8	Y	2	Both
249.0-250.0	T8S	R20E	27	277.3	Y	164.7	Y	2	Both	Gravity	277.3	Y	277.3	Y	2	Both	Gravity	12.7	Y	2	Both
248.0-251.0	T8S	R20E	59	731.2	Y	431.7	N	0	None	Ground Water	731.2	Y	731.2	Y	0	None	Ground Water	0.0	N	0	None
248.0-243.0	T8S	R3E	5	250.4	Y	134.3	N	0	None	Ground Water ^a	250.4	Y	250.4	Y	0	None	Ground Water	3.0	Y	0	None

Browns Park National Wildlife Refuge. Water enters bottomland through a control structure and then is gravity fed through the system. Water enters bottomland either from flooding over the bank and/or seepage from ground water. The water in some agricultural areas comes from eprinkler run-off. Cousay National Wildlife Refuge.

~ 5,050 acres inundated in May '93

Table E.2. The 10 parameters from 35-mm colored slide photo-imagery used to categorize and rank bottomlands important to endangered fishes in the Green River, Colorado and Utah (Part 2).

Basin: Upper Colorado River

Sub-basin: Green River

River: Green River

Site	Levy/Berm		Levy or Berm Dividing Mainstem and Bottomland Y/N	Width/Distance ft (m)	Vegetation		Proximity to Endangered Fishes ^a RM	Species	Life Stage	Bottomland Description	Land Ownership	Photo Imagery	
	Y	N			Type (s)	(Y/N)						Type	Catalog No. ^b
Gears Cr.-Allen Ranch BPNVR ^c	Y		Y	Man-Made	Emergent ^d	Y	81.5'	Razorback	Larval	Flood-plain Depression	Private	35mm	6132-16, 4213-17
Cr.-Allen Ranch BPNVR	Y		Y	Man-Made	Emergent	Y	71.5'	Razorback	Larval	Flood-plain Depression	State	35mm	6122, 42126-10
Butch Cassidy Lake, BPNVR	Y		Y	Natural	Emergent	Y	69.5'	Razorback	Larval	Flood-plain Depression	Federal	35mm	6124, 42133-37, 4312-4
Hog Lake, BPNVR	N		Y	None	Emergent	Y	67.0'	Razorback	Larval	Flood-plain Depression	Federal	35mm	6127-28, 4316-7
Warren Bottom, BPNVR	Y		Y	Man-Made	Emergent	Y	66.0'	Razorback	Larval	Flood-plain Depression	Federal	35mm	6129-30, 4318-10
Spitzie Bottom, BPNVR	Y		Y	Natural	Emergent	Y	65.0'	Razorback	Larval	Flood-plain Depression	Federal	35mm	6131, 43111-12
J S Hoy Bottom, BPNVR	Y		Y	Man-Made	Emergent	Y	64.0'	Razorback	Larval	Flood-plain Depression	Federal	35mm	6134-36, 43113-22
Grimes Bottom, BPNVR	Y		Y	Man-Made	Emergent	Y	58.5'	Razorback	Larval	Flood-plain Depression	Federal	35mm	611-2, 43125-27
Westlake Ranch	Y		Y	Man-Made	Emergent	Y	8.5'	Razorback	Larval	Flood-plain Depression	Private	35mm	911-2, 43125-27
Westlake Ranch	Y		Y	Natural	Emergent	Y	8.0'	Razorback	Larval	Flood-plain Depression	Private	35mm	9111, 47111-15
Gravel Ponds at Jensen	N		N	-	-	N	9.0'	Razorback	Larval	Gravel Pond	Private	35mm	9112-13, 47118-19
Stewart Lake	Y		Y	Man-Made	Emergent	Y	11.0'	Razorback	Larval	Flood-plain Depression	State	35mm	9114-15, 47122-31
Ashley Creek confluence	Y		Y	Natural	Emergent	Y	12.5'	Razorback	Larval	Stream Confluence	State	35mm	9116, 47130-31
Little Stewart Lake (McCarrel)	Y		Y	Man-Made	Emergent	Y	13.5'	Razorback	Larval	Flood-plain Depression	Private/Federal	35mm	9117-18, 47132-35
Spring Hollow	N		Y	-	Emergent	Y	15.0'	Razorback	Larval	Flood-plain Terrace	Private	35mm	9119, 47136, 4811
Walker Hollow	N		Y	-	Emergent	Y	16.0'	Razorback	Larval	Flood-plain Terrace	Private	35mm	9120, 48125-8
Walker Hollow	N		Y	-	Emergent	Y	16.0'	Razorback	Larval	Flood-plain Terrace	Private	35mm	9121, 48125-8
Gravel Pit/Site	N		N	-	Emergent	N	18.0'	Razorback	Larval	Gravel Pond	Private	35mm	9123-24, 4819-11
Bonanza Bridge area	N		N	-	Emergent	Y	22.0'	Razorback	Larval	Flood-plain Depression	Federal	35mm	9126, 48111-16
Collier Draw	N		Y	-	Emergent	Y	24.5'	Razorback	Larval	Flood-plain Depression	Federal	35mm	9129, 48118-19
Horseshoe Bend area	N		Y	-	Emergent	Y	27.0'	Razorback	Larval	Flood-plain Terrace	Federal/Private	35mm	9130-37, 48121-32
The Stirrup	N		Y	-	Emergent	Y	34.0'	Razorback	Larval	Flood-plain Terrace	Federal	35mm	1014, 48133-36
Hemlock Bottom/Besser Bend	N		Y	-	Emergent	Y	37.0'	Razorback	Larval	Flood-plain Depression	Private	35mm	1017-8, 4911-5
Downstream of Besser Bend	N		Y	-	Emergent	Y	39.0'	Razorback	Larval	Flood-plain Depression	Private	35mm	1018, 4911-14
Upstream of Besser Bend	N		Y	-	Emergent	Y	40.0'	Razorback	Larval	Flood-plain Depression	Federal	35mm	1012, 49116-10
Upstream of Brennan Bottom	N		Y	-	Emergent	Y	44.0'	Razorback	Larval	Flood-plain Depression	Federal	35mm	1015, 49111-14
Brennan Bottom	N		Y	Natural	Emergent	Y	45.0'	Razorback	Larval	Flood-plain Depression	Private/Pederal	35mm	1016-17, 49115-19
Johnson Bottom, ONWR	Y		Y	Man-Made	Emergent	Y	46.0'	Razorback	Larval	Flood-plain Depression	Federal	35mm	1018-19, 49120-21
Lots Ponds, ONWR	Y		Y	Man-Made	Emergent	Y	49.0'	Razorback	Larval	Flood-plain Depression	Federal	35mm	1020-30, 49122-30
Wassket Lake, ONWR	Y		Y	Natural	Emergent	Y	54.0'	Razorback	Larval	Flood-plain Depression	Federal	35mm	1028-30, 49131-37
Sheppard Bottom, ONWR	Y		Y	Man-Made	Emergent	Y	55.0'	Razorback	Larval	Flood-plain Depression	Federal	35mm	1028-32, 49131-37
Old Charlie Wash (Dillon), ONWR	Y		Y	Man-Made	Emergent	Y	29.0'	Razorback	Larval	Flood-plain Depression	Federal/Tribe	35mm	1031-33, 5014-6
Old Charlie Wash (Dillon), ONWR	Y		Y	Man-Made	Emergent	Y	40.0'	Razorback	Larval	Flood-plain Terrace	Federal/Tribe	35mm	1034-36, 5017-9
Oursay Ute Pasture area	N		N	-	Emergent	Y	60.0'	Razorback	Larval	Flood-plain Terrace	Tribe	35mm	1036, 1111-2, 50111-12
Ouchsena River confluence area	Y		Y	Man-Made	Emergent	Y	62.0'	Razorback	Larval	Flood-plain Terrace	Tribe	35mm	

^aThe only known spawning site is located at river mile 311 at Razorback bar.
^bCatalog No. = photograph roll number followed by photograph number (e.g. 611-3 = roll 6, pictures 1 through 3).
^cWorona Park National Wildlife Refuge consists of Cattail, Inland Salt Grass, and White Top.
^dLocated upstream of Razorback bar spawning area.
^eA portion of the bottomland is privately and Federally owned.

Table E.1. Continued.

Basin: Upper Colorado River
 Sub-basin: Green River
 River: Green River

Page: 3

Runoff (May 1993)										Post-Runoff (September 1993)										
Site	Location	RM	T	R	S	Total			Hydrological Connection to River			Other Water Sources	Total			Hydrological Connection to River			Other Water Sources	
						Potential Inundated Area (ha)	Inundated (Y/N)	Area (ha)	Up/Down Stream	No.	(Y/N)		Potential Inundated Area (ha)	Inundated (Y/N)	Area (ha)	Up/Down Stream	No.	(Y/N)		
White River confluence area	248.0-249.0	T95	R208	4		614.7	Y	496.4	N	0	None	Ground Water	N	614.7	N	0.0	N	0	None	Ground Water
West Branch area	243.0-247.0	T95	R38	7		880.9	Y	615.1	N	0	None	Ground Water	Y	880.9	Y	37.4	N	0	None	Ground Water
Tia Juana Bottom	242.0-244.0	T95	R38	14		625.1	Y	319.3	N	0	None	Ground Water	Y	625.1	Y	64.6	N	0	None	Ground Water
Mouth of Willow Creek	239.0-241.0	T95	R198	14		156.3	Y	105.8	N	0	None	Ground Water	N	156.3	N	0.0	N	0	None	Ground Water
Pariette Draw area	238.0-241.0	T95	R198	14		143.1	Y	105.8	N	0	None	Ground Water	N	143.1	N	0.0	N	0	None	Ground Water
Desert Spring/Moon Bottom	227.0-228.0	T105	R198	19		11.7	Y	10.0	N	0	None	Ground Water	N	11.7	N	0.0	N	0	None	Ground Water
Fourmile Spring/King Bottom	226.0-227.0	T105	R198	19		11.7	Y	10.0	N	0	None	Ground Water	N	11.7	N	0.0	N	0	None	Ground Water
Indian Pasture/Nydas Bottom	224.0-225.0	T105	R198	26		65.0	Y	30.8	N	0	None	Ground Water	N	65.0	N	0.0	N	0	None	Ground Water
Between Long and Boat Bottom	219.0-220.0	T105	R188	27		45.4	Y	25.0	N	0	None	Ground Water	N	45.4	N	0.0	N	0	None	Ground Water
Nine Mills Creek	213.0-214.0	T115	R188	12		11.4	Y	5.0	N	0	None	Ground Water	N	11.4	N	0.0	N	0	None	Ground Water
Nutter Hole	211.0-213.0	T115	R188	35		56.1	Y	6.6	N	0	None	Ground Water	N	56.1	N	0.0	N	0	None	Ground Water
Duchess Hole	209.0-211.0	T125	R188	3		90.0	N	0.0	N	0	None	Ground Water	N	90.0	N	0.0	N	0	None	Ground Water
Tabasago Canyon/Little Horse	206.0-208.0	T125	R188	9		22.9	Y	0.0	N	0	None	Ground Water	N	22.9	N	0.0	N	0	None	Ground Water
Little Horse Bottom	204.0-207.0	T125	R188	17		85.2	Y	0.0	N	0	None	Ground Water	N	85.2	N	0.0	N	0	None	Ground Water
Downstream Maverick Canyon	203.0-204.0	T125	R188	17		147.6	Y	0.0	N	0	None	Ground Water	N	147.6	N	0.0	N	0	None	Ground Water
Gold Hole	201.0-202.0	T125	R188	30		46.7	Y	29.1	N	0	None	Ground Water	N	46.7	N	0.0	N	0	None	Ground Water
Rock House Bottom	200.0-201.0	T125	R188	30		80.7	Y	2.2	N	0	None	Ground Water	N	80.7	N	0.0	N	0	None	Ground Water
Little Rock Canyon	199.0-200.0	T125	R188	31		61.6	Y	8.2	N	0	None	Ground Water	N	61.6	N	0.0	N	0	None	Ground Water
Stamper Flat	198.0-199.0	T125	R188	31		184.6	Y	0.0	N	0	None	Ground Water	N	184.6	N	0.0	N	0	None	Ground Water
Woodoo Ponds	197.0-198.0	T135	R188	6		184.7	Y	0.0	N	0	None	Ground Water	N	184.7	N	0.0	N	0	None	Ground Water
Between Snap-Three Canyons	170.0-171.0	T155	R178	21		20.6	Y	0.0	N	0	None	Ground Water	N	20.6	N	0.0	N	0	None	Ground Water
Chandler Fall Canyon Rapids	165.5-167.0	T165	R178	31		33.4	Y	0.0	N	0	None	Ground Water	N	33.4	N	0.0	N	0	None	Ground Water
Trail and Bull Canyon Rapids	164.5-165.5	T165	R178	9		23.6	Y	0.0	N	0	None	Ground Water	N	23.6	N	0.0	N	0	None	Ground Water
Joa Nutch Canyon/Rapids	160.0-161.0	T165	R178	33		16.8	Y	0.0	N	0	None	Ground Water	N	16.8	N	0.0	N	0	None	Ground Water
Florence Cr.-Three Forks	157.0-159.0	T175	R178	9		11.0	Y	0.0	N	0	None	Ground Water	N	11.0	N	0.0	N	0	None	Ground Water
Downstream of Three Forks	153.5-155.0	T175	R178	20		16.3	Y	0.0	N	0	None	Ground Water	N	16.3	N	0.0	N	0	None	Ground Water
Last Chance Canyon/Rapids	152.0-153.0	T175	R178	23		51.2	Y	0.0	N	0	None	Ground Water	N	51.2	N	0.0	N	0	None	Ground Water
Range Creek Rapids	150.5-152.0	T175	R178	32		10.7	Y	0.0	N	0	None	Ground Water	N	10.7	N	0.0	N	0	None	Ground Water
Little Big Horn Mesa	143.0-144.0	T185	R178	29		14.0	Y	0.0	N	0	None	Ground Water	N	14.0	N	0.0	N	0	None	Ground Water
Price River confluence	137.0-138.5	T195	R178	29		23.7	Y	0.0	N	0	None	Ground Water	N	23.7	N	0.0	N	0	None	Ground Water
Downstream of Price River	136.5-137.5	T195	R168	15		14.1	Y	0.0	N	0	None	Ground Water	N	14.1	N	0.0	N	0	None	Ground Water
Short Canyon/Rapids	131.5-132.5	T205	R168	3		4.3	Y	0.0	N	0	None	Ground Water	N	4.3	N	0.0	N	0	None	Ground Water
Willow Bend-Tusher Rapid	128.5-129.5	T205	R168	19		152.2	Y	0.0	N	0	None	Ground Water	N	152.2	N	0.0	N	0	None	Ground Water
Upriar of Green River Utah	121.5-125.5	T215	R168	10		513.9	Y	0.0	N	0	None	Ground Water	N	513.9	N	0.0	N	0	None	Ground Water
Little Grand Wash	114.5-115.5	T225	R168	3		43.0	Y	0.0	N	0	None	Ground Water	N	43.0	N	0.0	N	0	None	Ground Water
Downstream of Grand Wash	113.0-114.0	T225	R168	3		62.2	Y	0.0	N	0	None	Ground Water	N	62.2	N	0.0	N	0	None	Ground Water

*Grooms Park National Wildlife Refuge.

*Water enters bottomland through a water control structure and then is gravity fed through the system.

*Water enters bottomland either from flooding over the bank and/or seepage from ground water. The water in some agricultural areas comes from sprinkler run-off.

*Gour National Wildlife Refuge.

Table E.2. Continued.

Basin: Upper Colorado River
 Sub-basin: Green River
 River: Green River

Page: 4

Site	Levy/Berm Mainstem and Bottomland Y/N	Levy/Berm Natural/Man-Made	Width/ Distance ft (m)	Vegetation Predominant Type (s)	Proximity to Endangered Fishes ¹		Bottomland Description	Land Ownership	Photo Imagery Type Catalog No. ²	
					RM	Species				
White River confluence area	Y	Man-Made	W-10	Emergent	62.0	Razorback	Flood-Plain Terrace	Tribe	35mm	10-35, 50-13-14
West Branch area	N	-	-	Emergent	66.0	Razorback	Flood-Plain Terrace	Tribe	35mm	11-3-5, 50-13-17
South Fork Willow Creek	N	-	-	Emergent	67.0	Razorback	Flood-Plain Terrace	Tribe	35mm	11-3-5, 50-13-22
Mouth of Willow Creek	Y	-	-	Emergent	70.0	Razorback	Flood-Plain Terrace	Federal	35mm	11-6-9, 50-13-28
Pariette Draw area	N	-	-	Emergent	70.0	Razorback	Flood-Plain Terrace	Private	35mm	11-12, 50-13-28
Desert Spring/Moon Bottom	N	-	-	Emergent	83.0	Razorback	Flood-Plain Terrace	Federal	35mm	11-12, 50-13-30
Fourmile Wash/King Bottom	N	-	-	-	84.0	Razorback	Flood-Plain Terrace	Federal	35mm	11-25-26, 51-11-2
Indian Pasture/Hydes Bottom	N	-	-	-	86.0	Razorback	Flood-Plain Terrace	Federal	35mm	11-25, 51-3-4
Raye and Long Bottom	N	-	-	-	89.0	Razorback	Flood-Plain Terrace	Federal	35mm	11-25, 51-3-4
Between Long and Boat Bottom	N	-	-	-	90.0	Razorback	Flood-Plain Terrace	Federal	35mm	11-25, 51-3-4
Between Long and Boat Bottom	N	-	-	-	97.0	Razorback	Flood-Plain Terrace	Federal	35mm	11-25, 51-3-4
Between Long and Boat Bottom	N	-	-	-	97.0	Razorback	Flood-Plain Terrace	Federal	35mm	11-25, 51-3-4
Nutter Hole	N	-	-	-	98.0	Razorback	Flood-Plain Terrace	Federal	35mm	12-4, 51-13-14
Duches Hole	N	-	-	-	100.0	Razorback	Flood-Plain Terrace	Federal	35mm	12-6, 51-13-18
Tabago Canyon/Little Horse	N	-	-	-	103.0	Razorback	Flood-Plain Terrace	Federal	35mm	12-6, 51-13-18
Little Horse Bottom	N	-	-	-	104.0	Razorback	Flood-Plain Terrace	Federal	35mm	12-6, 51-13-18
Downstream of Maverick Canyon	N	-	-	-	107.0	Razorback	Flood-Plain Terrace	Tribe	35mm	12-10, 52-8-9
Gold Hole	N	-	-	-	109.0	Razorback	Flood-Plain Terrace	Federal/Tribe	35mm	12-11-12, 52-11-10
Little Horse Bottom	N	-	-	-	109.0	Razorback	Flood-Plain Terrace	Tribe	35mm	12-11-12, 52-11-10
Little Horse Bottom	N	-	-	-	110.0	Razorback	Flood-Plain Terrace	Tribe	35mm	12-11, 52-11-17
Stampede Flat	N	-	-	-	111.0	Razorback	Flood-Plain Terrace	Federal	35mm	12-11, 52-11-17
Hoodoo Forms	N	-	-	-	112.0	Razorback	Flood-Plain Terrace	Federal	35mm	12-11, 52-11-17
Between Snap-Three Canyons	N	-	-	-	113.0	Razorback	Flood-Plain Terrace	Tribe	35mm	12-11, 52-11-19
Chandler Falls Canyon	N	-	-	-	140.0	Razorback	Flood-Plain Terrace	Federal	35mm	12-13-20, 52-11-19
Trail and Bull Canyon Rapids	N	-	-	-	144.0	Razorback	Flood-Plain Terrace	Federal	35mm	13-1, 53-25
Joe Hutch Canyon and Rapids	N	-	-	-	145.5	Razorback	Flood-Plain Terrace	Federal	35mm	13-1, 53-25
Joe Hutch Canyon and Rapids	N	-	-	-	145.5	Razorback	Flood-Plain Terrace	Federal	35mm	13-1, 53-25
Joe Hutch Canyon and Rapids	N	-	-	-	153.0	Razorback	Flood-Plain Terrace	Federal	35mm	13-1, 53-25
Downstream of Three Forks	N	-	-	-	153.0	Razorback	Flood-Plain Terrace	Federal	35mm	13-1, 53-25
Downstream of Three Forks	N	-	-	-	156.0	Razorback	Flood-Plain Terrace	Federal	35mm	13-13, 54-1-4
East Chance Canyon and Rapids	N	-	-	-	158.0	Razorback	Flood-Plain Terrace	Federal	35mm	13-13, 54-1-4
Range Creek Rapids	N	-	-	-	159.0	Razorback	Flood-Plain Terrace	Federal	35mm	13-13, 54-1-4
Little Big Horn Mesa	N	-	-	-	167.0	Razorback	Flood-Plain Terrace	Federal	35mm	13-13, 54-1-4
Frica River confluence	N	-	-	-	172.5	Razorback	Flood-Plain Terrace	Federal	35mm	13-13, 54-1-4
Downstream of Frica River	N	-	-	-	172.5	Razorback	Flood-Plain Terrace	Federal	35mm	13-25, 54-24-25
Short Canyon and Rapids	N	-	-	-	173.5	Razorback	Flood-Plain Terrace	Federal	35mm	13-25, 54-24-25
Upstream of Short Canyon	N	-	-	-	178.5	Razorback	Flood-Plain Terrace	Federal	35mm	13-25, 54-24-25
Upstream of Short Canyon	N	-	-	-	180.5	Razorback	Flood-Plain Terrace	Federal	35mm	13-25, 54-24-25
Upstream of Green River, Utah	N	-	-	-	185.5	Razorback	Flood-Plain Terrace	Private	35mm	14-15, 55-12-14
Little Grand Wash	N	-	-	-	185.5	Razorback	Flood-Plain Terrace	Private	35mm	14-10-13, 55-12-14
Downstream of Gland Wash	N	-	-	-	197.0	Razorback	Flood-Plain Terrace	Federal	35mm	14-20, 56-11-13
Downstream of Gland Wash	N	-	-	-	197.0	Razorback	Flood-Plain Terrace	Federal	35mm	14-22, 56-13-14

¹The only known spawning site is located at river mile 311 at Razorback bar.
²Catalog No. = photograph roll number followed by photograph number (e.g. 6:1-3 = roll 6, pictures 1 through 3).
³Crowns Farm National Wildlife Refuge consists of Cattail, Inland Salt Grass, and White Top.
⁴Located upstream of Razorback bar spawning area.
⁵A portion of the bottomland is privately and Federally owned.

Table E.1. Continued.

Basin: Upper Colorado River
 Sub-basin: Green River
 River: Green River

Page: 5

Site	Location	RM	T	R	S	Runoff (May 1993)					Post-Runoff (September 1993)				
						Total Inundated Area (ha)	Inundated Area (ha)	Y/N	Up/Down Stream	Hydrological Connection to River	Total Inundated Area (ha)	Inundated Area (ha)	Y/N	Up/Down Stream	Hydrological Connection to River
Fivemile Wash/Little Valley	111.0-112.0	T225	R16E		8	15.0	N	N	0	None	Ground Water	15.0	N	0	None
Downstream of Ninemila Wash	109.0-110.0	T225	R16E		20	20.5	N	N	0	None	Ground Water	20.5	N	0	None
Anvil Bottom	101.5-102.5	T235	R16E		11	9.7	Y	Y	0	None	Ground Water	9.7	N	0	None
Upstream of San Rafael R.	98.0-99.0	T235	R16E		23	26.4	Y	Y	0	None	Ground Water	26.4	N	0	None
San Rafael River confluence	96.5-97.5	T235	R16E		36	44.0	Y	Y	0	None	Ground Water	44.0	N	0	None
White and Red Wash	95.0-96.5	T235	R17E		31	16.5	Y	Y	0	None	Ground Water	16.5	N	0	None
Between Red Wash-Bull Bottom	93.0-94.0	T245	R16E		16	19.7	Y	Y	0	None	Ground Water	19.7	N	0	None
Between Red Wash-Bull Bottom	92.0-93.0	T245	R16E		12	39.4	N	N	0	None	Ground Water	39.4	N	0	None
Between Red Wash-Bull Bottom	91.0-92.0	T245	R16E		13	36.4	N	N	0	None	Ground Water	36.4	N	0	None
Labyrinth Canyon confluence	88.5-91.5	T245	R16E		13	65.2	N	N	0	None	Ground Water	65.2	N	0	None
Theresa Canyon/Trin-Alcove Bend	87.0-88.0	T245	R16E		23	35.9	N	N	0	None	Ground Water	35.9	N	0	None
Junes Bottom	85.0-86.0	T245	R17E		31	36.5	N	N	0	None	Ground Water	36.5	N	0	None
Bull Hollow	84.0-85.0	T245	R17E		31	23.4	Y	Y	0	None	Ground Water	23.4	N	0	None
Bull Hollow-Tennilla Canyon	81.0-83.0	T255	R17E		6	37.8	Y	Y	0	None	Ground Water	37.8	N	0	None
Tennilla Canyon/Bottom	79.0-81.0	T255	R17E		8	48.1	Y	Y	0	None	Ground Water	48.1	N	0	None
Kemp Canyon	76.0-78.5	T255	R17E		9	48.1	Y	Y	0	None	Ground Water	48.1	N	0	None
Hay Joa Canyon	74.5-76.0	T255	R17E		16	25.0	Y	Y	0	None	Ground Water	25.0	N	0	None
Hay Joa Canyon-Spring Canyon	74.0-75.0	T255	R17E		16	33.6	N	N	0	None	Ground Water	33.6	N	0	None
Spring Canyon Point	69.5-74.0	T255	R17E		20	89.3	N	N	0	None	Ground Water	89.3	N	0	None
Bowknot Bend	62.0-70.0	T255	R17E		28	261.1	Y	Y	0	None	Ground Water	261.1	N	0	None
Twomile Canyon-Deadman Point	60.0-62.0	T255	R17E		32	36.4	Y	Y	0	None	Ground Water	36.4	N	0	None
Deadman Point-Horseshoe Can.	58.0-60.0	T265	R17E		4	58.3	N	N	0	None	Ground Water	58.3	N	0	None
Deadman Point-Horseshoe Can.	55.0-56.5	T265	R17E		3	64.3	Y	Y	0	None	Ground Water	64.3	N	0	None
Cottonwood Bottom	54.0-55.0	T265	R17E		12	43.6	N	N	0	None	Ground Water	43.6	N	0	None
Cottonwood-Mineral Bottoms	53.0-54.0	T265	R17E		24	74.3	N	N	0	None	Ground Water	74.3	N	0	None
Mineral-Tidwell Bottoms	51.0-52.5	T265	R17E		25	162.9	N	N	0	None	Ground Water	162.9	N	0	None
Tidwell Bottom	50.0-51.5	T265	R17E		26	32.1	N	N	0	None	Ground Water	32.1	N	0	None
Horsathief Bottom	49.0-50.0	T265	R17E		26	91.1	N	N	0	None	Ground Water	91.1	N	0	None
Woodruff Bottom	48.0-49.0	T275	R17E		3	12.7	N	N	0	None	Ground Water	12.7	N	0	None
Point Bottom	45.5-47.5	T275	R17E		2	69.6	N	N	0	None	Ground Water	69.6	N	0	None
Saddle Bottom	45.0-46.0	T275	R17E		2	104.7	N	N	0	None	Ground Water	104.7	N	0	None
Horsathief Canyon	44.0-45.5	T275	R17E		12	78.3	Y	Y	0	None	Ground Water	78.3	N	0	None
Uphaval Bottom/Canyon	43.5-44.5	T275	R17E		13	30.7	N	N	0	None	Ground Water	30.7	N	0	None
Hardcrabla Bottom	42.5-43.5	T275	R17E		4	156.2	Y	Y	0	None	Ground Water	156.2	N	0	None
Hardcrabla-Fort Bottoms	38.5-42.5	T275	R17E		22	76.8	Y	Y	0	None	Ground Water	76.8	N	0	None
Potato Bottom	35.5-38.0	T275	R17E		26										

*Browns Park National Wildlife Refuge.

*Water enters bottomland through a water control structure and then is gravity fed through the system.

*Water enters bottomland either from flooding over the bank and/or seepage from ground water. The water in some agricultural areas comes from sprinkler run-off.

*Quay National Wildlife Refuge.

Table E.2. Continued.

Basin: Upper Colorado River
 Sub-basin: Green River
 River: Green River

Site	Levy/Berm		Width/ Distance ft (m)	Vegetation		Proximity to Endangered Fishes*		Bottomland Description	Land Ownership	Photo Imagery Catalog No. ^b	
	Levy or Berm Y/N	Mainstem and Bottomland Natural/Man-Made		(Y/N)	Predominant Type(s)	RM	Species			Type	
Fivemile Wash/Little Valley	N	-	-	N	-	199.0	Razorback	Flood-Plain Terrace	Federal	35mm	14124, 56115
Downstream of Niemile Wash	N	-	-	N	-	208.0	Razorback	Flood-Plain Terrace	Federal	35mm	14124, 56115
Upstream of San Rafael River	N	-	-	N	-	208.5	Razorback	Flood-Plain Terrace	Federal	35mm	14134, 56129-30
San Rafael River confluence	N	-	-	N	-	212.0	Razorback	Flood-Plain Terrace	Federal	35mm	14137, 56132
White and Red Wash	N	-	-	N	-	213.5	Razorback	Flood-Plain Terrace	Federal	35mm	1516, 56133
Between Red Wash and Bull Bottom	N	-	-	N	-	214.5	Razorback	Flood-Plain Terrace	Federal	35mm	1518, 56135-36
Bull Bottom-Labyrinth Canyon	N	-	-	N	-	215.5	Razorback	Flood-Plain Terrace	Federal	35mm	1519
Labyrinth Canyon confluence	N	-	-	N	-	218.0	Razorback	Flood-Plain Terrace	Federal	35mm	15111, 5712-3
Three Canyon/Twin-Alcove Bend	N	-	-	N	-	218.5	Razorback	Flood-Plain Terrace	Federal	35mm	15112, 5714
Three Canyon/Twin-Alcove Bend	N	-	-	N	-	221.0	Razorback	Flood-Plain Terrace	Federal	35mm	15112, 5714
Three Canyon/Twin-Alcove Bend	N	-	-	N	-	223.0	Razorback	Flood-Plain Terrace	Federal	35mm	15114, 5716-7
Three Canyon/Twin-Alcove Bend	N	-	-	N	-	225.0	Razorback	Flood-Plain Terrace	Federal	35mm	15115-16, 5718-12
Bull Nollow	N	-	-	N	-	226.0	Razorback	Flood-Plain Terrace	Federal	35mm	15117, 5713-17
Bull Nollow-Tensile Canyon	N	-	-	N	-	228.0	Razorback	Flood-Plain Terrace	Federal	35mm	15118, 57114-17
Tensile Canyon and Bottom	N	-	-	N	-	229.0	Razorback	Flood-Plain Terrace	Federal	35mm	15119, 57118-20
Reg Springs Canyon	N	-	-	N	-	232.5	Razorback	Flood-Plain Terrace	Federal	35mm	15121-22, 57121-22
Ney Joe Canyon-Spring Canyon	N	-	-	N	-	235.0	Razorback	Flood-Plain Terrace	Federal	35mm	15124, 57125-26
Springs Canyon	N	-	-	N	-	235.0	Razorback	Flood-Plain Terrace	Federal	35mm	15124, 57125-26
Spring Canyon Point	N	-	-	N	-	237.0	Razorback	Flood-Plain Terrace	Federal	35mm	15125-26, 57127-30
Booknot Bend	N	-	-	N	-	241.0	Razorback	Flood-Plain Terrace	Federal	35mm	15127-34, 57131-38, 5811-6
Twomile Canyon-Deadman Point	N	-	-	N	-	249.0	Razorback	Flood-Plain Terrace	Federal	35mm	15135, 5816-7
Deadman Point-Norreshoe Canyon	N	-	-	N	-	251.0	Razorback	Flood-Plain Terrace	Federal	35mm	15136, 5818
Downstream of Norreshoe Canyon	N	-	-	N	-	252.0	Razorback	Flood-Plain Terrace	Federal	35mm	15137, 5819-13
Cottonwood Bottom	N	-	-	N	-	254.5	Razorback	Flood-Plain Terrace	Federal	35mm	1611, 5814-16
Cottonwood-Mineral Bottoms	N	-	-	N	-	259.0	Razorback	Flood-Plain Terrace	Federal	35mm	1611, 5814-16
Mineral Bottoms	N	-	-	N	-	259.0	Razorback	Flood-Plain Terrace	Federal	35mm	1613, 5819-20
Rivall Bottom	N	-	-	N	-	258.5	Razorback	Flood-Plain Terrace	Federal	35mm	1614-6, 58121
Rivall Bottom	N	-	-	N	-	259.5	Razorback	Flood-Plain Terrace	Federal	35mm	1617, 58121-22
Horseshief Bottom	N	-	-	N	-	261.0	Razorback	Flood-Plain Terrace	Federal	35mm	1618, 58123
Woodruff Bottom	N	-	-	N	-	261.5	Razorback	Flood-Plain Terrace	Federal	35mm	1619-10
Point Bottom	N	-	-	N	-	263.5	Razorback	Flood-Plain Terrace	Federal	35mm	16111-12, 58124-26
Saddle Nose Bottom	N	-	-	N	-	263.5	Razorback	Flood-Plain Terrace	Federal	35mm	16113, 58125
Horseshief Canyon	N	-	-	N	-	265.0	Razorback	Flood-Plain Terrace	Federal	35mm	16115, 58127
Horseshief-Upheaval Bottom	N	-	-	N	-	265.5	Razorback	Flood-Plain Terrace	Federal	35mm	16115, 58127
Ward Bottom	N	-	-	N	-	267.5	Razorback	Flood-Plain Terrace	Federal	35mm	16116, 58128-30
Ward Bottom	N	-	-	N	-	267.5	Razorback	Flood-Plain Terrace	Federal	35mm	16117-18, 58131-35
Ward Bottom	N	-	-	N	-	268.5	Razorback	Flood-Plain Terrace	Federal	35mm	16119-20, 58136-37, 5911-4
Poteco Bottom	N	-	-	N	-	273.0	Razorback	Flood-Plain Terrace	Federal	35mm	

*The only known spawning site is located at river mile 311 at Razorback bar.
^aCatalog No. = photograph roll number followed by photograph number (e.g. 6:1-3 = roll 6, pictures 1 through 3).
^bBrown Park National Wildlife Refuge.
^cLocated upstream of Razorback bar spawning area.
^dA portion of the bottomland is privately and Federally owned.

Table E.1. Continued.

Basin: Upper Colorado River
 Sub-basin: Green River
 River: Green River

Page: 7

Runoff (May 1993)										Post-Runoff (September 1993)									
Site	Location	RM	T	R	S	Total			Hydrological			Total			Hydrological				
						Potential Inundated Area (ha)	Inundated (Y/N)	Acres (ha)	Up/Down Stream	Connection to River	Potential Inundated Area (ha)	Inundated (Y/N)	Acres (ha)	Up/Down Stream	Connection to River				
Potato-Beaver Bottoms	34.5-35.5	T285	R17E		2	44.2	N	0.0	None	Ground Meter		44.2	N	0.0	N	0	None	Ground Meter	
Beaver-Queen Ann Bottoms	33.5-34.5	T285	R17E		11	49.1	Y	5.1	0	Ground Meter		49.1	N	0.0	N	0	None	Ground Meter	
Queen Ann Bottom	33.0-34.0	T285	R17E		11	68.8	Y	2.4	0	None	Ground Meter	68.8	Y	0.0	N	0	None	Ground Meter	
Queen Ann-Anderson Bottoms	32.0-33.0	T285	R17E		14	15.0	N	0.0	None	Ground Meter	15.0	Y	0.0	N	0	N	0	None	Ground Meter
Anderson Bottom	32.0-32.5	T285	R17E		24	23.3	Y	2.1	0	None	Ground Meter	23.3	Y	0.0	N	0	None	Ground Meter	
Anderson-Anderson Bottoms	29.0-30.0	T285	R17E		12	112.2	N	0.0	None	Ground Meter	112.2	Y	0.0	N	0	N	0	None	Ground Meter
Anderson-Anderson Bottoms	27.0-28.0	T285	R18E		17	117.8	N	0.0	None	Ground Meter	117.8	Y	0.0	N	0	N	0	None	Ground Meter
Stillwater Canyon-Sphinx	26.0-28.5	T285	R18E		17	96.3	Y	1.3	0	None	Ground Meter	96.3	Y	0.0	N	0	None	Ground Meter	
Downstream Valentia Bottom	25.5-26.5	T285	R18E		19	67.6	Y	17.7	0	None	Ground Meter	67.6	Y	0.0	N	0	None	Ground Meter	
Valentia-Tukedo Bottoms	24.0-25.5	T285	R18E		20	96.9	N	0.0	None	Ground Meter	96.9	Y	0.0	N	0	N	0	None	Ground Meter
Tukedo Bottom-Turks Head	19.5-24.0	T285	R18E		32	235.9	Y	4.0	0	None	Ground Meter	235.9	Y	0.0	N	0	None	Ground Meter	
Deedhorse Canyon confluence	18.5-20.0	T285	R18E		32	26.3	Y	1.6	0	None	Ground Meter	26.3	Y	0.0	N	0	None	Ground Meter	
Deedhorse Canyon	17.0-19.0	T295	R18E		4	57.8	Y	2.6	0	None	Ground Meter	57.8	Y	0.0	N	0	None	Ground Meter	
Deedhorse-Horse Canyons	16.0-17.0	T295	R18E		20	21.3	N	0.0	None	Ground Meter	21.3	Y	0.0	N	0	N	0	None	Ground Meter
Horse Canyon	13.5-14.0	T295	R18E		27	51.7	Y	7.4	0	None	Ground Meter	51.7	Y	0.0	N	0	None	Ground Meter	
Downstream of Horse Canyon	12.5-13.5	T295	R18E		26	60.4	Y	2.3	0	None	Ground Meter	60.4	Y	0.0	N	0	None	Ground Meter	
Horse and Jasper Canyons	10.5-12.5	T295	R18E		35	55.8	Y	1.8	0	None	Ground Meter	55.8	Y	0.0	N	0	None	Ground Meter	
Jasper Canyon	8.0-10.5	T295	R18E		2	33.5	Y	0.6	0	None	Ground Meter	33.5	Y	0.0	N	0	None	Ground Meter	
Short Canyon-Colored River	2.0-3.5	T305	R19E		6	13.8	Y	2.0	0	None	Ground Meter	13.8	Y	0.0	N	0	None	Ground Meter	
Short Canyon-Colored River	1.0-2.0	T305	R18E		23	14.8	Y	0.6	0	None	Ground Meter	14.8	Y	0.0	N	0	None	Ground Meter	
Near Colorado R. confluence	0.5-1.5	T305	R18E		25	29.0	Y	0.5	0	None	Ground Meter	29.0	Y	0.0	N	0	None	Ground Meter	
Near Colorado R. confluence	0.0-0.5	T305	R19E		7	15.6	N	0.0	None	Ground Meter	15.6	Y	0.0	N	0	N	0	None	Ground Meter

*Brown Park National Wildlife Refuge.
 Meter enters bottomland through a water control structure and then is gravity fed through the system.

*Meter enters bottomland either from flooding over the bank and/or seepage from ground water. The water in some agricultural areas comes from sprinkler run-off.

*Quey National Wildlife Refuge.

Table E.2. Continued.

Basin: Upper Colorado RiverSub-basin: Green RiverRiver: Green RiverPage: 8

Site	Levy or Berm Dividing Mainstem and Bottomland		Width/ Distance ft (m)	Vegetation		Proximity To Endangered Fishes ^a RM	Species	Life Stage	Bottomland Description	Land Ownership	Photo Inventory	
	Y/N	Natural/Man-Made		(Y/N)	Predominant Type (s)						Type	Catalog No. ^a
Pocato-Beaver Bottoms	N	-	-	N	-	275.5	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	16:21
Beaver-Queen Ann Bottoms	N	-	-	N	-	276.5	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	16:22
Queen Ann Bottom	N	-	-	N	-	277.0	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	16:23
Queen Ann-Anderson Bottoms	N	-	-	N	-	278.0	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	16:24
Anderson Bottom	N	-	-	N	-	279.5	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	16:25
Unknown and Valentine Bottoms	N	-	-	N	-	279.0	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	16:26
Stallwater Canyon and Spinx	N	-	-	N	-	280.0	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	16:27
Stallwater Canyon	N	-	-	N	-	282.5	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	16:28-29, 13-18
Downstream of Valentine Bottom	N	-	-	N	-	284.5	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	16:30-31, 59:16-20
Valentine-Tuxedo Bottoms	N	-	-	N	-	285.5	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	16:32, 59:19-22
Tuxedo Bottom	N	-	-	N	-	287.0	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	16:33, 59:21-23
Deedhorse Canyon confluence	N	-	-	N	-	291.0	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	16:34-37, 17:1-3, 59:23-29
Downstream of Deedhorse Canyon	N	-	-	N	-	292.0	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	17:4, 59:29
Deedhorse and Horse Canyons	N	-	-	N	-	293.0	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	17:5-7, 59:30-32
Horse Canyon	N	-	-	N	-	297.0	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	17:8-11, 59:36-37
Downstream of Horse Canyon	N	-	-	N	-	297.5	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	17:12-13, 59:36-37
Horse Canyon and Jasper Canyon	N	-	-	N	-	298.5	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	17:14-15, 60:1-2
Jasper Canyon	N	-	-	N	-	300.5	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	17:16-20, 60:7
Short Canyon and Colorado River	N	-	-	N	-	303.0	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	17:21-23, 60:9-10
Upstream of Colorado River	N	-	-	N	-	307.5	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	17:24-25, 60:12-13
Colorado River confluence	N	-	-	N	-	309.0	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	17:26-29, 60:14
Colorado River	N	-	-	N	-	312.5	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	17:30-31, 60:15
Colorado River confluence	N	-	-	N	-	330.5	Resorbeck	Larvel	Flood-Plain Terrace	Federal	35mm	17:32-37, 60:16

^aThe only known spawning site is located at river mile 311 at Resorbeck bar.

^aCatalog No. = photograph roll number followed by photograph number (e.g. 6:1-3 = roll 6, pictures 1 through 3).

^aJuvenile Park National Wildlife Refuge.

^aThe most common emergent vegetation consists of Cattail, Inland Salt Grass, and White Top.

^aUnpublished data from the Green River National Wildlife Refuge.

^aA portion of the bottomland is privately and Federally owned.

Table E.3. The 10 parameters from 35-mm color slide photo-imagery used to categorize and rank bottomlands important to endangered fishes in the Yampa River, Colorado (Part 1).

Basin: Upper Colorado River

Sub-basin: Green River

River: Yampa River

Page: 9

Site	Location	Runoff (May 1993)						Post-Runoff (September 1993)										
		Total Potential Inundated Area			Hydrological Connection to River			Total Potential Inundated Area			Hydrological Connection to River							
		N	Y	S	Inundated Acres (ha)	(Y/N)	Up/Down Stream	Other Water Sources	N	Y	S	Inundated Acres (ha)	(Y/N)	Up/Down Stream	Other Water Sources			
Upstream Snake R. confluence	48.0-48.5	T6N	R98W	19	21.9	Y	21.9	Y	2	Both	Ground Water ^a	21.9	N	0.0	N	0	None	Ground Water

Browns Park National Wildlife Refuge.

Water enters bottomland through a water control structure and then is gravity fed through the system.

Water enters bottomland either from flooding over the bank and/or seepage from ground water. The water in some agricultural areas comes from sprinkler run-off. Ouray National Wildlife Refuge.

Table E.4. The 10 parameters from 35-mm color slide photo-imagery used to categorize and rank bottomlands important to endangered fishes in the Yampa River, Colorado (Part 2).

Basin: Upper Colorado River

Sub-basin: Green River

River: Yampa River

Page: 9

Site	Levy/Berm Levy of Berm Dividing Point and Bottomland Y/N	Width/ Distance Natural/Man-Made ft (m)	Vegetation		Proximity to Endangered Fishes* Life Stage	Predominate Geological Description	Land Ownership	Photo Imagery Type	Catalog No. *		
			(Y/N)	Type(s)							
Upstream Snake River confluence	N	-	N	-	82.5°	Razorback	Larval	Flooded Side Channel	Private	35mm	1:6, 38:12

The only known spawning site is located at river mile 311 at Razorback bar.

Catalog No. = photograph roll number followed by photograph number (e.g. 6:1-3 = roll 6, pictures 1 through 3).

Browns Park National Wildlife Ref
The most common emergent vegetati

The most common emergent vegetation consists of cattail, inland salt grass, and white top. Located upstream of Razorback bar spawning area.

A portion of the bottomland is privately and Federally owned.

Table E.5. The 10 parameters from 35-mm colored slide photo-imagery used to categorize and rank bottomlands important to endangered fishes in the White River, Colorado and Utah (Part 1).

Basin: Upper Colorado River
 Sub-basin: Green River
 River: White River

Page: 10

Site	Location	RM	T	R	S	Runoff (May 1993)				Post-Runoff (September 1993)			
						Total Potential Inundated Area (ha)	Total Area Inundated (ha)	Hydrological Connection to River		Total Potential Inundated Area (ha)	Total Area Inundated (ha)	Hydrological Connection to River	
						(Y/N)	(Y/N)	Up/Down Stream	No.	(Y/N)	(Y/N)	Up/Down Stream	No.
Oxbow downstream Highway 64	91.0-93.0	TIN	R102N	4	18.9	Y	18.9	Y	2	Y	17.6	Y	2
White/Green River confluence	0.0-2.5	T9S	R10E	4	614.7	Y	496.4	Y	0	N	0.0	N	0
								Ground Water ^a				Ground Water	
								Ground Water				Ground Water	

^aBrowns Park National Wildlife Refuge.
^bWater enters bottomland through a water control structure and then is gravity fed through the system.
^cWater enters bottomland either from flooding over the bank and/or seepage from ground water. The water in some agricultural areas comes from sprinkler run-off.
^dCourtesy National Wildlife Refuge.

Table E.6. The 10 parameters from 35-mm colored slide photo-imagery used to categorize and rank bottomlands important to endangered fishes in the White River, Colorado and Utah (Part 2).

Basin: Upper Colorado River
 Sub-basin: Green River
 River: Yampa River

Page: 10

Site	Levy of Berm Dividing Mainstem and Bottomland Y/N	Levy/Berm Natural/Man-Made	Width/Distance ft (m)	Vegetation Predominant Type(s)		Proximity to Endangered Fishes ^a RM	Species	Bottomland Description	Land Ownership	Photo Imagery Type		Catalog No. ^b
				(Y/N)								
				(Y/N)								
Oxbow downstream of Highway 65	N	-	W-10	N	Emergent	156.0	Razorback	Flooded Side Channel	Private	35mm	5:7	70:16-17
White/Green River confluence	Y	Man-Made	-	Y	-	65.0	Razorback	Flood-plain Terrace	Tribe	35mm	3:1-4	68:11-2

^aThe only known spawning site is located at river mile 311 at Razorback bar.
^bCatalog No. = photograph roll number followed by photograph number (a.g. 6:11-3 = roll 6, pictures 1 through 3).
^cBrowns Park National Wildlife Refuge.
^dThe bottomland consists of alluvial deposits of silt, sand, and gravel.
^eLocated upstream of Razorback bar spawning area.
^fA portion of the bottomland is privately and Federally owned.

Table E.7. The 10 parameters used to rank 158 potential floodplain restoration sites important to endangered fishes of the Upper Colorado River Basin.

Basin: Upper Colorado River
 Sub-basin: Colorado
 River: Colorado River

Page: 1

Runoff (May 1993)										Post-Runoff (September 1993)									
Site	Location	RM	T	R	S	Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River			Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River				
									(Y/N)	No.	Up/Down Stream				(Y/N)	No.	Up/Down Stream		
																		Other Water Sources	
Rifle West	238.0-241.1	6S	83W	16.17		150.7	Y	12.2	N	0	None	None	150.7	Y	12.1	N	0	None	None
Rifle I-70 West Interchange	238.5-238.0	6S	83W	13.18		126.4	Y	5.3	N	0	None	None	126.4	Y	5.3	N	0	None	None
		6S	94W	24															
CDOW/Gentry Property	234.2-235.8	6S	94W	23.24		66.8	Y	10.4	N	0	None	None	66.8	Y	10.2	N	0	None	None
		6S	94W	27		5.8	Y	5.3	Y	2	Both	None	5.8	Y	1.8	N	0	None	None
"Trash-can Pond"	230.2-230.8	6S	94W	30															
		6S	94W	25		54.4	Y	6.5	Y	4	Both	Springs	54.4	Y	2.3	Y	2	Both	Springs
Mahaffey/Lemon	228.2-229.8			35.38															Irrigation
Mahaffey's	228.5-229.5	6S	94W	25.26		30.5	Y	2.5	Y	1	Up	None	30.5	Y	2.2	Y	1	Up	None
Hoaglund's	227.1-227.7	6S	94W	2.35		18.4	Y	1.2	Y	2	Both	Springs	16.4	Y	< 0.1	Y	1	Dn	Springs
																			Irrigation
Dere/Ortiz	227.5-228.5	6S	94W	35		39.3	Y	6.1	Y	1	Up	None	39.3	Y	8.0	N	0	None	None
Parachute East	223.1-225.1	6S	94W	5		35.7	Y	14.2	Y	4	Both	None	35.7	Y	2.5	Y	4	Both	None
Knight's	224.5-225.4	6S	94W	4.5		28.2	Y	4.4	Y	1	Up	Springs	28.2	N	0	N	0	None	None
Parachute Bridge	223.0-223.4	7S	95W	7		13.7	Y	2.1	Y	2	Both	None	13.7	Y	0.7	Y	1	Dn	Springs
"No-name"	222.7-223.1	7S	95W	18		8.6	Y	0.4	N	0	None	Springs	8.6	Y	0.8	Y	1	Dn	Springs
"No-name"	221.1-222.9	7S	95W	7.18		23.4	Y	11.5	N	0	None	Groundwater	23.4	Y	11.3	N	0	None	None
Battlement Mesa	220.7-221.7	7S	96W	13.24		34.4	Y	10.2	Y	4	Both	Groundwater	34.4	Y	3.9	Y	1	Dn	Springs
EXXON	218.9-220.1	7S	96W	23		34.9	Y	3.1	Y	1	Dn	None	34.9	Y	0	N	0	None	None
Doyle Property	218.4-219.5	7S	96W	23		8.7	N	0	N	0	None	None	8.7	N	0	N	0	None	None
"No-name"	217.2-218.0	7S	96W	26.27		22.3	Y	1.4	Y	2	Both	None	22.3	Y	0	N	0	None	None

Table E.7. Continued.

Basin: Upper Colorado River
 Sub-basin: Colorado River
 River: Colorado River

Page: 2

Site	Levy/Berm		Width/ Length (m)	Vegetation		Proximity to Endangered Species		Predominant Habitat Type	Land Ownership	Photo Imagery	
	Y/N	Levy or Berm Dividing Mainstem and Bottomland Natural/Man-made		Y/N	Predominant Type(s)	RM	Species			Type*	Catalog No.*
Rifle West	Y	Man-made	3,901-L	Y	Cottonwood	185.1	CS	Gravel-Pit Ponds	Private	ASCS	20: 1-4; 74: 2-3
Rifle 1-70 West Interchange	Y	Man-made	1,463-L	N	None	185.1	CS	Gravel-Pit Ponds	Private	ASCS	20: 5-6; 74: 5-6
COOW/Gentry Property	Y	Man-made	650-L	N	None	185.1	CS	Gravel-Pit Ponds	Public	ASCS	20: 8-9; 74: 11-12
"Trash-can Pond"	Y	Man-made	76-L	Y	Cottonwood	185.1	CS	Gravel-Pit Pond	Private	COE	CR-27-3 (6/24/82)
Mahaffey/Lemon	N	—	—	Y	Tamarisk	185.1	CS	Terrace	Private	ASCS	74: 15
Mahaffey/s	N	—	—	Y	Tamarisk	185.1	CS	Gravel-Pit Pond	Private	ASCS	20: 13; 74: 16
Hoaglund's	N	—	—	Y	Tamarisk	185.1	CS	Terrace	Private	ASCS	20: 15; 74: 18-19
Dere/Ortiz	Y	Man-made	244-L	Y	Tamarisk	185.1	CS	Gravel-Pit Ponds	Private	ASCS	20: 14; 74: 17-18
Parachute East	N	—	—	Y	Cottonwood	185.1	CS	Terrace	Private	ASCS	20: 18; 74: 2
Knight's	Y	Man-made	81-L	N	—	185.1	CS	Ponds	Private	COE	CR 28-1 (6-24-82)
Parachute Bridge	N	—	—	Y	Tamarisk	185.1	CS	Terrace	Private	ASCS	74: 20
"No-name"	M	—	—	Y	Cottonwood/Tamarisk	185.1	CS	Terrace	Private	ASCS	20: 20; 74: 21
"No-name"	Y	Man-made	1,707-L	N	—	185.1	CS	Gravel-Pit Ponds	Private	ASCS	20: 19; 74: 22
Battlement Mesa	N	—	—	Y	Cottonwood	185.1	CS	Terrace	Private	ASCS	20: 20; 74: 23
EXXON	N	—	—	Y	Cottonwood	185.1	CS	Terrace	Private	COE	CR-24-1 (6/24/82)
Doyle Property	N	—	—	Y	Cottonwood	185.1	CS	Terrace	Private	ASCS	74: 24
"No-name"	Y	Man-made	1,087-L	N	—	185.1	CS	Terrace	Private	ASCS	20: 21; 75: 2
						185.1	CS	Terrace	Private	ASCS	20: 22; 75: 2
						185.1	CS	Terrace	Private	ASCS	20: 25; 75: 4

* ASCS = Consolidated Farm Service Agency (U. S. Department of Agriculture); COE = U. S. Army Corps of Engineers; U. M. = University of Montana; 1:6000 Colored Aerial Photographs, 4/28/83, 9/2/83
 * Catalog No. = photograph roll number followed by photograph number (e.g. 6: 1-3 = roll 6, pictures 1 through 3).

Table E.8. The 10 parameters used to rank 158 potential floodplain restoration sites important to endangered fishes of the Upper Colorado River Basin.

Basin: Upper Colorado River

Sub-basin: Colorado

River: Colorado River

Page: 3

Runoff (May 1893)										Post-Runoff (September 1893)							
Site	Location RM	T	R	S	Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River			Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River			
								(Y/N)	No.	Up/Down Stream				Other Water Sources	(Y/N)	No.	Up/Down Stream
Una North	218.4-217.4	7S	96W	33,34	40.7	Y	8.4	N	0	None	40.7	Y	8.3	N	0	None	None
Wallace Creek Island	215.9-218.5	7S	96W	33	13.3	Y	3.4	Y	2	Both	13.3	N	0	N	0	None	None
Stoddard Property	209.4-210.1	8S	87W	26,27	23.2	Y	8.1	N	0	None	23.2	Y	8.1	N	0	None	None
Debeque F70 Slough	209.5-211.4	8S	87W	22,23	40.4	Y	8.7	Y	2	Both	40.4	Y	3.8	Y	1	Dn	None
				24													
Latham's	207.9-208.8	8S	87W	33	26.1	Y	8.2	Y	2	Both	26.1	Y	1.8	N	0	None	Irrigation
"No-name"	206.2-207.5	8S	87W	32,33	22.9	Y	2.1	Y	2	Both	22.8	N	0	N	0	None	None
Elder's	204.7-206.8	8S	87W	33	65.1	Y	22.8	Y	2	Both	65.1	Y	22.8	Y	1	Dn	Irrigation
"No-name"	202.3-204.2	9S	87W	8,17	86.4	Y	6.9	Y	2	Both	86.4	Y	3.3	N	0	None	None
"No-name"	201.9-202.4	9S	87W	19,20	8.1	Y	0.2	Y	2	Both	8.1	N	0	N	0	None	None
Long Point Bottom	198.0-198.9	9S	87W	31,36	10.8	Y	2.3	Y	4	Both	2.3	Y	0.2	Y	1	Dn	None
Beavertail	194.3-195.8	10S	98W	12,13	12.8	Y	2.0	Y	0	None	12.8	Y	1.8	N	0	None	None
Island Acres	191.1-192.1	10S	98W	23,26	28.1	Y	11.4	N	0	None	28.1	Y	11.4	N	0	None	River Seepage
Cameo	190.0-191.0	10S	98W	26,27	24.8	Y	1.2	Y	3	Both	24.8	Y	1.2	Y	1	Dn	Irrigation
Palisade	184.2-185.0	1S	2E	3	22.0	Y	2.5	N	0	None	22.0	Y	2.1	N	0	None	None
"No-name"	183.3-184.3	1S	2E	3,4	17.2	Y	3.8	Y	1	Dn	17.2	Y	1.3	N	0	None	Irrigation
Labor Camp	182.9-183.8	1S	2E	4	19.8	Y	8.3	Y	4	Both	19.8	Y	2.7	Y	1	Dn	Irrigation
"No-name"	181.6-182.8	1S	2E	5,8	43.3	Y	0.6	N	0	None	43.3	N	0	N	0	None	None
"No-name"	181.7-182.2	1S	2E	8	8.9	Y	2.8	Y	2	Both	8.9	Y	2.8	Y	2	Both	Irrigation
"No-name"	180.4-181.5	1S	2E	7,8	21.2	Y	4.2	N	0	None	21.1	Y	1.4	N	0	None	Irrigation
"No-name"	180.4-181.5	1S	2E	7	122.5	Y	20.7	Y	1	Dn	122.5	Y	17.8	Y	1	Dn	Irrigation
Clifton Water Treatment	179.1-181.1	1S	1E	13													
Clifton Pond	177.7-178.2	1S	1E	14,23	81.7	Y	21.7	N	0	None	81.7	Y	15.8	Y	2	Both	Irrigation
Com Lake	176.6-177.7	1S	1E	22,23	38.6	Y	14.5	Y	2	Both	38.8	Y	8.3	Y	1	Dn	None
Humphrey's	175.3-178.8	1S	1E	20,21	86.2	Y	47.9	Y	4	Both	86.2	Y	25.0	Y	1	Dn	Irrigation

Table E.8. Continued.

Basin: Upper Colorado River
 Sub-basin: Colorado River
 River: Colorado River

Site	Levy/Berm		Width/ Length (m)	Vegetation		Proximity to Endangered Species		Predominant Flood plain Habitat Type	Land Ownership	Photo Imagery	
	Mainstem Y/N	Bottomland Natural/Man-made		Y/N	Predominant Type(s)	RM	Species			Type*	Catalog No.*
Una North	Y	Man-made	1,097-L	N	—	185.1	CS	Gravel-Pit Pond	Private	ASCS	20: 26; 765: 5
Wallace Creek Island	N	—	—	Y	Cottonwood	185.1	CS	Terrace	Private	ASCS	20: 27
Stoddard	Y	Man-made	810-L	N	—	185.1	CS	Gravel-Pit Pond	Private	ASCS	20: 37; 75: 14
Debeque I-70 Slough	Y	Man-made	769-L	Y	Cottonwood	185.1	CS	Side Channel	Public	ASCS	20: 36-37; 75: 13-14
Latham's	Y	Man-made	155-L	Y	Cottonwood	185.1	CS	Terrace/Ponds	Private	ASCS	21: 2; 75: 15
"No-name"	N	—	—	Y	Cottonwood	185.1	CS	Terrace	Private	ASCS	21: 8; 75: 17
Etter's	Y	Man-made	182-L	Y	Cottonwood/Tamarisk	185.1	CS	Gravel Pit Pond	Private	ASCS	21: 6-7; 75: 18
"No-name"	N	—	—	Y	Cottonwood	185.1	RZ7	Terrace	Public	ASCS	21: 9-10; 75: 21
"No-name"	N	—	—	Y	Tamarisk	185.1	CS	Terrace	Private	ASCS	21: 12; 75-22
Long Point Bottom	N	—	—	Y	Tamarisk	185.1	CS	Terrace	Public	ASCS	21: 16; 75: 24-25
Beavertail	Y	Man-made	305-L	Y	Cottonwood	185.1	CS	Terrace	Private; Public	ASCS	21-19; 78: 4
Island Acres	Y	Man-made	1,463-L	N	—	185.1	CS	Gravel-Pit Ponds	Public	ASCS	21: 23; 78: 7-8
Cameo	Y	Man-made	398-L	Y	Cottonwood	185.1	CS	Terrace	Public	ASCS	21: 24; 78: 9
Palisade	Y	Man-made	1,525-L	Y	Cottonwood	171.0-185.0	CS	Sewage Ponds	Private	ASCS	21: 32; 78: 14
"No-name"	N	—	—	Y	Cottonwood/Tamarisk	171.0-185.1	CS	Terrace	Private	ASCS	21: 32; 78: 14
Labor Camp	N	—	—	N	—	171.0-185.1	CS	Side Channel	Private	ASCS	21: 33; 78: 15
"No-name"	N	—	—	Y	Agricultural Row Crops	171.0-185.1	CS	Terrace; Ponds	Private	ASCS	21: 34; 78: 16
"No-name"	N	—	—	Y	Cottonwood	171.0-185.1	CS	Terrace	Private	ASCS	21: 35; 78: 17
"No-name"	N	—	—	Y	Cottonwood	171.0-185.1	CS	Terrace	Private	ASCS	21: 36; 78: 18
"No-name"	N	—	—	Y	Cottonwood	171.0-185.1	CS	Oxbow	Private	ASCS	22: 1-2; 78: 19-21
Clifton Weiler Treatment	Y	Man-made	1,433-L	Y	Cottonwood	171.0-185.1	CS	Gravel-Pit Pond	Private; Public	ASCS	22: 2-3; 78: 22-23
Clifton Pond	Y	Man-made	792-L	Y	Cottonwood	171.0-185.1	CS	Gravel-Pit Pond	Private; Public	ASCS	22: 4; 78: 24
Corn Lake	Y	Man-made	640-L	Y	Cottonwood	171.0-185.1	CS	Gravel-Pit Pond	Private; Public	ASCS	22: 5-6; 78: 25
Humphrey's	Y	Man-made	1,280-L	Y	Cottonwood	171.0-185.1	CS	Gravel-Pit Pond	Private; Public	ASCS	

* ASCS = Consolidated Farm Service Agency (U. S. Department of Agriculture); COE = U. S. Army Corps of Engineers; U. M. = University of Montana, 1:6000 Colored Aerial Photographs, 4/28/93, 8/2/93
 * Catalog No. = photograph roll number followed by photograph number (e.g. 6: 1-3 = roll 6, pictures 1 through 3).

Table E.9. The 10 parameters used to rank 158 potential floodplain restoration sites important to endangered fishes of the Upper Colorado River Basin.

Basin: Upper Colorado River
 Sub-basin: Colorado
 River: Colorado River

Runoff (May 1993)										Post-Runoff (September 1993)							
Site	Location RM	T	R	S	Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River			Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River			
								(Y/N)	No.	Up/Down Stream				Other Water Sources	(Y/N)	No.	Up/Down Stream
Griffith's	174.1-176.5	1S	1E	20,21 22,28	52.7	Y	24.2	Y	6	Both	Irri- gation	Y	2.6	Y	1	Dn	Irri- gation
"Hotspot Junction" (30-28 Road)	173.9-175.1	1S	1E	20	97.9	Y	11.3	N	1	Dn	Irri- gation Groundwater	N	9.4	N	0	None	Irri- gation
"No-name"	173.4-173.6	1S	1E	19	10.6	Y	1.4	Y	2	Both	Irri- gation Groundwater	Y	0.6	Y	2	Both	Irri- gation
"No-name"	173.0-173.6	1S	1E	19	7.6	N	0	N	0	None	None	N	0	N	0	None	None
Mill Tailing Site	172.5-173.4	1S	1E	19	40.6	Y	1.9	Y	2	Both	Irri- gation	Y	<0.1	N	0	None	Irri- gation
Watson Island	171.0-172.1	1S	1W	23,24	20.6	Y	9.3	Y	4	Both	None	Y	4.6	Y	3	Both	None
"No-name"	171.3-171.6	1S	1W	23	5.9	Y	3.0	Y	2	Both	None	Y	0.4	Y	1	Dn	Irri- gation
"No-name"	170.4-171.0	1S	1W	16,22	16.4	Y	11.2	Y	6	Both	None	N	0	N	0	None	None
Connected Lakes Area	168.7-170.3	1S	1W	6,7,6 9,15,16	220.6	Y	77.2	Y	6	Both	Irri- gation Return Redlands Canal	Y	59.3	Y	3	Dn	Irri- gation Reds Canal
Walter Walker South	164.4-166.0	11S	101W	14	60.3	Y	26.7	Y	6	Both	Irri- gation	Y	3.0	Y	1	Dn	Irri- gation
"No-name"	168.4-169.7	1S	1W	5,6,9 10,15	117.0	Y	30.7	Y	4	Both	Groundwater	Y	20.3	Y	3	Both	None
Appleton Drain East	165.1-166.4	1S	1W	5,6,6	61.1	Y	19.5	N	0	None	Irri- gation	Y	17.0	N	0	None	Irri- gation
Walter Walker SWA	162.7-165.1	1S	1W	6	144.5	Y	17.6	Y	2	Both	None	Y	4.7	Y	2	Dn	None
Panorama	163.1-163.6	11S	101W	14	20.6	Y	3.1	Y	1	Dn	Sewage Treatment Ponds	Y	2.6	Y	1	Dn	Sewage Treat Ponds
"No-name"	161.0-162.7	2N	2W	34,35	20.6	Y	2.5	Y	0	None	None	Y	0.9	Y	1	Dn	None
DuPont Island	161.0-162.0	2N	2W	26,27 34,35	73.9	Y	16.6	Y	3	Both	None	Y	3.6	Y	3	Both	None

Table E.9. Continued.

Basin: Upper Colorado River
 Sub-basin: Colorado River
 River: Colorado River

Site	Levy/Berm		Width/ Length (m)	Vegetation		Proximity to Endangered Species RM	Life Stage	Predominant Flood plain Habitat Type	Land Ownership	Photo Imagery	
	Manstem Y/N	Natural/Man-made		Y/N	Predominant Type(s)					Type*	Catalog No. ^b
Griffith's	Y	Man-made	81-L	Y	Cottonwood/Tamarisk	171.0-185.1	CS	Side Channels Terrace	Private	ASCS	22: 6-7; 78: 25; 79: 1
"Hotspot Junction" (30-29 Road)	Y	Man-made	814-L	N	None	171.0-185.1	CS	Gravel-Pit Ponds	Private; Public	ASCS	22: 7-8; 79: 1-2
"No-name"	Y	Man-made	81-L	Y	Cottonwood	171.0-185.1	CS	Gravel-Pit Pond	Private	ASCS	22: 9; 79: 3
"No-name"	N	—	—	Y	Cottonwood	171.0-185.1	CS	Terrace	Private	ASCS	22: 9; 79: 3
Mill Tailing Site	N	—	—	N	—	171.0-185.1	CS	Terrace	Private	ASCS	22: 10; 79: 4
Watson Island	Y	Man-made	183-L	Y	Cottonwood/Tamarisk	171.0-185.1	CS	Side Channel	Private; Public	ASCS	22: 11-12; 79: 5-6
"No-name"	N	—	—	Y	Tamarisk	171.0-185.1	CS	Side Channel	Private	ASCS	22: 12; 79: 6
"No-name"	N	—	—	Y	Tamarisk	152.1-171.0	CS	Side Channel	Private	ASCS	22: 13-14; 79: 7-8
Connected Lakes Area	Y	Man-made	3,810-L	Y	Cottonwood/Tamarisk	152.1-171.0	CS	Gravel-Pit Ponds	Private; Public	ASCS	22: 15-19; 79: 8-10
Walter Walker South	N	—	—	Y	Tamarisk	152.1-171.0	CS	Terrace	Private; Public	ASCS	22: 20-22; 79: 11-13
"No-name"	Y	Man-made	2,103-L	Y	Tamarisk	152.1-171.0	CS	Gravel-Pit Ponds	Private; Public	ASCS	22: 16-19; 79: 8-10
Appleton Drain East	Y	Man-made	1,798-L	N	—	152.1-171.0	CS	Side Channels	Private	ASCS	22: 19-20; 79: 11
Walter Walker SWA	Y	Man-made	881-L	Y	Tamarisk	152.1-171.0	CS	Gravel-Pit Ponds	Public	ASCS	22: 22-24; 79: 13-15
Panorama	N	—	—	Y	Tamarisk	152.1-171.0	CS	Pond	Private	ASCS	22: 23; 79: 14
"No-name"	N	—	—	Y	Tamarisk/Cottonwood	152.1-171.0	CS	Sewage Ponds	Private	ASCS	22: 25; 79: 16
DuPont Island	N	—	—	Y	Tamarisk/Cottonwood	152.1-171.0	CS	Side Channel	Private	ASCS	22: 26; 79: 16

* ASCS = Consolidated Farm Service Agency (U. S. Department of Agriculture); COE = U. S. Army Corps of Engineers; U. M. = University of Montana, 1:5000 Colored Aerial Photographs, 4/28/83, 8/2/83

^b Catalog No. = photograph roll number followed by photograph number (e.g. 8: 1-3 = roll 8, pictures 1 through 3).

Table E.10. The 10 parameters used to rank 158 potential floodplain restoration sites important to endangered fishes of the Upper Colorado River Basin.

Basin: Upper Colorado River
 Sub-basin: Colorado
 River: Colorado River

Page: 7

Runoff (May 1893)										Post-Runoff (September 1893)									
Site	Location RM	T	R	S	Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River			Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River					
								(Y/N)	No.	Up/Down Stream				Other Water Sources	(Y/N)	No.	Up/Down Stream	Other Water Sources	
DuPont's	159.1-161.9	2N	2W	26.27 28.33	86.8	Y	37.5	Y	4	Both	Irrigation Returns	86.8	Y	25.2	Y	3	Both	Irrigation Returns	
Paul Smith's	158.0-159.1	2N	2W	20.21 28.29	53.4	Y	13.8	Y	1	Dn	Irrigation Return	53.4	Y	2.4	Y	1	Dn	Irrigation Return	
		2N	2W	18.20	17.0	Y	0.8	Y	1	Dn	None	17.0	Y	0.4	Y	1	Up	None	
Fruit 340 Bridge	157.5-158.3	2N	2W	19.20	40.9	Y	16.6	N	0	None	Sewage Ponds	40.9	N	16.5	N	0	None	Sewage Ponds	
Fruit Sewage Ponds Snook's Bottom	155.9-157.1	2N	2W	18.19	38.0	Y	1.4	Y	2	Both	None	38.0	Y	0.6	Y	1	Dn	None	
		1N	3W	13	180.9	Y	4.9	Y	4	Both	Irrigation	180.9	Y	1.5	Y	2	Dn	Irrigation	
Horsethief SWA	151.9-154.7	1N	3W	9.10 11.14, 15	34.0	Y	5.2	N	0	None	None	34	N	4.7	N	0	None	None	
Spann's	151.4-152.4	1N	3W	9.10	27.5	N	0	N	0	None	None	27.5	N	0	N	0	None	None	
		1N	3W	16.17	96.0	N	0	N	0	None	Salt Wash	96.0	N	0	N	0	None	Salt Wash	
Crow Bottom	143.9-146.5	10S	103W	8.17	39.7	N	0	N	0	None	None	39.7	N	0	N	0	None	None	
"No-name"	145.5-146.6	10S	103W	16.17	61.5	N	0	N	0	None	None	61.5	N	0	N	0	None	None	
Vulture Bottom	139.7-142.1	10S	103W	18	23.2	N	0	N	0	None	None	23.2	N	0	N	0	None	None	
"No-name" Island	137.2-138.0	10S	104W	26.27	17.8	Y	0.8	Y	2	Both	None	17.8	N	0	N	0	None	None	
Black Rocks	136.7-137.1	10S	104W	27	52.0	Y	1.0	Y	1	Up	Gully Washers	52.0	N	0	Y	1	Up	Gully Washers	
Knowles Canyon	133.6-135.0	11S	104W	4															
		10S	104W	32.33															

Table E.10. Continued.

Basin: Upper Colorado River
 Sub-basin: Colorado River
 River: Colorado River

Page: 8

Site	Levy/Berm		Width/ Length (m)	Vegetation		Proximity to Endangered Species		Predominant Flood plain Habitat Type	Land Ownership	Photo Imagery	
	Y/N	Mainstem and Bottomland Natural/Man-made		Y/N	Predominant Type(s)	RM	Species Life Stage			Type*	Catalog No.*
DuPont's	Y	Man-made	2,134-L	Y	Tamarisk	152.1-171.0	CS Adult	Gravel-Pit Ponds	Private	ASCS U. M.	22: 28-28; 79: 16-17 115: 121-123
Paul Smith's	Y	Man-made	1,066-L	N	—	152.1-171.0	CS Adult	Gravel-Pit Ponds	Private	ASCS	22: 28-28; 79: 18
Fruit 340 Bridge	N	—	—	N	—	152.1-171.0	CS Adult	Terrace	Private	ASCS U. M.	22: 28 130
Fruit Sewage Ponds	Y	Man-made	1,463-L	N	—	152.1-171.0	CS Adult	Sewage Ponds	Private	ASCS	22: 30-31; 79: 19
Snook's Bottom	N	—	—	Y	Cottonwood/Tamarisk	152.1-171.0	CS Adult	Terrace	Public	N.A.I.	N.A.I. 3103: 10-8 (8/27/86)
Horseshoe SWA	Y	Man-made	3,350-L	Y	Agricultural Row Crops; Cottonwood/Tamarisk	152.1-171.0	CS Adult	Terrace	Public	ASCS	22: 34-36; 79: 22-23
Spann's	Y	Natural	60-L	Y	Tamarisk/Cottonwood	0.0-152.1	CS Adult	Terrace	Private	ASCS	22: 37; 79: 24
"No-name" Crow Bottom	N	—	—	Y	Tamarisk	0.0-152.1	CS Adult	Side Channel	Private	ASCS	23: 1; 79: 25
"No-name"	N	—	—	Y	Cottonwood/Tamarisk	0.0-152.1	CS Adult	Terrace	Private; Public	ASCS	23: 4-5; 80: 6, 8
Vulture Bottom	N	—	—	Y	Cottonwood/Tamarisk	135.0-137.0	HB Adult	Terrace	Private	ASCS	23: 4; 80: 6
"No-name" Island	N	—	—	Y	Cottonwood	135.0-137.0	HB Adult	Terrace	Public	ASCS	23: 10-11; 80: 12-13
Black Rocks	N	—	—	Y	Cottonwood	135.0-137.0	HB Adult	Side Channel	Public	ASCS	23: 14; 80: 15
Knowles Canyon	N	—	—	Y	Cottonwood	135.0-137.0	HB Adult	Terrace	Public	ASCS	23: 15-18; 80: 15
	N	—	—	Y	Tamarisk	135.0-137.0	HB Adult	Side Channel	Public	ASCS	23: 18; 80: 19-20

* ASCS = Consolidated Farm Service Agency (U. S. Department of Agriculture); COE = U. S. Army Corps of Engineers; U. M. = University of Montana; 1:6000 Colored Aerial Photographs, 4/28/83, 9/2/83; N.A.I. = Nichols Associates, Inc., Grand Junction, CO, 1: 24000 Black and White Aerial Photographs.

* Catalog No. = photograph roll number followed by photograph number (e.g. 6: 1-3 = roll 6, pictures 1 through 3).

Table E.11. The 10 parameters used to rank 158 potential floodplain restoration sites important to endangered fishes of the Upper Colorado River Basin.

Page: 9

Basin: Upper Colorado River
 Sub-basin: Colorado
 River: Colorado River

Runoff (May 1993)										Post-Runoff (September 1993)									
Site	Location RM	T	R	S	Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River			Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River			Other Water Sources		
								(Y/N)	No.	Up/Down Stream				(Y/N)	No.	Up/Down Stream			
Joufas Bottom	129.8-133.9	11S	104W	7.8	99.8	N	0	N	0	Up	99.8	N	0	N	0	None	None		
Wildass Canyon Ranch	127.3-131.3	19S	26E	28.30	189.0	Y	14.5	Y	4	Both	189.0	Y	1.9	Y	1	Dn	Irrigation		
		19S	26E	31															
		20S	26E	8															
		20S	25E	12															
"No-name"	125.8-126.8	20S	25E	12.13	54.0	Y	0.8	Y	2	Both	54.0	N	0	N	0	None	None		
Elizondo Ranches	126.5-127.8	20S	25E	1.12	87.1	Y	4.0	N	0	None	87.1	Y	1.7	N	0	None	Groundwater		
		20S	25E	14															
Westwater Wash	124.8-126.0	20S	25E	14	84.0	Y	18.8	Y	8	Both	84.0	Y	0.2	Y	1	Dn	Irrigation		
Clisco Landing Area	107.6-111.8	21S	24E	12.14	267.0	Y	4.0	Y	2	Both	267.0	N	0	N	0	None	Weswater Wash		
		22S	24E	22.26															
Fish Ford Area	103.0-105.9	22S	24E	2.3	100.5	Y	10.8	Y	2	Both	100.5	N	0	N	0	None	Gully Washers		
		22S	24E	10.11															
McGraw/Hotel Bottom	98.1-101.0	22S	24E	21.28	123.7	Y	24.7	Y	6	Both	123.7	Y	7	Y	7	?	?		
White Ranch	77.5-78.4	24S	22E	35	26.1	N	0	N	0	None	26.1	N	0	N	0	None	Castle Creek		
		25S	21E	27															
		25S	21E	1.2															
		25S	21E	28.27															
Counthouse Wash	81.5-84.0	25S	21E	34.35	354.0	Y	212.0	Y	> 8	Both	354.0	Y	81.0	Y	1	Dn	Gully Washer		
Moab Slough		25S	21E	28.27			(estimate)			Groundwater			(estimate)				Mill Creek		
Kane Spring	58.2	26S	20E	21.22	2.8	Y	1.8	Y	2	Both	2.8	N	0	N	0	None	None		
		26S	20E	1.12															
		26S	20E	6.7															
		26S	21E	6.7															
Billboard (Lake Bottom)	50.9-52.5	26S	20E	1.12	36.0	Y	9.9	N	0	None	36.0	N	0	N	0	None	None		

Table E.11. Continued.

Basin: Upper Colorado River
 Sub-basin: Colorado River
 River: Colorado River

Site	Levy/Berm		Vegetation		Proximity to Endangered Species		Predominant Habitat	Land Ownership	Photo Imagery	
	Levy or Berm Dividing Mainstem and Bottomland Y/N	Width/ Length (m)	Y/N	Predominant Type(s)	RM	Species			Type ^a	Catalog No. ^a
Jofitas Bottom	N	—	Y	Tamarisk/Cottonwood	0.0-152.1	CS	Terrace	Private; Public	ASCS	23: 20-22; 80: 21-23
Wildass Canyon Ranch	N	—	Y	Cottonwood/Tamarisk	135.0-137.0	HB	Terrace	Private	ASCS	23: 23-4; 80: 28
					115.0-125.0	HB	Side Channel			
No-name	N	—	Y	Cottonwood	0.0-152.1	CS	Terrace	Private; Public	ASCS	23: 27; 81: 2-3
Elzondo Ranches	Y	Man-made	Y	Agricultural Row Crops	115.0-125.0	HB	Terrace	Private	ASCS	23: 28; 85: 28
Westwater Wash	N	—	Y	Cottonwood	115.0-125.0	HB	Gravel Pit Ponds	Private	ASCS	23: 28; 85: 28
					0.0-152.1	CS	Side Channel	Private	ASCS	23: 28; 81: 3-4
Cisco Landing Area	Y	Man-made	Y	Cottonwood;	115.0-125.0	HB	Terrace	Private; Public	ASCS	24: 5-9; 81: 17-18
				Agricultural Row Crops	115.0-125.0	HB	Side Channel			
Fish Ford Area	N	—	Y	Cottonwood;	0.0-152.1	CS	Terrace	Public	ASCS	24: 12-13; 81: 20-21
				Tamarisk			Side Channel			
McGraw/Hotel Bottom	N	—	Y	Tamarisk	0.0-152.1	CS	Side Channels	Private; Public	ASCS	24: 17-19
White Ranch	Y	915-L	Y	Agricultural Crops	0.0-152.1	CS	Terrace	Private	ASCS	25: 6; 82: 21
Courthouse Wash	Y	27-L	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Private	ASCS	25: 21; 83: 11
Moab Slough	Y	810-L 3,350-L	Y	Tamarisk	0.0-152.1	CS	Depression	Public	ASCS	25: 22-23; 83: 12-13
Kane Spring	Y	30-L	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	25: 27; 83: 17
Billboard (Lake Bottom)	Y	1,500-L	Y	Tamarisk	0.0-152.1	CS	Depression	Public	ASCS	25: 33; 83: 20

^a ASCS = Consolidated Farm Service Agency (U. S. Department of Agriculture); COE = U. S. Army Corps of Engineers; U. M. = University of Montana, 1:6000 Colored Aerial Photographs, 4/28/83, 9/2/83
^b Catalog No. = photograph roll number followed by photograph number (e.g. 6: 1-3 = roll 8, pictures 1 through 3).

Table E.12. The 10 parameters used to rank 158 potential floodplain restoration sites important to endangered fishes of the Upper Colorado River Basin.

Basin: Upper Colorado River
 Sub-basin: Colorado
 River: Colorado River

Page: 11

Runoff (May 1993)										Post-Runoff (September 1993)								
Site	Location RM	T	R	S	Total Potential Inundated Area (ha)	Hydrological Connection to River			Total Area Inundated (ha)	Inundated (Y/N)	Total Potential Inundated Area (ha)	Hydrological Connection to River						
						(Y/N)	No.	Up/Down Stream				(Y/N)	No.	Up/Down Stream				
Jackson Bottom	47.4-49.0	26S	21E	24,25	34.0	Y	13.8	Y	2	Both	Groundwater	Groundwater	N	0	N	0	None	None
"No-name"	44.0-44.7	26S	20E	38	23.0	Y	11.4	N	0	None	Groundwater	Groundwater	N	0	N	0	None	None
"No-name"	42.8-43.7	27S	21E	6	22.0	Y	4.8	N	0	None	Groundwater	Groundwater	N	0	N	0	None	None
"No-name"	41.1-42.1	27S	20E	15	22.3	Y	10.7	N	0	None	Groundwater	Groundwater	N	0	N	0	None	None
"No-name"	40.2-41.3	27S	20E	22,23	21.0	Y	0.4	N	0	None	Groundwater	Groundwater	N	0	N	0	None	None
"No-name"	39.5-40.6	27S	20E	15,22	17.4	Y	1.2	N	0	None	Groundwater	Groundwater	N	0	N	0	None	None
"No-name" Canyon Mouth	38.6	27S	20E	15	1.2	Y	0.4	Y	1	Up	None	None	N	0	N	0	None	None
"No-name"	38.6-39.2	27S	20E	15,18	10.5	Y	0.8	N	0	None	Groundwater	Groundwater	N	0	N	0	None	None
Goose Neck	33.2-35.6	27S	20E	18,19	57.5	Y	10.1	Y	> 6	Both	Groundwater	Groundwater	N	0	N	0	None	None
Shaler Canyon Mouth	34.6	27S	20E	18	5.6	Y	1.2	Y	1	Up	Gully Washer	Gully Washer	N	0	N	0	None	Gully Washer
"No-name" Canyon Mouth	32.1	27S	20E	29	1.6	Y	0.4	Y	1	Up	Gully Washer	Gully Washer	N	0	N	0	None	Gully Washer
"No-name" Canyon Mouth	31.6	27S	20E	28	1.6	Y	0.2	Y	1	Up	Gully Washer	Gully Washer	N	0	N	0	None	Gully Washer
"No-name"	30.8-31.9	27S	20E	26,29	27.9	Y	1.9	N	0	None	Groundwater	Groundwater	N	0	N	0	None	None
Little Bridge Canyon	30.0	27S	20E	30	1.6	Y	0.8	Y	1	Up	Gully Washer	Gully Washer	N	0	N	0	None	Gully Washer
Lockhart Canyon	29.5	28S	20E	16	5.6	Y	0.8	Y	1	Up	Gully Washer	Gully Washer	N	0	N	0	None	Gully Washer
"No-name"	25.3-26.5	26S	20E	19,20	31.0	N	7.7	N	0	None	Groundwater	Groundwater	N	0	N	0	None	None
Lathrop Canyon	23.5	28S	18E	13	4.5	Y	0.4	Y	1	Up	Gully Washer	Gully Washer	N	0	N	0	None	Gully Washer
Buck Canyon	22.7	28S	18E	13	4.5	Y	0.6	Y	1	Up	Gully Washer	Gully Washer	N	0	N	0	None	Gully Washer
Gooseberry Canyon	21.7	28S	18E	24	2.6	Y	0.4	Y	1	Up	Gully Washer	Gully Washer	N	0	N	0	None	Gully Washer
Dogleg Canyon	21.2	28S	18E	25	3.2	Y	0.4	Y	1	Up	Gully Washer	Gully Washer	N	0	N	0	None	Gully Washer
Sheep Bottom	17.5-18.7	28S	20E	30	27.3	Y	0.8	N	0	None	Groundwater	Groundwater	N	0	N	0	None	None
Indian Creek Canyon Mouth	16.5	28S	18E	12	4.5	Y	0.8	Y	1	Up	Gully Washer	Gully Washer	N	0	N	1	Up	Gully Washer
Monument Creek Mouth	15.3	28S	18E	10	5.3	Y	0.2	N	0	None	Gully Washer	Gully Washer	N	0	Y	1	Up	Gully Washer
"No-name" Canyon Mouth	11.9	28S	18E	26	3.2	Y	< 0.1	N	0	None	Gully Washer	Gully Washer	N	0	Y	1	Up	Gully Washer

Table E.12. Continued.

Basin: Upper Colorado River
 Sub-basin: Colorado River
 River: Colorado River

Page: 12

Site	Levy/Berm		Vegetation		Proximity to Endangered Species		Predominant Habitat Type	Land Ownership	Photo Imagery	
	Levy or Berm Mainstem and Bottomland Y/N	Width/ Length (m)	Y/N	Predominant Type(s)	RM	Species			Type ^a	Catalog No. ^b
Jackson Bottom	N	—	Y	Tamarisk	0.0-152.1	CS	Depression	Private	ASCS	26: 3; 83: 25
"No-name"	Y	1,220-L	N	—	0.0-152.1	CS	Gravel-Pit Ponds	Private; Public	ASCS	26: 6-7; 84: 4
"No-name"	Y	1,859-L	Y	Tamarisk	0.0-152.1	CS	Depression	Public	ASCS	26: 8; 84: 7
"No-name"	Y	1,200-L	Y	Tamarisk	0.0-152.1	CS	Depression	Public	ASCS	26: 8; 84: 8
"No-name"	N	—	Y	Tamarisk	0.0-152.1	CS	Depression	Public	ASCS	26: 11; 84: 10
"No-name"	N	—	Y	Tamarisk	0.0-152.1	CS	Depression	Public	ASCS	26: 11; 84: 11
"No-name" Canyon Mouth	N	—	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	26: 11; 84: 12
"No-name"	Y	1,400-L	Y	Tamarisk	0.0-152.1	CS	Depression	Public	ASCS	26: 11; 84: 13
Goose Neck	Y	4,000-L	Y	Tamarisk	0.0-152.1	CS	Depression	Public	ASCS	26: 15-17; 84: 17
Shafter Canyon Mouth	N	—	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	26: 15; 84: 15
"No-name" Canyon Mouth	N	—	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	26: 18
"No-name" Canyon Mouth	N	—	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	26: 19; 84: 19
"No-name"	Y	1,737-L	Y	Tamarisk	0.0-152.1	CS	Depression	Public	ASCS	26: 19; 84: 18
Little Bridge Canyon Mouth	N	—	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	26: 20; 84: 21
Lockhart Canyon Mouth	N	—	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	26: 23; 84: 25
"No-name"	Y	2,200-L	Y	Tamarisk	0.0-152.1	CS	Depression	Public	ASCS	26: 23; 84: 25
Lathrop Canyon Mouth	N	—	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	26: 24; 85: 2
Buck Canyon Mouth	N	—	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	26: 25; 85: 3
Gooseberry Canyon Mouth	N	—	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	26: 26; 85: 5
Dogleg Canyon Mouth	N	—	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	26: 26; 85: 5
Sheep Bottom	Y	2,100-L	Y	Tamarisk	0.0-152.1	CS	Depression	Public	ASCS	26: 29; 85: 10
Indian Creek Canyon Mouth	N	—	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	26: 30; 85: 12
Monument Creek Canyon Mouth	N	—	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	26: 33; 85: 14
"No-name" Canyon Mouth	N	—	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	26: 36; 85: 20

^a ASCS = Consolidated Farm Service Agency (U. S. Department of Agriculture); COE = U. S. Army Corps of Engineers; U. M. = University of Montana; 1:8000 Colored Aerial Photographs, 4/28/83, 8/2/83
^b Catalog No. = photograph roll number followed by photograph number (e.g. 8; 1-3 = roll 8, pictures 1 through 3).

Table E.13. The 10 parameters used to rank 158 potential floodplain restoration sites important to endangered fishes of the Upper Colorado River Basin.

Basin: Upper Colorado River
 Sub-basin: Colorado
 River: Colorado River

Page: 13

Runoff (May 1993)										Post-Runoff (September 1993)									
Site	Location			Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River			Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River			Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Other Water Sources
	RM	T	R	S			(Y/N)	No.	Up/Down Stream				(Y/N)	No.	Up/Down Stream				
No-name Canyon Mouth	10.1	28S	18E	28	0.4	Y	Y	1	Up	0.4	N	0	Y	1	Up	0.4	N	0	Gully Washer
Salt Creek Canyon Mouth	3.4	28.5S	18E	33	2.8	Y	Y	1	Up	2.8	N	0	Y	1	Up	2.8	N	0	Gully Washer
Elephant Creek Canyon Mouth	3.0	29.5S	18E	33	1.7	Y	Y	1	Up	1.7	N	0	Y	1	Up	1.7	N	0	Gully Washer

Table E.13. Continued.

Basin: Upper Colorado River
 Sub-basin: Colorado River
 River: Colorado River

Site	Levy/Berm		Width/ Length (m)	Vegetation		Proximity to Endangered Species		Predominant Floodplain Habitat Type	Land Ownership	Photo Imagery	
	Levy or Berm Dividing Mainstem and Bottomland Y/N	Natural/Man-made		Y/N	Predominant Type(s)	RM	Species			Type*	Catalog No.*
No-name Canyon Mouth	N	---	---	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	27: 4; 85: 22
Salt Creek Canyon Mouth	N	---	---	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	27: 12; 86: 1
Elephant Creek Canyon Mouth	N	---	---	Y	Tamarisk	0.0-152.1	CS	Canyon Mouth	Public	ASCS	27: 2; 86: 1

* ASCS = Consolidated Farm Service Agency (U. S. Department of Agriculture); COE = U. S. Army Corps of Engineers; U. M. = University of Montana. 1:6000 Colored Aerial Photographs, 4/28/83, 9/2/83

* Catalog No. = photograph roll number followed by photograph number (e.g. 6: 1-3 = roll 6, pictures 1 through 3).

Table E.14. The 10 parameters used to rank 158 potential floodplain restoration sites important to endangered fishes of the Upper Colorado River Basin.

Basin: Upper Colorado River
 Sub-basin: Colorado
 River: Gunnison River

Runoff (May 1993)										Post-Runoff (September 1993)									
Site	Location RM	T	R	S	Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River			Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	(Y/N)	No.	Up/Down Stream	Other Water Sources	Springs	Gully/Washer
								(Y/N)	No.	Up/Down Stream									
"No-name"	74.3-74.7	14S	94W	38	24.0	N	0	N	0	None	24.0	N	0	N	0	None	None	None	None
Lawhead Gulch Bottom	70.1-71.8	15S	94W	1	13.8	Y	0.4	Y	1	Up	13.8	N	0	N	0	None	None	None	None
Ferganchick's	69.6-70.5	15S	94W	4	16.7	N	0	Y	1	Up	18.7	N	0	N	0	None	None	None	None
"No-name"	68.3-69.4	15S	94W	4.9	12.1	N	0	Y	1	Up	12.1	N	0	N	0	None	None	None	None
"No-name"	67.5-68.2	15S	94W	5	3.8	N	0	N	0	None	3.8	N	0	N	0	None	None	None	None
Austin County Bridge	65.7-68.4	15S	94W	8	17.2	N	0	N	0	None	17.2	N	0	N	0	None	None	None	None
Austin Hwy 92 Bridge	65.0-65.5	15S	95W	1	13.5	Y	4.1	Y	1	Up	4.1	N	0	N	0	None	None	None	None
"No-name"	63.3-65.3	15S	95W	1	52.2	Y	0.8	Y	1	Up	52.2	N	0	N	0	None	None	None	None
"No-name"	63.0-64.8	15S	95W	2.11	17.7	N	0	N	0	None	17.1	N	0	N	0	None	None	None	None
Colorado Hwy 65 Bridge	62.6-63.2	15S	95W	3	14.4	Y	9.9	Y	1	Up	14.4	N	0	N	0	None	None	None	None
Tongue Creek	81.5-82.5	15S	95W	3.4	35.6	Y	18.2	Y	1	Dn	35.6	Y	0	N	0	None	Tongue Creek	None	None
Hutchin's	60.1-60.8	15S	95W	5	17.3	Y	4.8	Y	2	Dn	17.3	Y	1.6	N	0	None	None	None	None
North Delta	57.6-59.5	15S	95W	7.8	49.3	Y	14.0	Y	2	Up	49.3	N	0	N	0	None	None	None	None
South Delta	57.6-58.8	15S	95W	7.8	39.9	Y	10.0	N	0	None	39.9	Y	9.9	N	0	None	None	None	None
Confluence Park	56.7-57.7	15S	96W	12.13	40.1	Y	8.4	Y	2	Both	40.1	Y	6.7	Y	1	Dn	None	None	None
Uncompahgre R. Confluence	56.3-57.0	15S	96W	10.15	34.4	Y	8.1	Y	1	Up	34.4	Y	8.3	Y	1	Up	Ponds	None	None
Delta City Sewage Plant	55.6-56.5	15S	96W	10.11	38.0	Y	14.7	Y	1	Up	38.1	Y	3.6	Y	1	Up	None	None	None
"No-name"	54.6-55.5	15S	96W	10.15	27.9	Y	1.9	Y	1	Up	27.9	Y	1.5	Y	1	Up	None	None	None
"No-name"	54.2-55.1	15S	96W	15	28.2	Y	9.8	Y	2	Both	28.2	Y	2.2	Y	1	Dn	None	None	None
"No-name"	53.1-54.1	15S	96W	15.16	21.7	Y	1.3	N	0	None	21.7	N	0	N	0	None	None	None	None

Table E.14. Continued.

Basin: Upper Colorado River
 Sub-basin: Colorado River
 River: Gunnison River

Page: 16

Site	Levy/Berm		Width/ Length (m)	Vegetation		Proximity to Endangered Species RM	Species	Life Stage	Predominant Habitat Type	Land Ownership	Photo Imagery	
	Meinstem Y/N	Levy or Berm Dividing Natural/Man-made		Y/N	Predominant Type(s)						Type*	Catalog No.*
"No-name"	N	—	—	Y	Tamarisk	59.1	CS	Adult	Terrace	Private	B.R.	12: 3
Lawhead Gulch Bottom	N	—	—	Y	Cottonwood/Tamarisk	59.1	CS	Adult	Terrace	Private	ASCS	71: 2,3 12: 6
Ferganchick's	Y	Man-made	762-L	Y	Fruit Trees	59.1	CS	Adult	Terrace	Private	ASCS	71: 6
"No-name"	Y	Man-made	2,042-L	Y	Fruit Trees	59.1	CS	Adult	Terrace	Private	ASCS	71: 6
"No-name"	N	—	—	Y	Fruit Trees	59.1	CS	Adult	Terrace	Private	ASCS	71: 8
Austin County Bridge	N	—	—	Y	Cottonwood	59.1	CS	Adult	Terrace	Private	B.R.	13: 5; 1: 1
Austin Hwy 92 Bridge	N	—	—	Y	Tamarisk	59.1	CS	Adult	Terrace	Private	B.R.	13: 7; 2: 2
"No-name"	N	—	—	Y	Agricultural	59.1	CS	Adult	Terrace	Private	B.R.	13: 8; 2: 3
"No-name"	N	—	—	Y	Cottonwood	59.1	CS	Adult	Terrace	Private	B.R.	14: 4; 2: 4
Colorado Hwy 65 Bridge	N	—	—	Y	Cottonwood	59.1	CS	Adult	Terrace	Private	B.R.	14: 5; 2: 5
Tongue Creek	N	—	—	Y	Cottonwood	59.1	CS	Adult	Terrace	Private	B.R.	14: 7; 3: 1
Hutchin's	Y	Man-made	457-L	Y	Cattail/Bulrush	59.1	CS	Adult	Depression	Private	B.R.	14: 7; 3: 2
North Delta	Y	Man-made	894-L	Y	Cottonwood	3.1-59.1	CS	Adult	Gravel-Pit Ponds	Private	B.R.	14: 11; 3: 4
South Delta	Y	Man-made	1,523-L	N	—	3.1-59.1	CS	Adult	Gravel-Pit Ponds	Private	B.R.	15: 7; 3: 5
Confluence Park	Y	Natural	—	Y	Cottonwood	3.1-59.1	CS	Adult	Oxbow	Private; Public	B.R.	15: 1; 4: 4
Uncompahgre R. Confluence	Y	Man-made	460-L	N	—	3.1-59.1	CS	Adult	Ponds	Public	B.R.	15: 3; 4: 5
Delta City Sewage Plant	Y	Man-made	381-L	Y	Cottonwood/Agr Crops	3.1-59.1	CS	Adult	Terrace	Private	B.R.	15: 3; 4: 5 15: 5; 4: 6
"No-name"	N	—	—	Y	Cottonwood	3.1-59.1	CS	Adult	Terrace	Private	B.R.	15: 8; 4: 8
"No-name"	N	—	—	Y	Cottonwood	3.1-59.1	CS	Adult	Terrace	Private	B.R.	15: 7; 4: 8
"No-name"	N	—	—	Y	Cottonwood	3.1-59.1	CS	Adult	Terrace	Private	B.R.	15: 8; 4: 9

* ASCS = Consolidated Farm Service Agency (U. S. Department of Agriculture); COE = U. S. Army Corps of Engineers; U. M. = University of Montana, 1:5000 Colored Aerial Photographs, 4/28/93, 9/2/93; B.R. = U. S. Bureau of Reclamation, 1:12000 Colored Infrared Photographs, 6/2/93, 9/28/93.

* Catalog No. = photograph roll number followed by photograph number (e.g. 6: 1-3 = roll 6, pictures 1 through 3).

Table E.15. The 10 parameters used to rank 158 potential floodplain restoration sites important to endangered fishes of the Upper Colorado River Basin.

Runoff (May 1993)										Post-Runoff (September 1993)									
Site	Location	RM	T	R	S	Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River			Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River				
									(Y/N)	No.	Up/Down Stream				Other Water Sources	(Y/N)	No.	Up/Down Stream	Other Water Sources
Johnson Boy's Slough Escalante SWA North	53.2-54.2	15S	96W	15,18		63.1	Y	9.9	Y	2	Both	None	63.1	Y	4.1	Y	1	Dn	None
	50.8-52.9	15S	96W	17,18	19	110.5	Y	28.7	Y	1	Dn	Irrigation	110.5	Y	8.0	Y	1	Up	Irrigation
Escalante SWA South	50.2-52.4	15S	96W	17,19	20	82.8	Y	48.8	Y	2	Both	Irrigation	82.8	Y	4.5	Y	1	Dn	Irrigation
'Blue Duck' Bottom 'No-name'	49.5-50.4	15S	96W	19		21.7	Y	6.2	Y	1	Up	None	21.7	N	0	N	0	None	None
	49.4-49.7	15S	96W	19	24	11.2	Y	3.5	Y	1	Up	None	11.2	N	0	N	0	None	None
'No-name' 'No-name' 'No-name'	48.6-49.3	15S	97W	24		14.1	Y	3.5	Y	1	Up	None	14.1	N	0	N	0	None	None
	46.6-47.5	15S	97W	14		14.2	Y	4.9	Y	2	Both	None	14.2	N	0	N	0	None	None
	41.7-42.8	15S	97W	10		15.1	Y	2.5	Y	2	Both	Escalante Creek	15.1	N	0	N	0	None	Escalante Ck
Escalante Ranches 'No-name'	39.9-41.0	4S	3E	33		33.8	Y	18.4	Y	2	Both	None	33.8	N	0	N	0	None	Irrigation
	36.4-36.9	14S	98W	27		11.0	Y	2.8	Y	4	Both	None	11.0	Y	1.3	Y	2	Both	None
	35.6-36.0	14S	98W	27		25.0	Y	9.4	Y	1	Up	None	25.0	N	0	N	0	None	None
Broughton McKendrick's People's South (VIP Camp)	34.8-36.3	14S	98W	22		44.9	Y	3.5	Y	2	Both	None	44.9	N	0	N	0	None	None
	34.0-35.2	14S	98W	22		23.1	Y	2.9	Y	2	Dn	None	23.1	N	0	N	0	None	None
	34.6-34.9	14S	98W	22		8.8	Y	5.7	Y	4	Both	None	8.8	Y	2.2	Y	2	Both	None
Sand Flat Tunnel Point Dads Flat	27.3-28.7	3S	2E	33		14.5	Y	0.4	N	0	None	None	14.5	N	0	N	0	None	None
	26.3-26.9	14S	99W	1,12		6.6	Y	2.8	Y	1	Up	None	6.6	Y	0	N	0	None	None
	24.7-26.3	14S	99W	1		38.9	Y	0.2	Y	1	Dn	None	38.9	N	0	N	0	None	None
Deer Run 'No-name' 'No-name'	23.0-23.6	13S	99W	35		9.9	Y	1.5	Y	2	Both	None	9.9	Y	0.2	Y	1	Dn	None
	22.1-22.7	13S	99W	26,27		10.0	N	0	N	0	None	None	10.0	N	0	N	0	None	None
	21.3-21.8	13S	99W	22,27		9.3	Y	5.5	Y	2	Both	None	9.3	Y	3.4	Y	2	Both	None
'No-name'	18.4-19.2	13S	99W	10,15		4.4	Y	3.5	Y	2	Both	None	4.4	N	0	N	0	None	None

Basin: Upper Colorado River
Sub-basin: Colorado
River: Gunnison River

Page: 11

Table E.15. Continued.

Basin: Upper Colorado River
 Sub-basin: Colorado River
 River: Gunnison River

Site	Levy/Berm		Width/ Length (m)	Vegetation		Proximity to Endangered Species		Predominant Habitat Type	Land Ownership	Photo Imagery	
	Levy or Berm Mainstem and Bottomland	Y/N		Y/N	Predominate Type(s)	RM	Species			Type ^a	Catalog No. ^b
Johnson Boy's Slough	Y	Y	725-L	Y	Greasewood	3.1-59.1	CS	Oxbow	Private	B.R.	15; 8; 4; 9
Escalante SWA North	Y	Y	814-L	Y	Cottonwood; Greasewood	3.1-59.1	CS	Oxbow; Side Channels	Public	B.R.	15; 9-10; 5; 4-5
Escalante SWA South	N	—	—	Y	Cottonwood	3.1-59.1	CS	Side Channels	Public; Private	B.R.	15; 9-10; 5; 4-5
Blue Duck Bottom	N	—	—	Y	Cottonwood	3.1-59.1	CS	Terrace	Public	B.R.	18; 3; 5; 7
No-name	N	—	—	Y	Cottonwood	3.1-59.1	CS	Terrace	Public	B.R.	18; 3; 5; 7
No-name	N	—	—	Y	Cottonwood	3.1-59.1	CS	Terrace	Public	B.R.	18; 4; 8; 4
No-name	N	—	—	Y	Cottonwood	3.1-59.1	CS	Terrace	Private	B.R.	18; 7; 6; 5
Escalante Ranches	Y	Y	81-L	Y	Cottonwood	3.1-59.1	CS	Terrace	Private	B.R.	18; 12; 6; 10
No-name	N	—	—	Y	Cottonwood	3.1-59.1	CS	Terrace	Private; Public	B.R.	18; 4; 6; 12
Donihuez	N	—	—	Y	Tamarisk	3.1-59.1	CS	Terrace	Private	B.R.	18; 9; 8; 8
Broughton	N	—	—	Y	Cottonwood/Tamarisk	3.1-59.1	CS	Terrace	Private	B.R.	18; 10; 8; 9
McKendrick's	Y	Y	1,524-L	Y	Fruit Trees	3.1-59.1	CS	Terrace	Private	B.R.	18; 11; 7; 2
People's South (VIP Camp)	N	—	—	Y	Cottonwood/Tamarisk	3.1-59.1	CS	Terrace	Private	B.R.	18; 12; 7; 2
People's Island	N	—	—	Y	Cottonwood/Tamarisk	3.1-59.1	CS	Terrace	Private; Public	B.R.	18; 11; 8; 11
Sand Flat	N	—	—	Y	Cottonwood/Tamarisk	3.1-59.1	CS	Terrace	Private	B.R.	18; 17; 8; 18
Tunnel Point	N	—	—	Y	Cottonwood/Tamarisk	3.1-59.4	CS	Terrace	Public	B.R.	18; 19; 8; 18
Dads Flat	N	—	—	Y	Cottonwood/Sagebrush	3.1-59.4	CS	Terrace	Private	B.R.	18; 19; 8; 18
Deer Run	N	—	—	Y	Tamarisk/Cottonwood	3.1-59.1	CS	Terrace	Public	B.R.	20; 3; 10; 4
No-name	N	—	—	Y	Cottonwood/Tamarisk	3.1-59.1	CS	Terrace	Public	B.R.	20; 4; 10; 5
No-name	N	—	—	Y	Tamarisk	3.1-59.1	CS	Terrace	Public	B.R.	20; 5; 10; 6
No-name	N	—	—	Y	Tamarisk/Cottonwood	3.1-59.1	CS	Terrace	Private; Public	B.R.	21; 2; 11; 3

*ASCS = Consolidated Farm Service Agency (U. S. Department of Agriculture); COE = U. S. Army Corps of Engineers; U. M. = University of Montana; 1:6000 Colored Aerial Photographs, 4/28/83, 9/28/83; B.R. = U. S. Bureau of Reclamation, 1:12000 Colored Infrared Photographs, 6/2/83, 9/28/83.

^a Catalog No. = photograph roll number followed by photograph number (e.g. 6; 1-3 = roll 6, pictures 1 through 3).

Table E.16. The 10 parameters used to rank 158 potential floodplain restoration sites important to endangered fishes of the Upper Colorado River Basin.

Basin: Upper Colorado River
 Sub-basin: Colorado
 River: Gunnison River

Runoff (May 1993)										Post-Runoff (September 1993)								
Site	Location RM	T			Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River			Total Potential Inundated Area (ha)	Inundated (Y/N)	Total Area Inundated (ha)	Hydrological Connection to River				
		R	S					(Y/N)	No.	Up/Down Stream				Other Water Sources	(Y/N)	No.	Up/Down Stream	Other Water Sources
Kannah Creek	18.0-18.6	13S	99W	9,10	13.0	Y	1.2	Y	2	Both	Kannah Creek	13.0	Y	0.4	Y	2	Both	Kannah Crk
Whitewater Blding Materials	13.3-18.0	12S	99W	28	70.0	Y	10.3	Y	1	Up	East Creek	70.0	N	0	N	0	None	East Crk
		2S	1E	14,15	32.0	Y	19.6	Y	1	Up	None	32.0	N	0	N	0	None	None
		2S	1E	15														
Duck Blind	12.7-13.8	12S	99W	28	38.5	Y	13.6	N	0	None	Bangs Canyon	38.5	Y	7.3	N	0	None	Bangs Cyn
Schroeder's Mule Farm	4.5-6.1 3.1-3.9	1S	1W	35,38	28.1	Y	10.3	Y	3	Both	Ponds	28.1	Y	0.8	N	0	None	Ponds
		1S	1W	35	31.0	Y	12.2	N	0	None	None	31.0	N	0	N	0	None	None
		Redlands	2.2-3.0	1S	1W	26,27	4.8	Y	1.3	N	0	None	Canal Seepage	4.8	N	0	N	0

Table E.16. Continued.

Basin: Upper Colorado River
 Sub-basin: Colorado River
 River: Gunnison River

Site	Levy/Berm		Width/ Length (m)	Vegetation		Proximity to Endangered Species		Predominant Flood plain Habitat Type	Land Ownership	Photo Imagery	
	Levy or Berm Mainstem and Bottomland V/N	Dividing Natural/Man-made		V/N	Predominant Type(s)	RM	Species Life Stage			Type*	Catalog No.*
Kannah Creek	Y	Man-made	610-L	Y	Tamarisk	3.1-59.1	CS Adult	Canyon Mouth	Private	B.R.	21: 2; 11: 3
Whitewater Biding Materials	Y	Man-made	1,523-L	Y	Tamarisk/Cottonwood	3.1-59.1	CS Adult	Gravel-Pit Ponds	Private	B.R.	21: 7; 11: 8
Duck Blind	Y	Man-made	305-L	Y	Cottonwood/Tamarisk	3.1-59.1	CS Adult	Terrace	Private	B.R.	21: 7; 11: 8
Bangs Canyon	Y	Man-made	640-L	N	—	3.1-59.1	CS Adult	Gravel-Pit Pond	Private	B.R.	22: 3; 12: 2
Schroeder's Mule Farm	Y Y	Man-made Natural	610-L 1,311-L	Y Y	Tamarisk/Cottonwood Cottonwood; Agricultural Row Crops	3.1-59.1 3.1-59.1	CS Adult CS Adult	Gravel-Pit Pond Terrace	Private Private	B.R. B.R.	22: 11; 12: 10 23: 4; 12: 12
Redlands	N	—	—	Y	Tamarisk	0.7-3.0 CO 152.1-185.1	CS Adult CS Adult	Terrace	Private	B.R.	23: 6; 13: 5

* ASCS = Consolidated Farm Service Agency (U. S. Department of Agriculture); COE = U. S. Army Corps of Engineers; U. M. = University of Montana; 1:6000 Colored Aerial Photographs, 4/28/83; 8/2/83; B.R. = U. S. Bureau of Reclamation, 1:12000 Colored Infrared Photographs, 8/2/83, 9/28/83.

* Catalog No. = photograph roll number followed by photograph number (e.g. 6: 1-3 = roll 6, pictures 1 through 3).

APPENDIX F
Scoring and ranking of bottomland sites
using selection criteria

Table F.1. Selection criteria and associated sub-criteria raw score values multiplied by the average weighted point values to produce the total score and rank for each bottomland habitat site in the Green, Yampa, and White rivers, Colorado and Utah, 1993. Top 11 ranked bottomland habitats denoted by ▶

Site Description	River Mile	Sub-Criteria Raw Score Values*			Sub-Criteria Raw Score Values multiplied by Average Weighted Point Values			Total Score ^a	Rank ^d		
		Status ownership	Proximity bar	June to river	Potential connection of sites	Status ownership	Proximity bar			June to river	Potential connection of sites
Green River											
Sears Cr.-Allen Ranch, Browns Park NWR	388.0-392.5	1	1	3	4	1	12	28	56	13	
Crouse Creek Browns Park NWR	381.5-382.5	3	1	3	4	3	12	28	58	12	
Butch Cassidy Lake Browns Park NWR	379.5-380.5	3	1	3	4	3	12	28	58	12	
Hog Lake Browns Park NWR	376.5-378.0	3	1	3	4	3	12	28	48	14	
Warren Bottom, Browns Park NWR	375.5-377.0	3	1	3	4	3	12	28	58	12	
Spitzie Bottom, Browns Park NWR	375.0-376.0	3	1	3	4	3	12	28	48	14	
J S Hoy Bottom, Browns Park NWR	372.0-374.0	3	1	3	4	3	12	28	58	12	
Grimes Bottom Browns Park NWR	367.5-369.5	3	1	3	4	3	12	28	58	12	
Escalante Ranch	302.5-309.5	1	5	1	3	1	60	5	87	4 ^a	
Merill Snow Ranch	302.0-303.0	1	5	1	3	1	60	5	21	87	4
Gravel Ponds at Jensen	301.0-302.0	1	5	1	3	1	60	5	21	87	4
Stewart Lake	299.0-300.0	2	5	1	3	2	60	5	21	88	3
Ashley Creek confluence area	297.0-298.5	2	5	1	3	2	60	5	21	88	3
Little Stewart Lake	295.5-297.5	3	5	1	3	3	60	5	21	89	2
Spring Hollow	295.0-296.0	1	5	1	3	1	60	5	21	87	4
Walker Hollow	294.0-295.0	1	5	1	2	1	60	5	14	80	6
Alhambra Ferry Site	292.0-294.0	1	5	1	2	1	60	5	14	80	6
Gravel Pits	292.0-293.0	1	5	1	2	1	60	5	14	80	6
Bonanza Bridge Area	288.5-289.0	1	5	1	2	1	60	5	14	82	5
Collier Draw	285.5-286.5	3	5	1	2	3	60	5	14	82	5
Horseshoe Bend Area	277.0-284.0	1	4	1	1	1	48	5	7	61	11
The Stirrup	274.0-277.0	1	4	1	1	1	48	5	7	63	10
Hamacker Bottom/Baesser Bend	271.0-274.0	1	4	1	3	1	48	5	21	75	9
Downstream of Baesser Bend	269.0-272.0	3	4	1	3	3	48	5	21	77	8
Upstream of Brennan Bottom	267.0-269.0	3	4	1	3	3	48	5	21	77	8
Upstream of Brennan Bottom	266.0-267.0	3	4	1	3	3	48	5	21	77	8
Brennan Bottom	262.0-266.0	1	4	1	5	1	48	5	35	89	2
Johnson Bottom, Ouray NWR	263.0-265.0	3	4	3	5	3	48	15	35	101	1
Leota Pond Complex, Ouray NWR	257.0-262.0	3	4	3	5	3	48	15	35	101	1
Wascket Lake, Ouray NWR	253.0-257.0	3	3	3	5	3	36	15	35	89	2
Sheppard Bottom, Ouray NWR	254.0-256.0	3	3	3	5	3	36	15	35	89	2
Old Charlie Wash (main)/Woods Bottom, Ouray NWR	249.0-252.0	3	3	3	5	3	36	15	35	89	2
Old Charlie Wash (diked)/Woods Bottom, Ouray NWR	249.0-250.0	3	3	3	5	3	36	15	35	89	2
Ouray Ute pasture land	248.0-251.0	1	3	1	5	1	36	5	35	77	8
Ducheanne River confluence area	248.0-249.0	1	3	1	5	1	36	5	35	77	8
White River confluence area	248.0-249.0	1	3	1	5	1	36	5	35	77	8
West Branch area	243.0-247.0	1	3	1	5	1	36	5	35	77	8
Tia Juana Bottom	242.0-244.0	1	3	1	5	1	36	5	35	77	8
Mouth of Willow Creek	238.0-241.0	3	3	3	5	3	36	5	35	79	7
Pariette Draw	237.0-241.0	3	3	3	5	3	36	5	35	79	7
Desert Spring/Moon Bottom	227.0-228.0	3	2	1	1	3	24	5	7	39	17
Fourmile Wash/King Bottom	226.0-227.0	3	2	1	1	3	24	5	7	39	17
Indian Pasture/Hydes Bottom	224.0-225.0	3	2	1	1	3	24	5	7	39	17
Rays and Long Bottom	220.0-222.0	3	2	1	1	3	24	5	7	39	17
Between Long and Boat Bottom	219.0-220.0	3	2	1	1	3	24	5	7	39	17
Nine Mile Creek	213.0-214.0	3	2	1	1	3	24	5	7	39	17

*Each selection criteria is further sub-divided into sub-criteria and given a raw score value (see methods page ?).
^aEqual to the sub-criteria raw score value * the average weighted point value (e.g., for Sears Creek-Allen Ranch this equals 1*1=1 for the status of land ownership criteria, 1*12=12 for the proximity to the spawning bar criteria, 3*5=15 for the June connection to the river criteria, and 4*7=28 for the potential of network of sites criteria).
^bEqual to the sum of the sub-criteria raw score values * the average weighted point values (e.g., for Sears Creek-Allen Ranch this equals 1+12+15+28=56).
^cThe total score was sorted from lowest to highest with the highest total score receiving the lowest rank and lowest total score the highest rank. The lowest rank was the top priority and the highest rank the bottom priority.

Table F.1. Continued.

Site Description	River Mile	Sub-Criteria Raw Score Values*			Sub-Criteria Raw Score Values multiplied			Rank ^d		
		Status of land ownership	Proximity bar to river	June Potential to river of sites	BY Average Weighted Point Values					
					Status ownership	Proximity bar to river	June Potential to river of sites			
Green River										
Nutter Hole	211.0-213.0	3	2	1	3	24	5	7	39	17
Duches Hole	209.0-211.0	3	2	1	3	24	5	7	39	17
Tabyago Canyon/Little Horse	206.0-208.0	3	2	1	3	24	5	7	39	17
Little Horse Bottom	204.0-207.0	1	2	1	1	24	5	7	37	18
Downstream Maverick Canyon	203.0-204.0	1	2	1	1	24	5	7	37	18
Gold Hole	201.0-202.0	1	2	1	1	24	5	7	37	18
Rock House Bottom	200.0-201.0	3	2	1	3	24	5	7	39	17
Little Rock Canyon	199.0-200.0	3	2	1	3	24	5	7	39	17
Stampede Flat	198.0-199.0	1	2	1	1	24	5	7	37	18
Hoodoo Forms	197.0-198.0	1	2	1	1	24	5	7	37	18
Between Snap-Three Canyons	170.0-171.0	3	2	1	3	24	5	7	39	17
Chandler Falls Canyon	165.0-167.0	3	2	1	3	24	5	7	39	17
Trail and Bull Canyon Rapids	164.5-165.5	3	2	1	3	24	5	7	39	17
Joe Hutch Canyon/Rapids	160.0-161.0	3	2	1	3	24	5	7	39	17
Florence Creek/Three Forks	157.0-158.0	3	2	1	3	24	5	7	39	17
Downstream of Three Forks	153.5-155.0	3	2	1	3	24	5	7	39	17
Last Chance Canyon/Rapids	152.0-153.0	3	2	1	3	24	5	7	39	17
Range Creek Rapids	150.5-152.0	3	2	1	3	24	5	7	39	17
Little Big Horn Mesa	143.0-144.0	3	2	1	3	24	5	7	39	17
Price River confluence	137.0-138.5	3	2	1	3	24	5	7	39	17
Downstream of Price River	136.5-137.5	3	2	1	3	24	5	7	39	17
Short Canyon/Rapids	131.5-132.5	3	2	1	3	24	5	7	39	17
Willow Bend-Tusher Rapid	128.5-129.5	1	2	1	1	24	5	14	44	16
Upriver of Green River Utah	121.5-125.5	1	2	1	1	24	5	7	37	18
Little Grand Wash	114.5-115.5	3	2	1	3	24	5	7	39	17
Downstream of Grand Wash	113.0-114.0	3	2	1	3	24	5	7	39	17
Pivemile Wash/Little Valley	111.0-112.0	3	2	1	3	24	5	7	39	17
Downstream of Ninemile Wash	109.0-111.0	3	2	1	3	24	5	7	39	17
Anvil Bottom	101.5-102.5	3	2	1	3	24	5	7	39	17
Upstream of San Rafael River	98.0-99.0	3	2	1	3	24	5	7	39	17
Sand Rafael River confluence	96.5-97.5	3	2	1	3	24	5	7	39	17
White and Red Wash	95.0-96.5	3	2	1	3	24	5	7	39	17
Between Red Wash-Bull Bottom	94.0-95.5	3	2	1	3	24	5	7	39	17
Bull Bottom-Labyrinth Canyon	92.0-93.0	3	2	1	3	24	5	7	39	17
Labyrinth Canyon confluence	91.5-92.5	3	2	1	3	24	5	7	39	17
Three Canyon/Trin-Alcove Bend	88.5-91.5	3	2	1	3	24	5	7	39	17
Junes Bottom	87.0-88.0	3	2	1	3	24	5	7	39	17
Junes Bottom-Bull Hollow	85.0-86.0	3	2	1	3	24	5	7	39	17
Bull Hollow	84.0-85.0	3	2	1	3	24	5	7	39	17
Bull Hollow-Tennille Canyon	81.0-83.0	3	2	1	3	24	5	7	39	17
Tennille Canyon/Bottom	79.0-82.0	3	2	1	3	24	5	7	39	17
Key Springs Canyon	75.0-78.5	3	2	1	3	24	5	7	39	17
Key Joe Canyon	74.5-76.0	3	2	1	3	24	5	7	39	17
Key Joe Canyon-Spring Canyon	74.0-75.0	3	2	1	3	24	5	7	39	17
Spring Canyon Point	69.5-74.0	3	2	1	3	24	5	7	39	17
Bowknot Bend	62.0-70.0	3	2	1	3	24	5	7	39	17
Twomile Canyon-Deadman Point	60.0-62.0	3	2	1	3	24	5	7	39	17
Deadman Point-Horseshoe Canyon	59.0-60.0	3	2	1	3	24	5	7	39	17
Downstream of Horseshoe Canyon	58.0-59.0	3	2	1	3	24	5	7	39	17
Cottonwood Bottom	55.0-56.5	3	2	1	3	24	5	7	39	17
Cottonwood-Mineral Bottoms	54.0-55.0	3	2	1	3	24	5	7	39	17
Mineral-Tidwell Bottoms	53.0-54.0	3	2	1	3	24	5	7	39	17
Tidwell Bottom	51.0-52.5	3	2	1	3	24	5	7	39	17
Horsechief Bottom	50.0-51.5	3	2	1	3	24	5	7	39	17
Woodruff Bottom	49.0-50.0	3	2	1	3	24	5	7	39	17
Point Bottom	47.5-49.5	3	2	1	3	24	5	7	39	17
Saddle Horse Bottom	45.5-47.5	3	2	1	3	24	5	7	39	17
Horsechief Canyon	45.0-46.0	3	2	1	3	24	5	7	39	17
Horsechief-Upheaval Bottom	44.0-45.5	3	2	1	3	24	5	7	39	17
Upheaval Bottom/Canyon	43.5-44.5	3	2	1	3	24	5	7	39	17
Hardscrabble Bottom	42.5-43.5	3	2	1	3	24	5	7	39	17
Hardscrabble-Port Bottoms	38.5-42.5	3	2	1	3	24	5	7	39	17

Table F.1. Continued.

Site Description	River Mile	Sub-Criteria Raw Score Values*			Sub-Criteria Raw Score Values multiplied By Average Weighted Point Values†			Total Score‡	Rank⁴		
		Status of land ownership	Proximity bar to river	June Potential of sites connection network	Status of land ownership	Proximity bar to river	June Potential of sites connection network				
<u>Green River</u>											
Potato Bottom	35.5-38.0	3	2	1	1	3	24	5	7	39	17
Potato-Beaver Bottoms	34.5-35.5	3	2	1	1	3	24	5	7	39	17
Beaver-Queen Ann Bottoms	33.5-34.5	3	2	1	1	3	24	5	7	39	17
Queen Ann Bottom	33.0-34.0	3	2	1	1	3	24	5	7	39	17
Queen Ann-Anderson Bottoms	32.0-33.0	3	2	1	1	3	24	5	7	39	17
Anderson Bottom	30.0-31.5	3	2	1	1	3	24	5	7	39	17
Unknown and Valentine Bottoms	29.0-32.0	3	2	1	1	3	24	5	7	39	17
Valentine Bottom	27.0-29.0	3	2	1	1	3	24	5	7	39	17
Stillwater Canyon-Sphinx	26.0-28.5	3	2	1	1	3	24	5	7	39	17
Downstream Valentine Bottom	25.5-26.5	3	2	1	1	3	24	5	7	39	17
Valentine-Tuxedo Bottoms	24.0-25.5	3	2	1	1	3	24	5	7	39	17
Tuxedo Bottom-Turks head	19.5-24.0	3	2	1	1	3	24	5	7	39	17
Deadhorse Canyon confluence	18.5-20.0	3	2	1	1	3	24	5	7	39	17
Downstream Deadhorse Canyon	17.0-19.0	3	2	1	1	3	24	5	7	39	17
Deadhorse-Horse Canyons	16.0-17.0	3	2	1	1	3	24	5	7	39	17
Horse Canyon	13.5-14.0	3	2	1	1	3	24	5	7	39	17
Downstream of Horse Canyon	12.5-13.5	3	2	1	1	3	24	5	7	39	17
Horse-Jasper Canyons	10.5-12.5	3	2	1	1	3	24	5	7	39	17
Jasper Canyon	8.0-10.5	3	2	1	1	3	24	5	7	39	17
Jasper Canyon-Short Canyon	5.0-9.0	3	2	1	1	3	24	5	7	39	17
Short Canyon-Colorado River	2.0-3.5	3	2	1	1	3	24	5	7	39	17
Upstream of Colorado River	1.0-2.0	3	2	1	1	3	24	5	7	39	17
Near Colorado River confluence	0.5-1.5	3	2	1	1	3	24	5	7	39	17
Colorado River confluence	0.0-0.5	3	2	1	1	3	24	5	7	39	17
<u>Yampa River</u>											
Upstream of Snake River confluence	48.0-48.5	1	1	3	1	1	12	15	7	35	19
<u>White River</u>											
Oxbow downstream Highway 64 bridge	91.0-93.0	1	2	3	1	1	24	15	7	47	15
White/Green river confluence	0.0-2.5	1	3	1	3	1	36	5	21	63	10

Table F.2. Selection criteria and associated sub-criteria raw score values multiplied by the average weighted point values to produce the total score and rank for each bottomland site in the Colorado and Gunnison rivers, Colorado and Utah, 1993. Top 13 ranked bottomland sites denoted by ‡.

Site Description	River Mile	Sub-Criteria Raw Score Values*						Sub-Criteria Raw Score Values Multiplied by Average Weighted Point Values ^b						Total Score ^c	Rank ^d
		Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites	Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites						
Colorado River															
Rifle West	238.0-241.1	1	1	1	2	1	1	1	4	12	28	25			
Rifle I-70 West Interchange	236.5-238.0	1	1	1	2	1	1	1	4	12	28	25			
CDOW/Gentry Property	234.2-235.8	2	1	1	2	2	1	1	4	12	29	24			
Trash-can Pond	230.2-230.8	1	1	3	2	1	1	1	12	12	36	19			
Mahaffey/Lemon	228.2-229.8	1	1	3	2	1	1	1	12	12	36	19			
Mahaffey's	228.5-229.5	1	1	3	2	1	1	1	12	12	36	19			
Hoaglund's	227.1-227.7	1	1	3	2	1	1	1	12	12	36	19			
Dere/Ortiz	227.5-228.5	1	1	3	2	1	1	1	12	12	36	19			
Parachute East	223.1-225.4	1	1	3	2	1	1	1	12	12	36	19			
Knight's	224.5-225.4	1	1	3	2	1	1	1	12	12	36	19			
Parachute Bridge	223.0-223.4	1	1	3	2	1	1	1	12	12	36	19			
No-name	227.7-223.1	1	1	1	2	1	1	1	4	12	28	25			
No-name	221.1-222.9	1	1	1	2	1	1	1	4	12	28	25			
*Battlement Mesa	220.7-221.7	2	2	3	2	2	2	2	12	12	48	10			
*EXXON	218.9-220.1	1	2	3	2	1	1	1	12	12	47	11			
Doyle Property	218.4-219.5	1	1	1	2	1	1	1	4	12	28	25			
No-name	217.2-218.0	1	1	1	2	1	1	1	4	12	28	25			
Una North	216.4-217.4	1	1	1	2	1	1	1	4	12	28	25			
*Wallace Creek Island	215.9-216.5	1	2	3	2	1	1	1	12	12	47	11			
*Stoddard Property	209.4-210.1	1	3	1	3	1	1	1	4	18	56	7			
*Debeque I-70 Slough	209.6-211.4	2	3	3	3	2	2	2	12	18	65	4			
*Latham's	207.9-208.8	1	2	3	3	2	2	2	12	18	53	9			
No-name	206.2-207.5	1	2	3	3	1	1	1	12	18	53	9			

* Each selection criteria is further sub-divided into sub-criteria and given a raw score value (see Methods section).

^b Equal to the sub-criteria raw score value multiplied by the average weighted point value (e.g., for Walter Walker SWA this equals 1 X 2=2 for the status of land ownership criteria, 5 X 11=55 for the proximity to adult RZ captures criteria, 4 X 3=12 for the June connection to the river criteria, and 6 X 3=18 for the potential network of sites criteria).

^c Equal to the sum of the sub-criteria raw score values multiplied by the average weighted point values (e.g., for Walter Walker SWA this equals 2+55+12+18=87).

^d The total score was sorted from the lowest to highest ordinal with the highest total score receiving the lowest ordinal rank and lowest total score the highest ordinal rank. The lowest ordinal rank was the top priority site and the highest ordinal rank the bottom priority site.

Table F.2. Continued.

Site Description	River Mile	Sub-Criteria Raw Score Values ^a					Sub-Criteria Raw Score Values Multiplied by Average Weighted Point Values ^b					Total Score ^c	Rank ^d
		Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites	Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites				
Colorado River													
*Etter's	204.7-206.6	1	2	3	3		1	22	12	18	53	9	
*"No-name"	202.3-204.2	1	2	3	2		1	22	12	12	47	11	
*"No-name"	201.9-202.4	1	2	3	2		1	22	12	12	47	11	
Long Point Bottom	198.0-198.9	3	1	3	1		1	11	12	6	30	23	
Beavertail	194.3-195.6	3	1	1	1		3	11	4	6	24	26	
Island Acres	191.1-192.1	2	1	1	1		2	11	4	6	23	27	
Cameo	190.0-191.0	3	1	3	1		1	11	12	6	30	23	
Palisade	184.2-185.0	2	2	1	2		1	22	4	12	39	16	
*"No-name"	183.3-184.2	1	2	3	2		1	22	12	12	47	12	
*Labor Camp	182.9-183.6	2	3	3	2		2	33	12	12	59	5	
"No-name"	181.6-182.6	1	2	1	2		1	22	4	12	39	16	
*"No-name"	181.7-182.2	1	2	3	2		1	22	12	12	47	11	
"No-name"	180.4-181.5	1	2	1	2		1	22	4	12	39	16	
*Clifton Water Treatment	179.1-181.1	1	2	1	3		1	22	4	18	45	13	
*Clifton Pond	177.7-178.2	2	4	1	3		2	44	4	18	68	3	
Com Lake	176.6-177.7	2	1	1	2		2	11	4	12	29	24	
*Humphrey's	175.3-176.6	2	2	3	3		1	22	12	18	53	9	
*Griffith's	174.1-176.5	1	2	3	3		1	22	12	18	53	9	
*"Hotspot Junction" (30-29 Road)	173.9-175.1	2	2	1	3		2	22	4	18	46	12	
"No-name"	174.4-173.8	1	1	3	2		1	11	12	12	36	19	
"No-name"	173.0-173.6	1	1	1	2		1	11	4	12	28	25	
Mill Tailing Site	172.5-173.4	1	1	3	2		1	11	12	12	36	19	
Watson Island	171.0-172.1	2	1	3	2		2	11	12	12	37	18	
"No-Name"	171.3-171.8	1	2	3	1		1	22	12	6	41	15	
"No-name"	170.4-171.0	1	2	3	1		1	22	12	6	41	15	
Connected Lakes Area	168.7-170.3	2	2	1	1		2	22	4	6	34	20	

^a Each selection criteria is further sub-divided into sub-criteria and given a raw score value (see Methods section).

^b Equal to the sub-criteria raw score value multiplied by the average weighted point value (e.g., for Walter Walker SWA this equals 1 X 2=2 for the status of land ownership criteria, 5 X 11=55 for the proximity to adult RZ captures criteria, 4 X 3=12 for the June connection to the river criteria, and 6 X 3=18 for the potential network of sites criteria).

^c Equal to the sum of the sub-criteria raw score values multiplied by the average weighted point values (e.g., for Walter Walker SWA this equals 2+55+12+18=87).

^d The total score was sorted from the lowest to highest ordinal with the highest total score receiving the lowest ordinal rank and lowest total score the highest ordinal rank. The lowest ordinal rank was the top priority site and the highest ordinal rank the bottom priority site.

Table F.2. Continued.

Site Description	River Mile	Sub-Criteria Raw Score Values ^a					Sub-Criteria Raw Score Values Multiplied by Average Weighted Point Values ^b					Total Score ^c	Rank ^d
		Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites	Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites				
Colorado River													
Walter Walker South	164.4-166.0	1	2	3	2	2	22	12	12	48	10		
No-name ^e	166.4-169.6	1	2	3	2	2	22	12	12	48	10		
Appleton Drain East	165.1-166.4	1	2	1	2	1	22	4	12	39	16		
Walter Walker SWA	162.7-165.1	2	5	3	3	2	55	12	18	87	1		
Panorama	163.1-163.6	1	3	3	2	1	33	12	12	58	6		
No-name ^e	161.0-162.7	1	2	3	2	1	22	12	12	47	11		
DuPont Island	161.0-162.0	1	1	3	2	1	11	12	12	36	19		
DuPont's	159.1-161.9	1	1	3	2	1	11	14	12	38	17		
Paul Smith's	158.0-159.1	1	1	3	2	1	11	12	12	36	19		
Fruit 340 Bridge	157.5-158.3	1	1	3	2	1	11	12	12	36	19		
Snook's Bottom	155.9-157.1	3	1	3	2	3	11	12	12	38	17		
Horsethief SWA	151.9-154.7	2	1	3	2	2	11	12	12	37	18		
Spann's	151.4-152.4	1	1	1	1	1	11	4	6	22	28		
No-name ^e	137.2-137.1	3	1	1	1	3	11	4	6	24	26		
Crow Bottom	143.9-146.5	3	1	1	1	3	11	4	6	24	26		
No-name ^e	145.5-146.8	1	1	1	1	1	11	4	6	21	29		
Vulture Bottom	139.7-142.1	3	1	1	1	3	11	4	6	24	26		
No-name ^e Island	137.2-138.0	3	1	1	1	3	11	4	6	24	26		
Black Rocks	136.7-137.1	3	1	3	1	3	11	12	6	32	21		
Knowles Canyon	133.6-135.0	3	1	3	6	3	11	12	6	24	26		
Jouffas Bottom	129.8-133.9	3	1	1	1	3	11	4	6	24	26		
Wildass Canyon Ranch	127.3-131.3	1	1	3	1	3	11	12	6	30	23		
No-name ^e	125.8-126.8	3	1	3	1	3	11	12	6	32	21		
Elizondo Ranches	126.5-127.8	1	1	1	1	1	11	4	6	22	28		
Westwater Wash	124.8-126.0	1	1	3	1	1	11	12	6	30	23		
Cisco Landing Area	107.6-111.6	3	1	3	2	3	11	12	12	38	17		

^a Each selection criteria is further sub-divided into sub-criteria and given a raw score value (see Methods section).

^b Equal to the sub-criteria raw score value multiplied by the average weighted point value (e.g., for Walter Walker SWA this equals 1 X 2=2 for the status of land ownership criteria, 5 X 11=55 for the proximity to adult RZ captures criteria, 4 X 3=12 for the June connection to the river criteria, and 6 X 3=18 for the potential network of sites criteria).

^c Equal to the sum of the sub-criteria raw score values multiplied by the average weighted point values (e.g., for Walter Walker SWA this equals 2+55+12+18=87).

^d The total score was sorted from the lowest to highest ordinal with the highest total score receiving the lowest ordinal rank and lowest total score the highest ordinal rank. The lowest ordinal rank was the top priority site and the highest ordinal rank the bottom priority site.

Table F.2. Continued.

Site Description	River Mile	Sub-Criteria Raw Score Values ^a					Sub-Criteria Raw Score Values Multiplied by Average Weighted Point Values ^b					Total Score ^c	Rank ^d
		Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites	Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites				
Colorado River													
Fish Ford Area	103.0-105.9	3	1	3	2	3	11	12	12	38	17		
McGraw/Hotel Bottom	98.1-101.0	2	1	3	2	2	11	12	12	31	22		
White Ranch	77.5-78.4	1	1	1	1	1	11	4	6	22	28		
Courthouse Wash	63.8	1	1	3	1	1	11	12	6	30	23		
Moab Slough	61.5-64.0	2	1	3	1	2	11	12	6	31	22		
Kane Spring	58.2	3	1	3	1	3	11	12	6	32	21		
Billboard (Lake Bottom)	59.9-52.5	3	1	1	1	3	11	4	6	24	26		
Jackson Bottom	47.4-49.0	1	1	3	1	1	11	12	6	30	23		
"No-name"	44.0-44.7	3	1	1	1	3	11	4	6	24	26		
"No-name"	42.8-43.7	3	1	3	1	3	11	12	6	32	21		
"No-name"	41.1-42.1	3	1	3	1	3	11	12	6	32	21		
"No-name"	40.2-41.3	3	1	3	1	3	11	12	6	32	21		
"No-name"	39.5-40.6	3	1	3	1	3	11	12	6	32	21		
"No-name" Canyon Mouth	38.8	3	1	3	1	3	11	12	6	32	21		
"No-name"	38.6-39.2	3	1	1	1	3	11	4	6	32	21		
Goose Neck	33.2-35.6	3	1	3	1	3	11	12	6	32	21		
Shafer Canyon Mouth	34.8	3	1	3	1	3	11	12	6	32	21		
"No-name" Canyon Mouth	32.1	3	1	3	1	3	11	12	6	32	21		
"No-name" Canyon Mouth	31.6	3	1	3	1	3	11	12	6	32	21		
"No-name"	30.8-31.9	3	1	1	1	3	11	4	6	32	21		
Little Bridge Canyon	30.0	3	1	3	1	3	11	12	6	32	21		
Lockhart Canyon	26.5	3	1	3	1	3	11	12	6	32	21		
"No-name"	25.3-26.5	3	1	1	1	3	11	4	6	24	26		
Lathrop Canyon	23.5	3	1	3	1	3	11	12	6	32	21		
Buck Canyon	22.7	3	1	3	1	3	11	12	6	32	21		
Gooseberry Canyon	21.7	3	1	3	1	3	11	12	6	32	21		

^a Each selection criteria is further sub-divided into sub-criteria and given a raw score value (see Methods section).

^b Equal to the sub-criteria raw score value multiplied by the average weighted point value (e.g., for Walter Walker SWA this equals $1 \times 2 = 2$ for the status of land ownership criteria, $5 \times 11 = 55$ for the proximity to adult RZ captures criteria, $4 \times 3 = 12$ for the June connection to the river criteria, and $6 \times 3 = 18$ for the potential network of sites criteria).

^c Equal to the sum of the sub-criteria raw score values multiplied by the average weighted point values (e.g., for Walter Walker SWA this equals $2 + 55 + 12 + 18 = 87$).

^d The total score was sorted from the lowest to highest ordinal with the highest total score receiving the lowest ordinal rank and lowest total score the highest ordinal rank. The lowest ordinal rank was the top priority site and the highest ordinal rank the bottom priority site.

Table F.2. Continued.

Site Description	River Mile	Sub-Criteria Raw Score Values ^a					Sub-Criteria Raw Score Values Multiplied by Average Weighted Point Values ^b					Total Score ^c	Rank ^d
		Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites	Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites				
<u>Colorado River</u>													
Dogleg Canyon	21.2	3	1	3	1	3	11	12	6	32	21		
Sheep Bottom	17.5-18.7	3	1	1	1	3	11	4	6	24	26		
Indian Creek Canyon Mouth	16.5	3	1	3	1	3	11	12	6	32	21		
Monument Creek Mouth	15.3	3	1	1	1	1	11	4	6	24	26		
"No-name" Canyon Mouth	11.9	3	1	1	1	1	11	4	6	24	26		
"No-name" Canyon Mouth	10.1	3	1	3	1	3	11	12	6	32	21		
Salt Creek Canyon Mouth	3.4	3	1	3	1	3	11	12	6	32	21		
Elephant Creek Canyon Mouth	3.0	3	1	3	1	3	11	12	6	32	21		
<u>Gunnison River</u>													
"No-name"	74.3-74.7	1	1	1	1	1	11	4	6	22	28		
Lawhead Gulch Bottom	70.1-71.6	1	1	3	1	1	1	12	6	30	23		
Ferganchick's	69.5-70.5	1	1	3	1	1	11	12	6	30	23		
"No-name"	68.3-69.4	1	1	2	1	1	11	12	6	30	23		
"No-name"	67.5-68.2	1	1	1	1	1	11	4	6	22	28		
Austin County Bridge	65.7-66.4	1	1	1	1	1	11	4	6	22	28		
Austin Hwy 92 Bridge	65.0-65.5	1	1	3	1	1	11	12	6	30	23		
"No-name"	63.3-65.3	1	1	3	1	1	11	12	6	30	23		
"No-name"	63.0-64.6	1	1	1	1	1	11	4	6	22	28		
Colorado Hwy 65 Bridge	62.6-63.2	1	1	3	2	1	11	12	12	36	19		
Tongue Creek	61.5-62.5	1	1	3	2	1	11	12	12	36	19		
Hutchin's	60.1-60.8	1	1	3	2	1	11	12	12	36	19		

^a Each selection criteria is further sub-divided into sub-criteria and given a raw score value (see Methods section).

^b Equal to the sub-criteria raw score value multiplied by the average weighted point value (e.g., for Walter Walker SWA this equals 1 X 2=2 for the status of land ownership criteria, 5 X 11=55 for the proximity to adult RZ captures criteria, 4 X 3=12 for the June connection to the river criteria, and 6 X 3=18 for the potential network of sites criteria).

^c Equal to the sum of the sub-criteria raw score values multiplied by the average weighted point values (e.g., for Walter Walker SWA this equals 2+55+12+18=87).

^d The total score was sorted from the lowest to highest ordinal with the highest total score receiving the lowest ordinal rank and lowest total score the highest ordinal rank. The lowest ordinal rank was the top priority site and the highest ordinal rank the bottom priority site.

Table F.2. Continued.

Site Description	River Mile	Sub-Criteria Raw Score Values ^a					Sub-Criteria Raw Score Values Multiplied by Average Weighted Point Values ^b					Total Score ^c	Rank ^d
		Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites	Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites				
Gunnison River													
North Delta	57.6-59.5	1	1	3	2	1	11	12	12	36	19		
South Delta	57.6-58.8	1	1	1	2	1	11	4	12	28	25		
Confluence Park	56.7-57.7	2	1	3	3	1	11	12	18	43	14		
Uncompahgre R. Confluence	56.3-56.5	2	1	3	2	1	11	12	12	36	19		
Delta City Sewage Plant	55.6-56.5	2	1	3	2	2	11	12	12	37	18		
"No-name"	54.6-55.5	1	1	3	2	1	11	12	12	36	19		
"No-name"	54.2-55.1	1	1	3	2	1	11	12	12	36	19		
"No-name"	53.1-54.1	1	1	1	2	1	11	4	12	28	25		
Johnson Boy's Slough	53.2-54.2	1	4	3	3	1	44	12	18	75	2		
Escalante SWA North	50.8-52.9	2	2	3	3	2	22	12	10	54	8		
Escalante SWA South	50.2-52.4	2	2	3	3	2	22	12	18	54	8		
"Blue Duck" Bottom	49.5-50.4	2	1	3	1	2	11	12	6	21	22		
"No-name"	49.4-49.7	2	1	3	1	1	11	12	6	30	23		
"No-name"	48.6-49.3	1	1	3	1	1	11	12	6	30	23		
"No-name"	46.6-47.5	1	1	3	1	1	11	12	6	30	23		
Escalante Ranches	41.7-42.8	1	1	3	1	1	11	12	6	30	23		
"No-name"	39.9-41.0	3	1	3	1	3	11	12	6	32	21		
Dominguez	36.4-36.9	1	1	3	1	1	11	12	6	30	23		
Broughton	35.6-36.0	1	1	3	1	1	11	12	6	30	23		
McKendrick's	34.8-36.3	1	1	3	1	1	11	12	6	30	23		
Peepie's South (VIP Camp)	34.0-35.2	1	1	3	1	1	11	12	6	30	23		
Peepie's Island	34.6-34.9	1	1	3	1	1	11	12	6	30	23		
Sand Flat	27.3-28.7	1	1	1	1	1	11	4	6	30	23		

* Each selection criteria is further sub-divided into sub-criteria and given a raw score value (see Methods section).

^b Equal to the sub-criteria raw score value multiplied by the average weighted point value (e.g., for Walter Walker SWA this equals 1 X 2=2 for the status of land ownership criteria, 5 X 11=55 for the proximity to adult RZ captures criteria, 4 X 3=12 for the June connection to the river criteria, and 6 X 3=18 for the potential network of sites criteria).

^c Equal to the sum of the sub-criteria raw score values multiplied by the average weighted point values (e.g., for Walter Walker SWA this equals 2+55+12+18=87).

^d The total score was sorted from the lowest to highest ordinal with the highest total score receiving the lowest ordinal rank and lowest total score the highest ordinal rank. The lowest ordinal rank was the top priority site and the highest ordinal rank the bottom priority site.

Table F.2. Continued.

Site Description	River Mile	Sub-Criteria Raw Score Values ^a					Sub-Criteria Raw Score Values Multiplied by Average Weighted Point Values ^b					Total Score ^c	Rank ^d
		Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites	Status of land ownership	Proximity to adult RZ captures	June connection to river	Potential network of sites				
Gunnison River													
Tunnel Point	26.3-26.9	3	1	3	1	3	11	12	6	32	21		
Dads Flat	24.7-26.3	1	1	3	1	1	11	12	6	30	23		
Deer Run	23.0-23.6	3	1	3	1	3	11	12	6	32	21		
"No-name"	22.1-22.7	3	11	3	1	3	11	12	6	32	21		
"No-name"	21.3-21.8	3	1	3	1	3	11	12	6	24	26		
"No-name"	18.4-19.2	3	1	3	1	3	11	12	6	32	21		
Kannah Creek	18.0-18.6	1	1	3	1	1	11	12	6	30	23		
Whitewater Bliding Materials	13.3-16.0	1	1	3	1	1	11	12	6	30	23		
Duck Blind	12.7-13.8	1	1	3	1	1	11	12	6	30	23		
Bangs Canyon	11.7-13.0	1	1	1	1	1	11	4	6	24	26		
Schroeder's	4.5-6.1	1	1	3	1	1	11	12	6	30	23		
Mule Farm	3.1-3.9	1	1	1	1	1	11	4	6	24	26		
Redlands	2.2-3.0	1	1	1	1	1	11	4	6	24	26		

* Each selection criteria is further sub-divided into sub-criteria and given a raw score value (see Methods section).

^b Equal to the sub-criteria raw score value multiplied by the average weighted point value (e.g., for Walter Walker SWA this equals 1 X 2=2 for the status of land ownership criteria, 5 X 11=55 for the proximity to adult RZ captures criteria, 4 X 3=12 for the June connection to the river criteria, and 6 X 3=18 for the potential network of sites criteria).

^c Equal to the sum of the sub-criteria raw score values multiplied by the average weighted point values (e.g., for Walter Walker SWA this equals 2+55+12+18=87).

^d The total score was sorted from the lowest to highest ordinal with the highest total score receiving the lowest ordinal rank and lowest total score the highest ordinal rank. The lowest ordinal rank was the top priority site and the highest ordinal rank the bottom priority site.

APPENDIX G
Land ownership acreage for the
Colorado and Gunnison rivers

Table G.1. Land ownership status by area (hectares) and percentage for 158 bottomland sites adjacent to the Colorado and Gunnison rivers (see Figures 25 and 26).

River Stream Reach (River Mile)	Land Ownership Status												
	Public						Public/Private ^b						
	Private	%	State	%	City	%	Federal ^a	%	State/ Federal	%	Public/ Private ^b	%	Total
Colorado													
Rifle-Debeque (240-203)	951	93	67	7	0	0	0	0	0	0	0	0	1,018
Palisade-Loma (185-152)	1,194	64	0	0	63	3	0	0	181	10	426	23	1,864
Loma-Stateline (152-132)	40	14	0	0	0	0	93	32	0	0	158	54	291
Stateline-Moab (132-63)	387	25	0	0	0	0	100	7	0	0	1,034	68	1,521
Moab-Green River Confluence (63-0.0)	0	0	22	7	0	0	265	83	11	3	23	7	321
TOTALS	2,572	51	89	2	63	1	458	9	192	4	1,641	33	5,015
Gunnison													
North Fork-Austin (75-67)	88	100	0	0	0	0	0	0	0	0	0	0	88
Austin-Railroad Bridge (67-50)	419	64	193	30	40	6	0	0	0	0	0	0	652
Railroad Bridge- Colorado R. Confluence (50-0.7)	392	72	11	2	0	0	42	8	36	7	66	12	547
TOTALS	899	70	204	16	40	3	42	3	36	3	66	5	1,287
^a Includes lands owned or controlled by Bureau of Land Management, Bureau of Reclamation or the National Park Service.													
^b Includes both Federal and State lands interspersed with private ownership.													

^a Includes lands owned or controlled by Bureau of Land Management, Bureau of Reclamation or the National Park Service.

^b Includes both Federal and State lands interspersed with private ownership.

APPENDIX H

Nine initial criteria and sub-criteria considered for
scoring and ranking bottomland sites

Table H.1. The initial nine criteria and sub-criteria considered for selection criteria for scoring and ranking bottomland sites in the Upper Colorado River Basin.

=====	
1. Area	
≥ 75 acres (Total Potential [historical] Inundated Area)	
< 75 acres (" " " " ")	
2. Ownership	
Public	
Private	
3. Proximity to Fish Use (Historical and Current)	
1- 25 River miles	
26- 50 " "	
51-100 " "	
>100 " "	
4. Water Permanency/Duration (June-September)	
Yes/No	
5. Present Hydrological Connection to River	
Yes/No	
6. Wetland Feature	
Natural	
Man-made	
7. Suitability or Appropriate for Fish/Water Control Structures	
Yes/No	
8. Diked/Riprap; Other Obstruction	
Yes/No	
9. Potential for Network of Sites (within 25 river miles of known historical or current fish use: RZ, CS)	
Number of Sites within 25 miles to comprise a network of sites	
0-1 Sites	
2-3 "	
≥4 "	
=====	

APPENDIX I
Mean daily flow exceedences




Figure I.1. Percent of time flow (cfs [cms]) is exceeded in the Green River before (1947-1966) and after (1966-1993) Flaming Gorge Dam.





Figure I.2. Mean daily flow exceedences for May (top) and June (bottom) for the Colorado River at the Cameo USGS stream gauge during the post-water development period, 1952-1993.





Figure I.3. Mean daily flow exceedences for May (top) and June (bottom) for the Colorado River at the Stateline USGS stream gauge during the post-water development period, 1952-1993.





Figure I.4. Mean daily flow exceedences for May (top) and June (bottom) for Plateau Creek (near the confluence with the Colorado River) USGS stream gauge during the post-water development period, 1953-1993.





Figure I.5. Mean daily flow exceedences for May (top) and June (bottom) for the Gunnison River at the Whitewater USGS stream gauge during the post-water development period, 1967-1993.





Figure I.6. Mean daily flow exceedences for May (top) and June (bottom) for the Gunnison River at the Delta USGS stream gauge during the post-water development period, 1977-1993.

