

RECEIVED

JUN 1 1 1969

Winthrop College Library  
Documents Department

# PECOS NATIONAL MONUMENT NEW MEXICO

## ITS GEOLOGIC SETTING





# PECOS NATIONAL MONUMENT NEW MEXICO

## ITS GEOLOGIC SETTING

By ROSS B. JOHNSON

*A description of  
rocks and landscapes  
and their effect on man*



CONTRIBUTIONS TO GENERAL GEOLOGY  
GEOLOGICAL SURVEY BULLETIN 1271-E

UNITED STATES DEPARTMENT OF THE INTERIOR

WALTER J. HICKEL, *Secretary*

GEOLOGICAL SURVEY

William T. Pecora, *Director*

U.S. GOVERNMENT PRINTING OFFICE, WASHINGTON : 1969

---

For sale by the Superintendent of Documents, U.S. Government Printing Office  
Washington, D.C. 20402

## CONTENTS

---

	Page
Introduction -----	E1
Rocks near the monument -----	2
Geographic extent of the formations of Glorieta Mesa -----	7
Sources of rock fragments and environments of deposition -----	9
Relation of Glorieta Mesa to the mountains -----	9
Use of rock products by the Indians and Spanish priests -----	10
References cited -----	11

---

## PLATE

---

1. Geology and topography of Glorieta Mesa and  
Pecos National Monument, N. Mex ----- In pocket

---

## FIGURES

---

	Page
1. Map showing location of Glorieta Mesa and Pecos National Monument -----	E3
2. Photograph showing rock sequence on Glorieta Mesa at Cerro de Escobas -----	4
3. Columnar section of generalized geologic se- quence of rock units on Glorieta Mesa at Cerro de Escobas near Pecos National Monu- ment -----	5



CONTRIBUTIONS TO GENERAL GEOLOGY

---

PECOS NATIONAL MONUMENT  
NEW MEXICO  
ITS GEOLOGIC SETTING

By Ross B. Johnson

The scenographical arrangement of the surrounding country is remarkably picturesque; the view of Pecos, as it now lies, without the least addition, would form a beautiful picture, and more than a picture, for every cloud, every degree that the sun moves, gives such varied effects to the landscape, that one has a thousand pictures; but their effects are so fleeting, that although they last long enough to delight the spectator, it would yet perplex the artist to catch these changes. [Aber<sup>t</sup> 1848.]

### Introduction

The ruins of the pueblos and missions of Pecos lie on the east bank of Glorieta Creek near its junction with the Pecos River at the south end of the Sangre de Cristo Mountains in north-central New Mexico. Here the Pecos River and Glorieta Creek have formed a broad rolling valley in which the red adobe walls of the mission church stand as a striking monument to a historic past.

This is beautiful country; the bright hues of red rocks are complemented by the varied greens of the junipers, piñons, and ponderosa pines. Northward the Sangre de Cristo Mountains stretch for miles in a blue mist toward the Truchas Peaks and forests of the Pecos Wilderness. A few miles south of the ruins the steep high escarpment of Glorieta Mesa marks, in a general way, the southern termination of the Rocky Mountain System, which here is represented by the Sangre de Cristos.

The escarpment of Glorieta Mesa has been formed largely by the Pecos River and its tributaries eroding the soft sedimentary layers. The Pecos flows southward from the high mountains in the north, parallels the mesa escarpment for 15 miles, and breaches the mesa near San Jose. About  $1\frac{1}{2}$  miles southwest of the Pecos ruins at Cerro de Escobas is the highest point on Glorieta Mesa. It is the most conspicuous feature of the local landscape and rises to an elevation of 8,212 feet—1,270 feet above the ruins. The slope of the escarpment here is very steep, rising 6 feet in every 10 horizontal feet.

Along the north side of the Glorieta Mesa escarpment is a 30-mile-long natural pass around the south end of the Sangre de Cristos that extends from Cañoncito on the west to Starvation Peak on the east (fig. 1). The elevation of the pass is greater than 6,000 feet at all places, and it reaches its summit of 7,432 feet near the village of Glorieta near the west end of the pass. This pass has been used as a major travel route for more than 800 years by the Indians, Spanish, and Americans. The famous Santa Fe Trail passed through here and was superseded by the railroad, whose main-line tracks closely parallel the traces of the old wagon ruts. The modern four-lane divided highway, Interstate Highway I-25, carries high-speed automotive traffic through Glorieta Pass alongside the Santa Fe Railway. Glorieta Pass has been the locale of many important historical events, including the passage of Coronado's expedition in 1540-41; the construction of the two large mission churches at Pecos Pueblo; the capture and imprisonment of the men of the Texas Expedition in 1841; the passage of the American Army under General Kearny on its way to Santa Fe, Chihuahua, and California in 1846; and the Civil War Battle of Glorieta Pass in 1862.

## Rocks Near the Monument

The rocks of Glorieta Mesa and Pecos National Monument are all of sedimentary origin; that is, they consist mostly of fragments or grains of older rocks and were deposited by streams, in the oceans, or even in lakes. The fragments that make up the rocks of Glorieta Mesa were deposited 200 to 300 million years ago in nearly horizontal layers; these layers lie in almost the same position today.

As one can see from the photograph of Glorieta Mesa at Cerro de Escobas (fig. 2), the nearly horizontal layers of rocks can be




Figure 1.—Glorieta Mesa and Pecos National Monument.









Figure 2.—Rock sequence on Glorieta Mesa at Cerro de Escobas; six rock units viewed southwest from Pecos National Monument.

grouped readily into several units of similar rocks. The six units of rocks that have been separately named by geologists are shown in figure 2. Such units are called formations when the rocks can be separated in this way and traced in the manner shown on the geologic map (pl. 1). A sedimentary formation may be mostly one kind of rock fragment, or alternating layers of at least two kinds of rock fragments either deposited under the same environment or having similar physical characteristics, such as color and thickness of the layers.

The six units exposed on the escarpment of Glorieta Mesa have been named and mapped throughout New Mexico and one even into south-central Colorado. Each name was formally assigned to a rock unit in a scientific publication by the geologist first describing and mapping the unit. Each name is reserved for its particular unit, and it may not be used again for any other rock unit in the United States. The names of these six formations have been used and recognized in geologic literature for many years. A formation is named usually for a geographic feature near which it was first described or mapped. These formations were named for two mountain ranges, three mesas, and a city. They are all familiar names in New Mexico and without exception show the influence of the Southwest's Spanish heritage on the names of its geographic features.

The youngest formation on Glorieta Mesa is the Santa Rosa Sandstone of Late Triassic age (190–210 million years old; fig. 3) that caps Cerro de Escobas. In this area the upper part of the formation has been removed by recent erosion, and only 50 feet or so remains. At other places in New Mexico the Santa Rosa may reach a thickness of 450 feet. As its name implies, the formation consists of layers of rocks made up mostly of sand grains, that is, sandstone. The sand grains are clean clear quartz cemented together by reddish-brown clay that makes the sandstone a light tan. The Santa Rosa also contains smaller amounts of layered red shale, which consists of extremely small particles of clay and

SYSTEM	AGE (my*)	THICKNESS (FEET)	FORMATION	GEOLOGIC SECTION	LITHOLOGIC DESCRIPTION
TRIASSIC	190-210	50+	Santa Rosa Sandstone		Light-gray and tan sandstone beds UNCONFORMITY
PERMIAN	235-270	250-270	Bernal Formation		Purplish-red sandstone and siltstone beds. Some local beds of gypsum
			San Andres Limestone		Gray limestone beds
		15-20	Glorieta Sandstone		Light-gray sandstone beds
		125-135	Yeso Formation		Bright-red sandstone and siltstone with a few thin gray limestone beds
		380-400			Brownish-red and gray conglomerate, buff and red sandstone, red siltstone, red and greenish-gray shale, and gray limestone beds
500+	Sangre de Cristo Formation		Brownish-red and gray conglomerate, buff and red sandstone, red siltstone, red and greenish-gray shale, and gray limestone beds		
PENNSYLVANIAN	270-300				

\*Million years

Figure 3.—Generalized geologic sequence of rock units on Glorieta Mesa at Cerro de Escobas near Pecos National Monument.

other minerals; the red is due to oxidation, or rusting, of iron-bearing mineral fragments in the shale.

The Bernal Formation of Permian, probably Late Permian, age (about 235 million years old) underlies the Santa Rosa Formation. The units are not quite parallel to each other, and thus a relationship known as an unconformity exists between the two units. Rocks of Early Triassic age are missing on Glorieta Mesa, and it is deduced that Early Triassic time was an age when no sediments were deposited in northern New Mexico and when the older rocks of the Bernal Formation were slightly eroded. Here on Glorieta Mesa the Bernal consists of purplish-red sandstone and siltstone. Near the village of Bernal the formation contains some thin beds of gypsum.

Beneath, and older than, the rocks of the Bernal Formation are the gray limestone beds of the San Andres Limestone, also of Permian age. Here there is no unconformity between the two units, for deposition of sediments was continuous and uninterrupted from the San Andres into the Bernal. The limestones of the San Andres are dense and earthy and contain a few sandy layers.

The Glorieta Sandstone is the most conspicuous formation on Glorieta Mesa because of its contrasting light-gray to white color and because it caps the mesa at most places. The Glorieta is also of Permian age, and grades upward into the younger San Andres Limestone without interruption. The clean quartz sand grains of the Glorieta are cemented together so well that the rock is extremely hard and consequently strongly resists the chemical and physical attacks of weathering agents. Throughout its exposures in northern New Mexico, it stands out as a steep cliff face and usually forms the caprock of mesas.

Underlying the Glorieta Sandstone are the bright-red sandstone and siltstone layers of the Yeso Formation of Permian age. Although the change upward from the bright-red sandstone of the Yeso to the light-gray sandstone of the Glorieta is striking, sedimentation was continuous; and changes in thickness of layers and in kinds and sizes of fragments are gradual. The Yeso also contains some thin beds of red shale and a few local beds of limestone. In some areas it also contains beds of gypsum. In the vicinity of Glorieta Mesa, however, conditions were not suitable for the deposition of gypsum.

The Sangre de Cristo Formation underlies the Yeso Formation without a break in sedimentation. The Sangre de Cristo

Formation is the oldest formation exposed on Glorieta Mesa and is the oldest Permian formation in the region. The lower part is still older—of Pennsylvanian age. The upper part is about 270 million years old, but the lowermost part may be as old as 300 million years. The upper 100 feet or so of the Sangre de Cristo is very well exposed on the lower face of the Glorieta escarpment. However, in the broad valley formed by the Pecos River and Glorieta Creek, the lower four-fifths of the formation is covered at most places by a mantle of thick red soil derived from the weathering and decomposition of the formation itself. The formation consists of layers of varying thickness of brownish-red and gray conglomerate, buff and red sandstone, red siltstone, red and greenish-gray shale, and gray limestone beds.

Older rocks underlie those of the Sangre de Cristo Formation in the area of Pecos National Monument, but are not at the surface anywhere in the vicinity. These older rocks may be seen along the Pecos River north of the village of Pecos. They consist generally of limestone, shale, sandstone, and conglomerate of Pennsylvanian, Mississippian, and perhaps Devonian age (350–400 million years old) and ancient crystalline rocks of Precambrian age (more than 600 million years old).

## Geographic Extent of the Formations of Glorieta Mesa

The six formations as delineated and mapped on Glorieta Mesa extend over many thousands of square miles. They appear to be coextensive from the limited viewpoint at Pecos National Monument, but beyond that view they are not. Locally, throughout New Mexico and southern Colorado, several, one, or none of these formations may be present.

The Santa Rosa Sandstone was named for exposures near the city of Santa Rosa, N. Mex. The Santa Rosa crops out along the canyons of the Canadian and Pecos Rivers of northeastern and east-central New Mexico. At most places on Glorieta Mesa, the upper rocks of the formation have been eroded. However, the complete unit is present along the east flank of the Sangre de Cristo Mountains and extends at least as far north as Ocate Creek, 50 miles north of Las Vegas where it is covered by lava flows of much younger age.

Bernal Mesa east of the village of Bernal near Starvation Peak is the locality where the Bernal Formation was first named and

described. It is a local unit, crops out only on Glorieta Mesa, and caps low mesas a short distance south and southeast of Starvation Peak. It also extends along the east flank of the Sangre de Cristo Mountains, at least as far north as Ocate Creek. To the south, rocks equivalent to this unit are known by another name.

The San Andres Limestone was named for exposures in the San Andres Mountains of south-central New Mexico. Of all the rock units occurring on Glorieta Mesa, the San Andres probably covers the surface over more square miles than any other unit. The San Andres Limestone crops out in large separated areas across southern New Mexico into trans-Pecos Texas. The formation has also been mapped in the Zuni Mountains and the Lucero uplift west of Belen, N. Mex. North of Glorieta Mesa the limestone thins to a point where it can no longer be distinguished as a unit.

The name for the Glorieta Sandstone was derived from exposures on Glorieta Mesa just  $2\frac{1}{2}$  miles south of Cerro de Escobas. Because the Glorieta forms the main caprock of Glorieta Mesa, it is beautifully and conspicuously exposed for many miles; and thus the mesa very appropriately gives its name to the formation. The Glorieta Sandstone crops out in large separated areas that cover many thousands of square miles of New Mexico. It extends northward along the east flank of the Sangre de Cristo Mountains at least as far as Ocate Creek. The Glorieta Sandstone has been mapped in the Zuni Mountains of west-central New Mexico, the Lucero uplift west of Belen, and much of central New Mexico.

The Yeso Formation was named from Mesa del Yeso, 12 miles northeast of Socorro, N. Mex. A thick bed of gypsum, or yeso, crops out on the slopes of Mesa del Yeso, and therefore furnishes the name for both the geologic formation and the mesa. From Glorieta Mesa, rocks of the Yeso Formation extend southward and westward and form the surface rocks over large parts of west-central, central, and south-central New Mexico. North from Glorieta Mesa the sandstones and siltstones of the Yeso Formation grade into rocks typical of the Sangre de Cristo Formation and thus are included as part of the Sangre de Cristo Formation in northern New Mexico and southern Colorado.

Although the Sangre de Cristo Mountains occupy a large part of northern New Mexico, the Sangre de Cristo Formation was first named and described not in New Mexico but in Colorado. The name was first used in 1899 for rocks of this formation on the east slopes of the Sangre de Cristo Mountains northwest of the Spanish Peaks— 130 miles north of Pecos National Monument. The formation is appropriately named, for its outcrops are confined almost

entirely to the Sangre de Cristo Mountains of northeastern New Mexico and south-central Colorado.

## Sources of Rock Fragments and Environments of Deposition

Geologists can tell the environment in which the sedimentary rocks were deposited by studying various features in the rocks, such as fossils, kinds of fragments and minerals, shapes of the individual units, and depositional structures within the layers. The geologists sometimes can also tell the source of the fragments that make up the sedimentary rocks if they have a complete knowledge of the older rocks in the region.

From data such as these, geologists have concluded that the sands that make up the Santa Rosa Sandstone and the Glorieta Sandstone were deposited just offshore on the beaches of seas that were advancing over the land. The sand grains were derived from the winnowing action of waves on the sediments of the underlying formations—the Bernal and Yeso Formations.

The sediments that formed the rocks of the Bernal and Yeso Formations appear to be of shallow-water marine origin. Perhaps they were laid down on a broad, flat coastal plain. The San Andres Limestone is a marine limestone and probably was deposited during the wide advance of the Permian sea over the coastal plain after the deposition of the beach sands of the Glorieta Sandstone.

In contrast to these water-laid deposits, the Sangre de Cristo Formation consists mainly of terrestrial or land-laid sediments deposited by streams as alluvial fans, flood plains, deltas, and channel fills. Thin layers of limestone occur throughout the formation and contain fossils that indicate that the limestone layers are of both fresh-water and marine origin. Therefore, from time to time, lakes and ponds or ocean bays covered large parts of the region. Most of the fragments of the conglomerates, sandstones, siltstones, and shales of the Sangre de Cristo Formation are derived from the ancient Precambrian crystalline rocks that form the core of the high mountains a few miles to the north. From the unweathered appearance of the coarser fragments, it is evident that the source of these fragments was not very far away.

## Relation of Glorieta Mesa to the Mountains


The sedimentary rock layers of Glorieta Mesa slope gently south from the Sangre de Cristo Mountains. The slope is so gentle at



105°42'30"

R. 12 E

105°40'



105°42'30"

R. 12 E


INTERIOR—GEOLOGICAL SURVEY, WASHINGTON, D. C.—1969—G68396

105°40'


EXPLANATION




Santa Rosa Sandstone




Bernal Formation



San Andres Limestone



Glorieta Sandstone



Yeso Formation



Sangre de Cristo Formation

TRIASSIC

PERMIAN

PENNSYLVANIAN

Contact



APPROXIMATE MEAN DECLINATION, 1969

Base from U.S. Geological Survey,  
Pecos, 1961

SCALE 1:24 000

Geology by R. B. Johnson, 1967



CONTOUR INTERVAL 40 FEET  
DATUM IS MEAN SEA LEVEL

GEOLOGY AND TOPOGRAPHY OF GLORIETA MESA AND  
PECOS NATIONAL MONUMENT, NEW MEXICO

